

*The*  
American Journal  
of Medicine



December 1951





**FOR  
THE  
FIRST  
TIME**

## **aqueous natural vitamin A in capsules**

**AQUASOL A CAPSULES**  
is the first and only product to provide  
water-soluble natural vitamin A  
in capsules... and is made by the "oil-  
in-water" technique developed in  
the Research Laboratories of the U. S.  
Vitamin Corporation (U. S. Pat. 2,417,299).

# **AQUASOL A CAPSULES**

two potencies:

**25,000 U. S. P. Units**  
natural vitamin A per capsule  
... in water-soluble form

**50,000 U. S. P. Units**  
natural vitamin A per capsule  
... in water-soluble form



Bottles of 100, 500 and 1000 capsules

Samples upon request

**u. s. vitamin corporation**

casimir funk laboratories, inc. (affiliate)  
250 east 43rd st. • new york 17, n. y.

### **advantages:**

**up to 500%  
greater absorption  
80% less excretion  
85% higher liver storage**

### **indications:**

for more rapid,  
more effective therapy  
in all vitamin A  
deficiencies... particularly  
those associated with  
conditions characterized  
by poor fat absorption  
(dysfunction of the  
liver, pancreas, biliary  
tract and intestines;  
celiac and other  
diarrheal diseases).

Proven effective in  
ACNE and other dermal  
lesions responsive to  
high potency vitamin A.

**"DIHYDROTACHYSTEROL...**

has proved to be the most valuable remedy for the control of chronic hypoparathyroidism and by its use patients may be kept free of the symptoms of tetany as well as of the trophic manifestations of chronic hypocalcemia.\*\*


Orally administered, Hytakerol facilitates the absorption of calcium from the bowel and increases the excretion of phosphate by the kidney.

Sulkowitch's simple urinary test may be used instead of repeated blood analyses in checking calcium levels.\*

**In Chronic  
Hypoparathyroidism...  
Hypocalcemia**

# Hytakerol<sup>®</sup>

Brand of  
Dihydrotachysterol



\*Orellman, A.: Essentials of Endocrinology. Philadelphia, J. B. Lippincott Co., 1947, 2nd ed., p. 269.

**DOSE**

Orally from 3 to 10 cc. (or from 6 to 20 capsules) daily until tetany is relieved; weekly maintenance dose from 1 to 7 cc. (or from 2 to 14 capsules) depending upon the blood and urine calcium levels. From 10 to 15 Gm. calcium lactate or gluconate may be given daily as supplement through the period of therapy.

**SUPPLIED**

**Hytakerol in Oil**, bottles of 15 cc.  
**Hytakerol Capsules** (each equivalent to 0.5 cc.), bottles of 50.

*Winthrop-Stearns* INC.  
NEW YORK 18, N. Y. WINDSOR, ONT.

Hytakerol, trademark reg. U. S. & Canada



## C O N T E N T S

## The American Journal of Medicine

Vol. XI DECEMBER, 1951 No. 6

*Editorial*

- Alterations in Normal Bacterial Flora of Man and Secondary Infections during Antibiotic Therapy . . . . . CHESTER S. KEEFER 665
- Notice to Subscribers . . . . . 666

*Clinical Studies*

- Clinical and Physiologic Features of Some Types of Pulmonary Diseases with Impairment of Alveolar-capillary Diffusion. The Syndrome of "Alveolar-capillary Block"
- ROBERT AUSTRIAN, JOHN H. McCLEMENT, ATTILIO D. RENZETTI, JR., KENNETH W. DONALD, RICHARD L. RILEY AND ANDRÉ Cournand 667

In recent years impairment of diffusion of oxygen across the alveolar-capillary interface has come to the fore as an important factor in pulmonary dysfunction. This "alveolar-capillary block," as the present authors designate it, may occur in a variety of diseases affecting the lungs and may play a significant role in the causation of dyspnea. A composite physiological, clinical and pathological analysis of the condition is given in this interesting study.

- Primary Pulmonary Hypertension. I. Clinical and Hemodynamic Study
- DAVID T. DRESDALE, MARTIN SCHULTZ AND ROBERT J. MIGHTOM 686

This is an interesting and constructive paper in which the authors define more clearly the concept of primary pulmonary hypertension on the basis of distinct clinical, hemodynamic, roentgenographic, electrocardiographic and morphologic findings. It should now be possible to make the diagnosis more frequently during life, indeed presumptive diagnosis would seem to be feasible on clinical grounds alone. Of therapeutic interest is the effect of priscoline in lowering the pulmonary artery blood pressure and the possibility that sympathectomy may be beneficial.

- Pulmonary Function Studies in Polycythemia Vera. Results in Five Probable Cases
- WALTER NEWMAN, JAMES A. FELTMAN AND BLANCHE DEVLIN 706

This study of pulmonary function tests in five cases of probable or possible polycythemia vera revealed interesting alterations in pulmonary ventilation and gas exchange, and in respiratory center responsiveness, which may play an important part in the symptomatology of the disorder. The importance of phlebotomy is stressed.

*Contents continued on page 5*

# diagnosis during THE SILENT PERIOD

Routine testing with Ames diagnostic tablets

in the office, home or hospital is

frequently rewarded by the discovery of

unsuspected pathology during the

"silent period"—when disease in its incipency

may be most easily managed or cured.

Ames tablet tests are completely self-contained,  
rapid and clinically reliable.

## CLINITEST

(BRAND)

for the detection of  
glycosuria



### Occurrence

- diabetes mellitus
- hyperthyroidism
- renal or alimentary glycosuria

## BUMINTEST

(BRAND)

for the detection of  
albuminuria



- nephritis
- renal tuberculosis
- renal tumors
- drug poisoning
- toxemia of pregnancy
- liver disease with  
and without jaundice

## HEMATEST

(BRAND)

for the detection of  
occult blood



- inflammatory, neoplastic and  
degenerative lesions of the  
gastro-intestinal, genito-  
urinary and pulmonary systems

## ACETEST

(BRAND)

for the detection of  
acetonuria



- diabetic acidosis
- Von Gierke's disease
- cyclic or excessive vomiting
- diarrhea

*Available through your pharmacy or physician supply house*

*Clinitest, Bumintest, Hematest, Acetest, trademarks reg. U. S. and Canada*



**AMES COMPANY, INC • ELKHART, INDIANA**

Ames Company of Canada, Ltd., Toronto

DI-3



## C O N T E N T S

## The American Journal of Medicine

Vol. XI DECEMBER, 1951 No. 6

*Contents continued from page 3*

## Effect of Cortisone and ACTH on Fluid and Electrolyte Distribution in Man

MARVIN F. LEVITT AND MORTIMER E. BADER 715

Using inulin distribution as a measure of extracellular volume, the authors found a significant increase in extracellular water, sodium and chloride after ACTH or cortisone administration, which reached a peak in eight or nine days and then regressed despite continuation of therapy. This interesting effect of the adrenal cortex has significant implications.

## Experience with Methimazole (Tapazole) in the Treatment of Hyperthyroidism. A

Report of Thirty-five Cases . . . BERNARD L. HALLMAN AND PHILIP K. BONDY 724

Methimazole (1-methyl-2-mercaptoimidazole) has approximately 100 times the antithyroid activity of thiouracil. According to the experience with thirty-five patients described in this paper, this drug in small dosage rapidly effects a euthyroid state and appears to be of a low order of toxicity. Despite the diminishing indications for the use of antithyroid drugs, methimazole would seem to deserve more extensive trial.

*Seminars on Arteriosclerosis*

Arteriosclerosis—Some Clinical Implications . . . . . WILLIAM DOCK 730

In this essay Dr. Dock discusses the clinical implications of arteriosclerotic involvement of the brain, retina, kidneys, heart, legs, aorta, iliac and splanchnic arteries. As always, he is stimulating and provocative, bringing out new facts and new facets of old facts about this vital subject.

Diet and Lipotropic Agents in Arteriosclerosis . . . . . JACK D. DAVIDSON 736

To close the Seminars on Arteriosclerosis, Dr. Davidson gives a much needed critical analysis of the results of dietary restriction and administration of lipotropic agents in experimental and clinical arteriosclerosis. Proper evaluation of effects is difficult even in the most rigidly controlled animal experiments, of which there are few, to say nothing of clinical trials. It is concluded that restriction of diet with respect to cholesterol and fat seems justifiable in patients with arteriosclerosis despite the lack of convincing proof of beneficial effect. The usefulness of choline and inositol in prevention or cure of arteriosclerosis remains to be demonstrated.

*Conference on Therapy*

Use of Curare and Curare-like Agents . . . . . 749

Conferences on Therapy (Cornell University Medical College)—This timely clinic summarizes the present status of *d*-tubocurarine, flaxedil, syncurine and other agents employed as muscle relaxants in conjunction with anesthetics. The pharmacological action of these drugs, their clinical indications and the dangers inherent in their use, particularly in regard to respiration, are discussed clearly and informatively.

*Contents continued on page 7*

## *Anxiety and Autonomic Lability as the Basis of Functional Disorders*

Emotional response to the stress of life is the primary source of illness in a steadily increasing number of cases. *Weiss* and *English*<sup>1</sup> estimate that as high as two-thirds of all patients have disorders due either entirely or in part to emotional factors and anxiety. *Ebaugh*<sup>2</sup> refers to anxiety as "... the universal disease of our times".

Complete examination discloses no organic basis for the symptoms in these cases, yet the clinical picture may mimic a true organic disease.

The symptom-complex usually involves several or-

gan systems.<sup>2,3,4</sup> In such cases, the anxiety is channeled into organ dysfunction via the autonomic nervous system.<sup>2,3,5,6</sup> Some of the effects produced by exaggerated activity of a labile autonomic system are tabulated below. Many of these, it will be noted, are related to the symptoms which feature prominently in functional disorders. The symptoms in any one case are not necessarily limited to one organ system. Usually some are referable to sympathetic hypertonicity, others to parasympathetic hypertonicity.

ORGAN SYSTEM	SYMPATHETIC HYPER- TONICITY	PARASYM- PATHETIC HYPER- TONICITY	SYMPTOMS OF FUNCTIONAL DISORDER	AUTONOMIC LABILITY
GASTRO- INTESTINAL	Hypomotility Hyposecretion Intestinal Atony	Increased Salivation Hypermotility Hypersecretion	Belching Heartburn Nausea & vomiting Mucous diarrhea	When a patient exhibits a clinical picture suggestive of non-organic dysfunction, the diagnosis of Functional Disorder can be facilitated by use of the following indications of Autonomic Lability: Variable Blood Pressure Temperature Variations Changing Pulse Rate Deviations in B.M.R. Exaggerated Cold Pressure Reflex Oculo-cardiac Reflex Abnormalities Glucose Tolerance Alterations
CARDIO- VASCULAR	Rapid heart rate Peripheral vaso- constriction Slight rise in blood pressure	Reduced heart rate Vasodilatation Lowered blood pressure	Palpitation Sinus tachycardia Premature systoles B.P. low in some; elevated in others	
RESPIRATORY	Dry nasopharyngeal mucous membrane Bronchial relaxation	Increased nasophar- yngeal secretion Bronchial constrict- ion Laryngospasm	Dry mouth and throat Difficulty in breathing Sighing respiration	
GENITO- URINARY	Bladder detrusor relaxed; Sphincter contracted Ureter tone and motility decreased	Bladder detrusor contracted; Sphincter relaxed Ureter tone and motility increased	Urinary frequency Difficulty in urinating Dysmenorrhea Menstrual irregularity	

This tabulation is based on data available in references 1 to 6 stated below.

Primarily, the patient visits his physician out of concern over his symptoms. At this point, he is either unaware of his basic emotional problem or ignores it. A complete examination will rule out organic disease and thus reassure the patient. Then, treatment is directed along two lines: First, relieve the patient of subjective distress by drug therapy.\* He will then be more cooperative in discussing his emotional problems. Then, having uncovered the basic problem, guidance is given toward correcting

unhealthy situations and attitudes.

\*The fact that autonomic dysfunction plays a large part in mediating the disturbance suggests autonomic sedation. A number of independent studies indicate that this therapeutic approach is effective.<sup>7,8,9</sup> The investigators used ergotamine tartrate (adrenergic blockade), levo-alkaloids of belladonna (cholinergic blockade) and phenobarbital (central sedation) in the form of **Bellergal** tablets. The total effect is an integrated sedation of the entire A.N.S.

1. WEISS, E., and ENGLISH, O.: *Psychosomatic Medicine*, ed. 2. Saunders Co., 1949.
2. EBAUGH, F. G.: *Postgrad. Med.* 4: 208, 1948.
3. WILLIAMS, E. Y., et al: *J. Nat. M. A.* 42: 32, 1950.
4. WOOLEY, L.: *South. Med. & Surg.* 102: 157, 1940.
5. KATZ, L. N., et al: *Ann. Int. Med.* 27: 261, 1947.
6. KROGER, W. S., et al: *Am. J. Obst. & Gynec.* 39: 328, 1950.
7. KARNOSH, L. J., and ZUCKER, E. M.: *A Handbook Of Psychiatry*, Mosby Co., 1945.
8. HARRIS, L. J.: *Canad. M. A. J.* 58: 251, 1948.
9. SLAGLE, G. W.: *J. Michigan M. Soc.* 41: 119, 1942.

**Sandoz  
Pharmaceuticals**

DIVISION OF SANDOZ CHEMICAL WORKS, INC.  
68 CHARLTON STREET, NEW YORK 14, NEW YORK



## C O N T E N T S

## The American Journal of Medicine

Vol. XI DECEMBER, 1951 No. 6

*Contents continued from page 5**Clinico-pathologic Conference*

## Miliary Pulmonary Infiltration with Hydrothorax and Ascites . . . . . 757

Clinico-pathologic Conference (Washington University School of Medicine)—With the growing realization that histoplasmosis and other disseminated fungus infections occur more commonly than has been appreciated, differentiation from miliary tuberculosis has become more and more important. The problem is complicated by the not infrequent association of fungus infection with tuberculosis and a variety of other debilitating diseases. The present clinic, which deals with just this question, is therefore timely and of unusual interest.

*Case Reports*

## Mobilization of Gouty Tophi by Protracted Use of Uricosuric Agents

T. F. YÜ AND ALEXANDER B. GUTMAN 765

The experience described in this report would make it appear that by carefully regulated use of uricosuric agents for prolonged periods it may be possible not only to prevent formation of new tophi in chronic tophaceous gout but also to mobilize old established tophi.

## Localized Amyloid Deposition in the Lower Respiratory Tract

AARON SCHOTTENFELD, LEON M. ARNOLD, JOHN G. GRUHN AND A. DAVID ETESS 770

This interesting paper calls attention to the occurrence of amyloid deposits in the trachea and bronchi, not associated with amyloidosis elsewhere, causing bronchostenosis and difficulty in respiration.

## Author Index to Volume XI . . . . . 777

## Subject Index to Volume XI . . . . . 780

*Advertising Index on 3rd Cover**Change of address must reach us one month preceding month of issue.*

# FOR THE *Pantry* HOSTESS



"IT'S AS THOUGH someone has opened a faucet and is draining the life out of me" is her way of describing her feeling of chronic fatigue and listlessness. Her blood picture will probably reveal hypochromic microcytic anemia.

Inorganic iron is specific in such cases, but the IBEROL formula is based on the proposition that iron alone is not enough—that hemoglobin formation involves more than iron—that although an iron deficiency is established, other deficiencies are also possible—that recovery may be more rapid and complete if other important elements are supplied to supplement the diet. Notice the formula.

Just three IBEROL Tablets a day, in spite of the high potencies contained, is the average adult dose. For prophylaxis in pregnancy, old age or convalescence, one or two tablets daily are usually sufficient. IBEROL is supplied in bottles of 100, 500 and 1000 tablets. **Abbott**

## IBEROL<sup>®</sup> Tablets

(Iron, B<sub>12</sub>, Folic Acid, Stomach-Liver Digest, with Other Vitamins, Abbott)


### THREE IBEROL TABLETS:

the average daily therapeutic dose for adults, supply:


Ferrous Sulfate.....	1.05 Gm.
(representing 210 mg. elemental iron, the active ingredient for the increase of hemoglobin in the treatment of iron-deficiency anemia)	
Plus these nutritional constituents:	
Thiamine Mononitrate (6 times MDR*).....	6 mg.
Riboflavin (3 times MDR*).....	6 mg.
Nicotinamide (2 times RDA†).....	30 mg.
Ascorbic Acid (5 times MDR*).....	150 mg.
Pyridoxine Hydrochloride.....	3 mg.
Pantothenic Acid.....	6 mg.
Vitamin B <sub>12</sub> .....	30 mcg.
Folic Acid.....	3.6 mg.
Stomach-Liver Digest.....	1.5 Gm.

\*MDR—Minimum Daily Requirement

†RDA—Recommended Daily Dietary Allowance







**COMBINATION**

*for  
supplementary effects  
wherever estrogen-androgen therapy is indicated...*

- i.e.* In fractures and osteoporosis in either sex to promote bone development, tissue growth, and repair.
- i.e.* In the female climacteric in certain selected cases.
- i.e.* In dysmenorrhea in an attempt to suppress ovulation on the basis that anovulatory bleeding is usually painless.
- i.e.* In the male climacteric to reduce follicle-stimulating hormone levels.

A steroid combination which permits utilization of both the complementary and the neutralizing effects of estrogen and androgen when administered concomitantly. Thus certain properties of either sex hormone may be employed in the opposite sex with a minimum of side effects. Each tablet provides estrogens in their naturally occurring, water-soluble, conjugated form expressed as sodium estrone sulfate, together with methyltestosterone.

No. 879—Conjugated estrogens equine  
("Premarin") ..... 1.25 mg.  
Methyltestosterone ..... 10.0 mg.  
Bottles of 100 tablets (yellow)

No. 878—Conjugated estrogens equine  
("Premarin") ..... 0.625 mg.  
Methyltestosterone ..... 5.0 mg.  
Bottles of 100 tablets (red)



**METHYLTESTOSTERONE**

*for combined estrogen-androgen therapy*

Ayerst, McKenna & Harrison Limited  
22 East 40th Street, New York 16, N. Y.



It takes  
more than  
a jugful  
of milk

to equal the riboflavin content of  
"Beminal" Forte with Vitamin C.  
One capsule No. 817 provides 12.5 mg.  
of riboflavin. More than 7 quarts  
of milk would be needed to furnish the  
same amount. This is but one feature  
of "Beminal" Forte with Vitamin C  
which also contains therapeutic  
amounts of other important  
B complex factors and ascorbic acid.

**"Beminal"**  
Forte with  
Vitamin C

No. 817 Each dry-filled  
capsule contains:  
Thiamine HCl (B<sub>1</sub>), 25.0 mg.  
Riboflavin (B<sub>2</sub>), 12.5 mg.  
Nicotinamide, 100.0 mg.  
Pyridoxine HCl (B<sub>6</sub>), 1.0 mg.  
Calc. pantothenate, 10.0 mg.  
Vitamin C  
(ascorbic acid), 100.0 mg.

Supplied in bottles  
of 30, 100, and 1,000.




Ayerst

Ayerst, McKenna & Harrison Limited  
22 East 40th Street, New York 16, N. Y.

5116



# 10 *clinical advantages*




WITH JUST

2

DROPS

## *Privine* .05%



## *in decongestion*

1. potent 2. prompt 3. prolonged 4. free from significant C.N.S. effects 5. few or no systemic effects 6. virtually non-allergenic 7. consonant with local physiologic function 8. non-irritating 9. safe 10. fosters patient cooperation.

PRIVINE® (BRAND OF NAPHAZOLINE) HYDROCHLORIDE 2/1712 M

**Ciba** SUMMIT, NEW JERSEY



# The High Protein, Low Calorie Diet...

## The Modern Dietary Answer to the Problem of Weight Reduction

It is now universally recognized that overweight is the inevitable response of the body to a persistent caloric intake in excess of metabolic needs. Neither underactivity nor overactivity of an endocrine gland of itself ever produces obesity.<sup>1</sup> Except in rare instances, a sound weight reduction program excludes endocrine therapy<sup>2</sup> and involves only a caloric intake below the level of energy expenditure, while maintaining optimal protein, vitamin, and mineral nutrition. The high protein, low fat, low carbohydrate regimen constitutes a modern dietary answer to the problem of weight reduction.

Successful weight reduction has been quickly achieved by sharp restriction in caloric intake even to as low as 450 calories daily, while nitrogen balance was maintained by foods providing adequate protein.<sup>3</sup> Also, satisfactory weight reduction has been attained in obese patients on an ad libitum diet high in protein and adequate in vitamins and minerals,<sup>2</sup> but low in fat and carbohydrate. Such diets contain liberal amounts of broiled, boiled, or roasted lean meat; kidney, liver, and brain may be included. Meat enjoys this favored position because it leaves the feeling of having eaten well and contributes generously of biologically complete protein.

Meat also plays an important role in the reduction diet because it contributes important amounts of iron and of vitamin B complex, which includes niacin, pantothenic acid, pyridoxine, riboflavin, thiamine, and the newly discovered vitamin B<sub>12</sub>. The high palatability of meat, its stimulation of the digestive processes, its satiety value, and its easy and practically complete digestibility are other features of importance in the reduction diet.

1. Newburgh, L. H.: Obesity: In *Clinical Nutrition*, edited by Jolliffe, N.; Tisdall, F. F., and Cannon, P. R., New York, Paul B. Hoeber, Inc., 1950, chap. 28, p. 689.
2. Kunde, M. M.: The Role of Hormones in the Treatment of Obesity, *Ann. Int. Med.* 28:971 (May) 1948.
3. Strang, J. M.; McClugage, H. B., and Evans, F. A.: The Nitrogen Balance During Dietary Correction of Obesity, *Am. J. M. Sc.* 181:336, 1931.

The Seal of Acceptance denotes that the nutritional statements made in this advertisement are acceptable to the Council on Foods and Nutrition of the American Medical Association.



**A m e r i c a n   M e a t   I n s t i t u t e**  
Main Office, Chicago... Members Throughout the United States



IN PENICILLIN THERAPY!

# COMPENAMINE

## A NEW HYPOALLERGENIC PENICILLIN SALT

Through the routine use of Compensamine, reactions to penicillin can be reduced significantly below that encountered with other available forms of penicillin G. This hypoallergenic characteristic of Compensamine permits its use even in known penicillin reactors; in this group it reduces the incidence of reactions by at least 80 per cent. Thus Compensamine brings new safety to penicillin therapy.

A research development of C.S.C. Pharmaceuticals, Compensamine is generically designated as *l*-ephedrine penicillin G. Its clinical behavior and therapeutic performance are identical, unit for unit, with comparable dosage forms of procaine penicillin. Nearly insoluble in water and oil, its dosage forms are of the repository type.

Compensamine is priced identically with procaine penicillin G. Hence economic considerations are no obstacle to its routine use.

*C.S.C. Pharmaceuticals*

A DIVISION OF COMMERCIAL SOLVENTS CORPORATION, 17 E. 42nd St., New York 17, N. Y.

### THREE DOSAGE FORMS

#### COMPENAMINE IS CURRENTLY AVAILABLE IN THREE REPOSITORY DOSAGE FORMS:

- COMPENAMINE (for aqueous injection), in vials.
- COMPENAMINE AQUEOUS, in vials and disposable and permanent syringe cartridges.
- COMPENAMINE in PEANUT OIL, in vials and disposable and permanent syringe cartridges.

The

# weight reduction problem

a more complete solution . . .

## OBEDRIN<sup>plus</sup> THE 60-10-70 DIET


Obedrin Tablets permit full utilization of the appetite depressing action of methamphetamine hydrochloride but eliminate the central nervous stimulation, so the patient does not suffer from nervous irritability and insomnia.

The 60-10-70 Basic diet provides the basic minimum of proteins to maintain nitrogen balance, the basic minimum of carbohydrates to "burn off" excessive fat in storage.

Obedrin Tablets and the 60-10-70 basic diet will permit loss of weight with minimum discomfort, thus inviting patient cooperation.

### FORMULA

\*Semoxydrine  
 Hydrochloride . . . . . 5 mg.  
 Pentobarbital Sodium . . . . . 20 mg.  
 Ascorbic Acid . . . . . 100 mg.  
 Thiamine  
 Hydrochloride . . . . . 0.5 mg.  
 Riboflavin . . . . . 1 mg.  
 Niacinamide . . . . . 5 mg.  
 \*Methamphetamine Hydrochloride  
 Obedrin is supplied in bottles of 100, 500 and 1,000 yellow grooved tablets.



NAME \_\_\_\_\_ 60-10-70 DIET  
 WEIGHT \_\_\_\_\_ DATE \_\_\_\_\_  
 DO NOT EAT EVEN SMALL AMOUNTS OF ANY FOOD NOT ON THIS LIST

**BREAKFAST**  
 Choose only one item on this list. No Sugar — No Butter.  
 Small orange  
 Orange juice, 1/2 glass (1 oz.)  
 1 Medium apple  
 Pineapple (1 slice)  
 1/2 cup strawberries  
 Plum, fruit, 3  
 Bread, 1 slice (white, rye, or whole wheat)  
 Add any one item on this list.  
 Cream crackers, 1  
 Soda crackers, 10  
 Choose one beverage: Sweet tea with sugar—artificial  
 Tea or coffee, no cream, no sugar—artificial  
 Skimmed milk—1 glass

**LUNCH**  
 Choose one, either meat or fish, not both.  
 1 slice = 4 x 7 inches and 1/2" thick  
 Kidney, 1/2 cup  
 Liver, 1/2 cup  
 Veal, 1 slice  
 Hamburger (small cake)  
 Beef (lean, boiled), 2 slices  
 Chicken, game, duck or turkey (1/4 lb.)  
 Corned beef hash, canned, 1/2 cup  
 1 slice = 4 x 2 inches and 1/2" thick  
 Ham, 2 slices  
 Loaf, 2 slices  
 Salmon, 1/4 cup  
 Scallops, 1 cup


**Plats**  
 Ban, 2 slices  
 Cream, 1 cup  
 Cod, 2 slices  
 Crabmeat, 1 cup

A COMPLIMENTARY PAD OF DIET SHEETS AND A TRIAL SUPPLY OF OBEDRIN SENT TO PHYSICIANS ON REQUEST.

# Obedrin

**MASSENGILL** BRISTOL, TENNESSEE





the simplest method of outpatient maintenance

parenteral

control of edema...

just one or two tablets daily

plus an

occasional

injection

*tablets*

**MERCUHYDRIN**

(brand of meralluride)

*with ascorbic acid*

**ORAL MERCURIAL DIURETIC**

To secure the greatest efficacy and all the advantages of Tablets MERCUHYDRIN with Ascorbic Acid, a three-week initial supply should be prescribed ..... 25 to 50 tablets.


**dosage:** One or two tablets daily—morning or evening—preferably after meals.

**available:** Bottles of 100 tablets. Each tablet contains meralluride

60 mg. (equivalent to 19.5 mg. mercury) and ascorbic acid 100 mg.

*L*akeside  
laboratories, INC., MILWAUKEE 1, WISCONSIN

M-18



***1,000,000 globules  
on the head of a pin***



A lipid emulsion whose suspended globules are 1 micron in diameter has been achieved by cooperative medical and pharmaceutical research. 1,000,000 particles of this emulsion can fit on the head of a pin. A by-product of investigations in fat metabolism, it affords an unusually well tolerated, well absorbed source of concentrated calories. Another case where collaboration between investigators in leading medical centers and Upjohn researchers has "paid off" in newer knowledge—and better control of metabolic problems.

## ***Lipomul-Oral***

contains:

Vegetable Oil ..... 40% w/v  
Dextrose, Anhydrous ..... 10% w/v  
Preserved with Sodium Benzoate 0.1%

Lipomul-Oral provides 4 calories per cc., or approximately 120 calories per ounce.

Available in pint bottles.

\*Trademark

a product of


**Upjohn**

**Research for Medicine ... Produced with care ... Designed for health**

THE UPJOHN COMPANY, KALAMAZOO, MICHIGAN


# NEW 5<sup>TH</sup> EDITION

**COMING EARLY 1952**



**Bound in Red Fabri-**  
**coid. 1200 pages, size**  
**6" x 9 1/4" x 2 1/4" Pre-**  
**Publication Price,**  
**Postpaid \$12\* U.S.A.**  
**\$15 Foreign.**

**PRE-PUBLICATION  
SAVING of  
\$300  
IF YOU ORDER  
NOW**

**FREE**

—to every encyclo-  
pedia owner—

## MODERN DRUGS

—the quarterly supplement that keeps you up-to-date with the newest drug descriptions.

**522**     *Modern Drug Encyclopedia and Therapeutic Index*

---

**Administration:** *Orally*, for adults 1 rounded teaspoonful (5 to 7 Gm) thoroughly dissolved in glass of water, milk or fruit juice, followed by another glass of fluid, 1 to 3 times daily; children according to age and weight. Liberal amounts of liquid should be taken during the day.

**Supply:** METADREN (C.A.)—Powder—Package, 4 oz, 5 oz, 1 lb.

**METADREN**                      *Androgen therapy*                      (Chu)

Chemical Name: 17-methyltestosterone.

Formula:

CC12CCC3C(C1CC2=O)CCC4=CC(=C)C(C=C4)C

**Description:** The pure, synthetic, methyl ester of the steroid testosterone, with more marked androgenic properties than testosterone when given orally.

*Completely Rewritten*

to give you **UP-TO-DATE** Reference on nearly 4000 Ethical Drugs (including 1500 brand new listings) of 175 manufacturers.

*Indispensable*

to **PHYSICIANS** for its authoritative, finger-tip data...complete from description to prescription. Now compiled in seven special sections: **DRUGS • BIOLOGICALS • ALLERGENS • GENERIC NAME INDEX • THERAPEUTIC INDEX • MANUFACTURER'S INDEX • GENERAL INDEX.** The new 5th Edition of **The MODERN DRUG ENCYCLOPEDIA** provides self-pronouncing drug listings for the first time.

Edited by  
**MARION E. HOWARD, M.D., F.A.C.P.**  
Yale University Medical School  
New Haven, Conn.

**MORE THAN 50,000  
SATISFIED USERS**

**Saves hours of precious time. Proved by use as America's "best seller" medical encyclopedia.**

**MAIL THIS COUPON TODAY!**

**This Pre-Publication Saving Expires Feb. 1, 1952**  
**DRUG PUBLICATIONS, INC.**  
 49 West 45th Street, New York 36, N. Y.

**Enclosed is the sum of twelve dollars (\$12\* U.S.A.) for which please send me postpaid the new Fifth Edition of the MODERN DRUG ENCYCLOPEDIA AND THERAPEUTIC INDEX and MODERN DRUGS.**

NAME \_\_\_\_\_ M.D.

ADDRESS \_\_\_\_\_

CITY \_\_\_\_\_ ZONE \_\_\_\_\_ STATE \_\_\_\_\_

U.S.A. \$12\* FOREIGN \$15

\* Includes three-year supplementary service at \$2 per year.  
(New York City residents please add 3% sales tax.)

MN



*White's*

**Three  
Important  
Additions  
to  
Therapy  
with  
Penicillin**

**DRAMCILLIN**  
preparations now include:  
**PENICILLIN**

**DRAMCILLIN-500,**  
**DRAMCILLIN-250,**  
(250,000 units\* per tea-  
spoonful), **DRAMCILLIN**  
(100,000 units\* per tea-  
spoonful), **DROPCILLIN**  
(50,000 units\* per drop-  
perful).

**PENICILLIN-  
TRIPLE SULFONAMIDES**  
**DRAMCILLIN-250 with**  
**Triple Sulfonamides,**  
**DRAMCILLIN-250 Tablets**  
**with Triple Sulfonamides,**  
**DRAMCILLIN with Triple**  
**Sulfonamides** (100,000  
units of penicillin and 0.167  
grams each of sulfadiazine,  
sulfamerazine and sulfacet-  
imide per teaspoonful).

\*Buffered Crystalline Penicillin G  
Potassium

## DRAMCILLIN

- On convenient 8 to 12 hour dosage sched-  
ule, Dramcillin-500 produces optimal ther-  
apeutic effect in most infections
- More uniform absorption due to magnitude  
of dosage
- The most economical high potency, liquid  
oral penicillin

## DRAMCILLIN

- fully effective with just 4 doses daily
- the only high potency, liquid penicillin-  
sulfonamide combination containing rap-  
idly absorbed and highly soluble sulfaceti-  
mide

## DRAMCILLIN

- each tablet is equivalent to one teaspoon-  
ful of the above liquid form

With the addition of these three new preparations, DRAMCILLIN—one easily remembered name—now identifies an effective, palatable, and most complete group of products designed to meet every commonly encountered need of the physician in treatment with oral penicillin or penicillin-sulfonamides. White Laboratories, Inc., Kenilworth, N. J.

# 500

One half million units of penicillin\* per teaspoonful.

# 250

WITH TRIPLE SULFONAMIDES 250,000 units penicillin\* with 0.167 grams each sulfadiazine, sulfamerazine, sulfacetamide per teaspoonful.

# 250

TABLETS WITH TRIPLE SULFONAMIDES

they  
deserve  
the  
best..

Implicit in a happy healthy childhood is maximal nutrition—and one of the essential dietetic guideposts to vigorous adulthood is adequate vitamin C<sup>1,2,3</sup> (1/4-4 oz. for infants up to 1 year;<sup>10,11</sup> 4-8 oz. for older children).<sup>4</sup> Fortunately, most every youngster likes the taste of Florida orange juice and the "lift" its easily assimilable fruit sugars\* provide.<sup>6</sup> It is well-tolerated and virtually non-allergenic.<sup>3</sup> And, under modern techniques of processing and storage—it is possible for citrus fruits and juices (whether fresh, canned or frozen) to retain their ascorbic acid content,<sup>5,8</sup> and their pleasing flavor,<sup>7</sup> in very high degree and over long periods.

FLORIDA CITRUS COMMISSION • LAKELAND, FLORIDA

*Citrus fruits—among the richest known sources of Vitamin C—also contain vitamins A and B, readily assimilable natural fruit sugars, and other factors, such as iron, calcium, citrates and citric acid.*

... plenty of citrus fruit

#### References

1. Clinical Nutrition: ed. by N. Jolliffe et al., Hoeber, 1950.
2. Hanke, M. T.: Diet and Dental Health, Chicago Univ. Press, 1933.
3. Jeans, P. C.: J.A.M.A., 142:806, 1950.
4. Joslin, C. L. and Bradley, J. E.: Personal communication.
5. Krehl, W. A. and Cowgill, G. R.: Food Research, 15:179, 1950.
6. McLester, J. S.: Nutrition and Diet in Health and Disease, Saunders, Phila., 4th ed., 1944.
7. Moore, E. L. et al.: J. Home Econ., 37:290, 1945.
8. Roy, W. R. and Russell, H. E.: Food Industries, 20:1764, 1948.
9. Sherman, H. C.: Chemistry of Food and Nutrition, Macmillan, 7th ed., 1946.
10. Silverman, A. C.: New York J. Med., 47:1987, 1947.
11. Stevenson, S. S.: J. Pediat., 31:616, 1947.



**FLORIDA**  
Oranges • Grapefruit  
Tangerines





Now available ...  
a "chemical fence" for the alcoholic

# "ANTABUSE"

"Antabuse"—nearly three years under intensive clinical investigation—is now available for the treatment of alcoholism. By setting up a sensitizing effect to ethyl alcohol, "Antabuse" builds a "chemical fence" around the alcoholic...helps him develop a resistance to his craving. Its high degree of efficacy is confirmed by extensive clinical evidence.

"Antabuse" is safe therapy when properly administered. However, it should be employed only under close medical supervision. Complete descriptive literature is available and will be gladly furnished on request.

"Antabuse" is identical with the material used by the original Danish investigators, and is supplied under license from Medicinalco, Copenhagen, Denmark. U. S. Pat. No. 2,567,814.

Supplied in  
tablets of 0.5 Gm.,  
bottles of 50  
and 1,000.



Tested in more than 100  
clinics...by more than 800 qualified investigators  
...on more than 5,000 patients...and covered by  
more than 200 laboratory and clinical reports.

# "ANTABUSE"

...brand of specially prepared and highly purified tetraethylthiuram disulfide.

AYERST, McKENNA & HARRISON LIMITED  
New York, N. Y. Montreal, Canada



# Cortone<sup>®</sup>

## for topical use in eye diseases

CORTONE has proved remarkably effective in the treatment of many inflammatory eye diseases. *Topical* administration is indicated principally in disorders of the anterior segment—the cornea and anterior uvea.

### *Three products for Individualized Dosage:*

#### OPHTHALMIC SUSPENSION OF CORTONE Acetate

**0.5%—5 cc. vials:** For more superficial and less serious indicated conditions and for continuing treatment in severe eye conditions after preliminary therapy with the 2.5% strength has achieved the desired degree of improvement.

**2.5%—5 cc. vials:** For treatment of the more severe indications and for initial therapy of any indicated condition that potentially might lead to permanent ocular damage.

#### OPHTHALMIC OINTMENT OF CORTONE Acetate

**1.5%—3.5 Gm. tubes:** For use in conjunction with either of the ophthalmic suspension preparations, or alone, depending on the condition present; particularly useful for bedtime application.

Administered topically in recommended dosage, CORTONE is

***Nonirritating — Safe — Economical***

*Literature on Request*


**Cortone<sup>®</sup>**  
ACETATE  
(CORTISONE Acetate Merck)

For diseases of the deeper structures of the eye, it is recommended that adequate systemic dosage with the *Oral Tablets* or the *Parenteral Suspension* of CORTONE accompany or precede topical application.



**MERCK & CO., INC.**  
*Manufacturing Chemists*  
RAHWAY, NEW JERSEY  
In Canada: MERCK & CO. Limited—Montreal

CORTONE is the registered  
trade-mark of Merck & Co., Inc.  
for its brand of cortisone.



*in lobar pneumonia:*

The prompt response to Terramycin therapy in lobar pneumonia is consistent with results obtained in primary atypical pneumonia, bronchopneumonia and many other infections of the respiratory tract. In a typical series of pediatric cases, Terramycin-treated, "temperatures returned to normal in 24 to 48 hours after therapy was begun. The clinical appearance of marked improvement took place during the same period."

*Potterfield, T. G., and Starkweather, G. A.:  
J. Philadelphia General Hosp. 2:6 (Jan.) 1951*

CRYSTALLINE TERRAMYCIN HYDROCHLORIDE

available | Capsules, Elixir, Oral Drops, Intravenous,  
Ophthalmic Ointment, Ophthalmic Solution.

ANTIBIOTIC DIVISION



CHAS. PFIZER & CO., INC., Brooklyn 6, N. Y.





*known  
and  
relied on  
the world over*

# ADRENALIN<sup>®</sup>

ADRENALIN (epinephrine, Parke-Davis) is today, as it has been for many years, one of the most versatile and useful drugs, known and used the world over. Introduced to the medical profession by PARKE-DAVIS in 1901, ADRENALIN is widely used in many conditions—bronchial asthma, serum sickness, the Adams-Stokes syndrome, and anesthesia accidents.

Circulatory stimulant, vasoconstrictor, resuscitant, and hemostatic, this pure crystalline hormone is one of the truly basic drugs—an invaluable aid in office, in hospital, and in clinic. It is an important adjunct in local anesthesia, valuable in arresting superficial hemorrhage, and a standby for decongestion of engorged mucous membranes.

ADRENALIN is available as ADRENALIN CHLORIDE SOLUTION 1:1000, ADRENALIN CHLORIDE SOLUTION 1:100, ADRENALIN IN OIL 1:500 and in a variety of forms to meet all medical and surgical requirements.

**PARKE, DAVIS & COMPANY**



THOMAS NELSON & SONS

*announces*

*that its Medical Book Department  
has been acquired by*

THE WILLIAMS & WILKINS COMPANY

Mt. Royal and Guilford

Baltimore 2, Maryland

All medical titles, including Golden's

*Diagnostic Roentgenology*

and the 1951 Renewal Pages, formerly  
published by Nelson, may be procured from

Williams & Wilkins.

# Clinical success in postpartum hemorrhoids



From a clinical report of 79 cases of postpartum hemorrhoids, treated with RECTAL MEDICONE at a large New York institution, the following results were tabulated:

NO. OF CASES	TYPE	RESULTS
41	SUBACUTE	38 SATISFACTORY RESPONSE
22	ACUTE	22 RELIEF IN ALL CASES
16	CHRONIC	10 SATISFACTORY RESPONSE (4 of the 16 cases required surgery)

The explanation for these highly favorable results in this painful condition lies in the fact that RECTAL MEDICONE SUPPOSITORIES contain benzocaine for topical anesthesia—reinforced by other effective anti-hemorrhoidal agents, which promote retrogression and healing.



Ethically promoted by  
MEDICONE COMPANY  
225 Varick St., New York 14, N. Y.

**R E C T A L M E D I C O N E**



a practical  
*step forward*

in the therapy of  
rheumatic affections



Pabalate usually provides better therapy for rheumatic affections than pure salicylate itself, through its mutually synergistic combination of para-aminobenzoic acid and salicylate.<sup>1,2</sup> Reports of authoritative clinical tests show a *higher* degree of pain relief... to *more* patients... on *lower* dosage... over *longer* periods... with *greater freedom* from adverse reactions.<sup>3</sup>

**REFERENCES:** 1. Dry, T. J. et al.: Proc. Staff Meetings Mayo Clin., 21:497, 1946. 2. Hoagland, R. J.: Am. J. Med., 9:272, 1950. 3. Smith, R. T.: J. Lancet, 70:192, 1950.


**FORMULA:** Each enteric-coated tablet or each teaspoonful of chocolate-flavored liquid contains 0.3 Gm. (5 gr.) sodium salicylate U.S.P., and 0.3 Gm. (5 gr.) para-aminobenzoic acid (as the sodium salt).

**A. H. ROBINS CO., INC.**  
RICHMOND 20, VA.




**Pabalate®**


Also available as Pabalate Sodium-Free for cases in which sodium is contraindicated



Seven Years' Research that opened a **New Era**



# in Cough Therapy



The time encrusted *empirical* approach to cough therapy—with its “picture of confusion, contradiction and neglect”<sup>2</sup>—can now be replaced by sound, *rational* non-narcotic treatment, thanks to the pioneering studies of Boyd et al.,<sup>1,2</sup> demonstrating that glyceryl guaiacolate is the most powerful of expectorants commonly used. Robitussin—the antitussive-expectorant with specific drug action—provides glyceryl guaiacolate for increasing respiratory tract fluid, together with desoxyephedrine, for its bronchial-spasm-relieving<sup>3</sup> and its mood-improving actions—in an aromatic syrup that is highly patient-acceptable.

A. H. ROBINS CO., INC. • RICHMOND 20, VA.

## Robitussin®



*promotes useful cough...  
minimizes harmful cough*

### References:

1. Boyd, E. M. and Lapp, S.: J. Pharmacol. and Exper. Therap., 87:24, 1946.
2. Connell, W. F. et al.: Canad. M.A.J., 42:220, 1940.
3. Novelli, A. and Tainter, M. L.: J. Pharmacol., 77:324, 1943.

### Formula:

Each 5 cc. (1 teaspoonful) contains 100 mg. glyceryl guaiacolate and 1 mg. desoxyephedrine hydrochloride, in a palatable aromatic syrup.



(The Council on Pharmacy and Chemistry of the American Medical Association has adopted the following statement of Actions and Uses and of Dosage for publication in connection with a description of Banthine Bromide for inclusion in New and Nonofficial Remedies)

## METHANTHELIN BROMIDE.—*Banthine*<sup>®</sup> Bromide (Searle)

$\beta$ -diethylmethylaminoethyl 9-xanthenecarboxylate bromide

**Actions and Uses.**—Methantheline bromide, a parasympatholytic agent, produces both the peripheral action of anticholinergic drugs such as atropine and the ganglionic blocking action of drugs such as tetraethylammonium chloride. Tolerated amounts of methantheline bromide exert side effects typical of atropine-like drugs, but cause less tachycardia, and also less postural hypotension than does tetraethylammonium chloride. Toxic doses produce a curare-like action at the somatic neuromuscular junction.

Clinical studies indicate that the drug effectively inhibits motility of the gastrointestinal and genitourinary tracts and, to a variable degree, diminishes the volume of perspiration and salivary, gastric and pancreatic secretions. It also decreases mucoprotein secretion. Like atropine, it produces mydriasis and cycloplegia when applied locally to the eye or administered systemically, but until more clinical evidence becomes available, its local use for this purpose is not recommended. The value of the drug for preventing abnormal cardiac reflexes through the vagus during thoracic surgery, or as an agent for routine preoperative medication in place of atropine, requires further investigation before final conclusions can be reached.

Methantheline bromide is indicated for clinical use whenever anticholinergic spasmolytic action is desired, provided it is not contraindicated because of its atropine-like characteristics or because of a patient's intolerance to the unavoidable side effects of such therapy. It is useful as an adjunct in the management of peptic ulcer, chronic hypertrophic gastritis, certain less specific forms of gastritis, pylorospasm, hyperemesis gravidarum, biliary dyskinesia, acute and chronic pancreatitis, hypermotility of the small intestine not associated with organic change, ileostomies, spastic colon (mucous colitis, irritable bowel), diverticulitis, ureteral and urinary bladder spasm, hyperhidrosis or control of normal sweating which aggravates certain dermatoses, and control of salivation.

Methantheline bromide produces some degree of cycloplegia and mydriasis in therapeutic doses and

therefore should not be administered to patients with glaucoma. It sometimes decreases the ability to read fine print. Xerostomia (dryness of the mouth) is a common, sometimes transient, side effect. Urinary retention of varying degree may occur in elderly male patients with prostatic hypertrophy, and some patients may have difficulty emptying the rectum. Patients with edematous duodenal ulceration may experience nausea and vomiting during initial administration of the drug. These patients should take only liquids during the institution of drug therapy. All patients should be advised of the possible occurrence of side effects. Overdosage sufficient to produce a curare-like action may be counteracted by prompt subcutaneous injection of 2 mg. of neostigmine methylsulfate.

**Dosage.**—Methantheline bromide is administered orally or parenterally by either the intramuscular or intravenous route. Parenteral administration is not advised for patients able to take the drug orally. The average initial adult dose, oral or parenteral, is 50 mg. For patients with considerable intolerance, 25 mg. may be employed. In the management of peptic ulcer, a beginning schedule of 50 mg. three times daily before meals and 100 to 150 mg. on retiring is suggested. However, the usual effective dose is 100 mg. four times daily, although some patients may require more or less than this amount. The dosage may be increased to tolerance, using dryness of the mouth as a guide, and adjusted to meet the individual response of patients. Maintenance dosage in peptic ulcer is usually considered to be about one-half the therapeutic level. In the management of other hypermotile or hypersecretory states, the dosage should be adjusted to the smallest amount which will relieve the symptoms. When spastic conditions are secondary to inflammatory or other organic lesions, therapy directed toward the cause should be employed whenever possible.

G. D. SEARLE & CO.

Tablets Banthine Bromide: 50 mg.

Ampuls Banthine Bromide: 50 mg.





the new repository


**ACTH** preparation

means economy for more patients...



greater convenience of administration for home and office treatment





# ACTHAR<sup>\*</sup> *Gel*

## IS LONG-ACTING

ACTHAR *Gel*—a new repository ACTH preparation—is an important contribution to home and office treatment. A single injection in many cases provides an adequate daily dosage. Thus simplified administration plus a considerable price reduction of ACTHAR, which is fully reflected in the price of ACTHAR *Gel*, provides further economy of ACTH therapy.

ACTHAR *Gel* possesses all the efficacy of ACTHAR in aqueous solution and is well tolerated locally, whether administered intramuscularly or by deep subcutaneous injection.

**Indications:** Rheumatoid arthritis, rheumatic fever, acute lupus erythematosus, drug sensitivities, severe bronchial asthma, contact dermatitis, most acute inflammatory diseases of the eye, acute pemphigus, exfoliative dermatitis, ulcerative colitis, acute gouty arthritis, secondary adrenal cortical hypofunction.

**Supplied:** In 20 I.U. (mg.) and 40 I.U. (mg.) per cc. in 5 cc. vials.

\*THE ARMOUR LABORATORIES BRAND OF ADRENOCORTICOTROPIC HORMONE (A.C.T.H.)



**THE ARMOUR LABORATORIES** CHICAGO 11, ILLINOIS

*—world-wide dependability—*

PHYSIOLOGIC THERAPEUTICS THROUGH BIORESEARCH



# Dexedrine\* tablets & elixir

the antidepressant of choice, and the most effective  
drug for control of appetite in weight reduction



Smith, Kline & French Laboratories, Philadelphia

\*T.M. Reg. U.S. Pat. Off.



# ANNOUNCING HED-HEPARIN

HEPARIN SODIUM INJECTION U. S. P.

10 cc. vials 1000 and 5000 U.S.P. Units per cc.

The first of a group of improved medicinal agents produced by HED Pharmaceuticals, Inc., under the personal supervision of H. E. Dubin, Ph.D., formerly Director, Biochemical Department of H. A. Metz Laboratories and formerly Technical Director of U. S. Vitamin Corporation and H. E. Dubin Laboratories, Inc.

HED-HEPARIN is a superior product—at surprisingly economical cost to your patient—the result of efficiencies and improvements devised by HED Pharmaceuticals. Indications: Pulmonary Embolism, Coronary Thrombosis, Thrombophlebitis, Frostbite, Myocardial Infarction.

Newer Applications: Atherosclerosis, Angina Pectoris



"I have established HED Pharmaceuticals on the basic principle that there can be no compromise with quality. You may accept HED-HEPARIN with the knowledge that nothing has been left undone to assure therapeutic efficacy." *H. E. Dubin*

Your physicians supply dealer now has HED-HEPARIN. Should he be temporarily out of stock, HED PHARMACEUTICALS will gladly supply you direct and bill you through your dealer.

*"No. Compromise with Quality"*

**HED Pharmaceuticals, Inc.**  
NEW ROCHELLE • NEW YORK

...the best is yet to be



For those approaching middle life, the years ahead *can* be the best — provided normal metabolic functions are safeguarded. In such interrelated disorders as atherosclerosis, diabetes mellitus, and liver disease, the clinical findings are likely to include abnormal fat metabolism (with accompanying deposition of cholesterol) and abnormal capillary fragility.

Prophylaxis against these threats to the older patient may be established and maintained with . . .

schenley

VASCUTUM<sup>Trademark</sup>

*for the life that begins at forty*

VASCUTUM\* presents an unusually complete lipotropic combination *plus* the specific capillary protectants, rutin and ascorbic acid.

The average daily dose (6 tablets) provides:

Choline	1 Gm.	Pyridoxine HCl	4 mg.
Inositol	1 Gm.	Rutin	150 mg.
dl-Methionine	500 mg.	Ascorbic Acid	75 mg.

SUPPLIED: Bottles containing 100 tablets

SCHENLEY LABORATORIES, INC.  
LAWRENCBURG • INDIANA

for efficient  
nasal decongestion

*Wyeth*



**LOCAL ACTION ONLY**

- ▶ No elevation in blood pressure
- ▶ No central nervous stimulation

Your patient can take Wyamine even at night and secure restful sleep. Relief begins in minutes . . . lasts for hours.

*Available as:* Solution Wyamine Sulfate, bottles of 1 fl. oz.

Wyamine-Tyrothricin Nasal Solution, bottles of 1 fl. oz. - with dropper or JETOMIZER®

Wyamine-Penicillin, Capsules, Penicillin with Vasoconstrictor, for preparation of nasal solutions.

**WYAMINE®**

MEPHENTERMINE

N-METHYLPHENYL—TERTIARY—BUTYLAMINE WYETH

*Wyeth* Incorporated, Philadelphia 2, Pa.



# yes *the answer to more*

*effective treatment of*

## *many common infections is* **NEOPENZINE**

(Penicillin with Sulfonamides, Lilly)

Has this combination of penicillin in high dosage with mixed sulfonamides a wider spectrum of antibacterial activity than any single bacteriostatic or bactericidal agent?	→ yes
Can orally administered penicillin cause fewer hypersensitive reactions than injected penicillin?	→ yes
Is penicillin as effective by the oral route as by injection if given in adequate doses?	→ yes
Is there positive assurance that the penicillin in 'Neopenzine' is protected from potency loss?	→ yes
Has the mixture of sulfonamides in 'Neopenzine' been demonstrated to be safer systemic therapy than single sulfonamides?	→ yes
Can 'Neopenzine' save time for the busy practitioner?	→ yes

Each Tablet 'Neopenzine' contains 300,000 units of penicillin—G, crystalline-potassium. Each 5 cc. (approximately one teaspoonful) of Suspension 'Neopenzine' contain 200,000 units. The latter is prepared by combining the contents of the potency-protector package. Each tablet or each 5 cc. of suspension also contain 0.167 Gm. each of sulfadiazine, sulfamerazine, and sulfamethazine.

*Dosage for adults:* 2 tablets four times daily.

*Dosage for children:* Suspension (or tablets) in proportion to weight.

Detailed information and literature on 'Neopenzine' are personally supplied by your Lilly medical service representative or may be obtained by writing to

ELI LILLY AND COMPANY • INDIANAPOLIS 6, INDIANA, U.S.A.

## Editorial

### Alterations in Normal Bacterial Flora of Man and Secondary Infections during Antibiotic Therapy

IT is now firmly established that striking alterations in the bacterial flora in various regions of the body may occur following the use of antibiotics.<sup>1-3</sup> This change in the bacterial flora is independent of the effect of the antibiotic on the specific organism responsible for an infection. Thus when penicillin is given, susceptible and sensitive gram-positive bacteria may be eliminated from the mouth and nasopharynx and only non-sensitive and gram-negative organisms can be isolated. When streptomycin is administered, susceptible gram-negative organisms disappear and only gram-positive organisms survive. Following the exhibition of aureomycin, terramycin or chloramphenicol all of the susceptible gram-positive as well as gram-negative organisms may disappear, leaving only monilia, or other non-sensitive bacteria or yeasts. In any event it appears that the surviving bacteria will depend upon the susceptibility of the microorganisms in the original flora, the development or appearance of resistant strains or the introduction of new and resistant strains of microorganisms from outside.

In many instances these changes in the local bacterial flora are unimportant and inconsequential; thus they would fail to attract attention unless careful bacteriologic cultures were made. In other cases, however, a new infection

appears. These new infections may arise in one of several ways. (1) New organisms may be introduced from outside, i.e., from carriers or from contact with other infections, or by the accidental introduction of new organisms during the manipulations incident to proper administration of the antibiotic agent. (2) Organisms which are present in small numbers and which are not sensitive to the antibiotic may flourish and invade tissues after the susceptible organisms are inhibited. (3) Organisms appear rapidly which have become resistant after exposure to an antibiotic.

Weinstein<sup>4</sup> has reported cases of staphylococcal pneumonia, bacteremia and pyelonephritis due to staphylococci in patients who had an original Hemophilus influenzae infection and who were treated with streptomycin. A number of other observers have reported oral, intestinal, bronchopulmonary, genital, dermal and generalized infections due to monilia<sup>5</sup> following the

<sup>4</sup> WEINSTEIN, L. Alterations in normal bacterial flora of man and animals and secondary infection during streptomycin therapy. *Streptomycin, Its Nature and Practical Application*. Edited by S. A. Waksman. Baltimore, 1949. Williams and Wilkins. The spontaneous occurrence of new bacterial infections during the course of treatment with streptomycin or penicillin. *Am. J. M. Sc.*, 214: 56, 1947. The treatment of meningitis due to Hemophilus influenzae with streptomycin. *New England J. Med.*, 235: 101, 1946.

<sup>5</sup> CROSS, W. G. Oral reactions to penicillin. *Brit. M. J.*, 1: 171, 1949. HEWITT, W. L. and WILLIAMS, B. Chloromycetin (chloramphenicol) in the treatment of infections. *New England J. Med.*, 242: 119, 1950. HARRIS, H. J. Aureomycin and chloramphenicol in brucellosis, with special reference to side effects. *J. A. M. A.*, 142: 161, 1950. WILLIAMS, B., JR. Oral and pharyngeal complications of chloramphenicol (chloromycetin) therapy. *Am. Pract. & Digest. Treat.*, 1: 897, 1950. WOODS, J. W.,

<sup>1</sup> LIPMAN, M. O., COSS, J. A. and BOOTS, R. H. Changes in bacterial flora of the throat and intestinal tract during prolonged oral administration of penicillin. *J. Bact.*, 51: 594, 1946.

<sup>2</sup> SMITH, J. W. and BLOOMFIELD, A. L. The effect of penicillin on the aerobic bacterial flora of the normal throat. *Stanford M. Bull.*, 6: 469, 1948.

<sup>3</sup> SOMMER, L. S. and FAVOUR, C. B. Biologic complications of penicillin therapy. *Am. J. Med.*, 7: 511, 1949.

use of penicillin, aureomycin, chloramphenicol and terramycin therapy. Finland<sup>6</sup> records instances of staphylococcic diarrhea in patients receiving terramycin, and Weinstein and his associates<sup>7</sup> have recently called attention to the high incidence of suppurative otitis media and bronchopneumonia in patients with pertussis who received either aureomycin or chloramphenicol. Also, Chang and Jackson<sup>8</sup> have noted

MANNING, I. H., JR. and PATTERSON, C. N. Monilial infections complicating the therapeutic use of antibiotics. *J. A. M. A.*, 145: 207, 1951. GEIGER, A. J., WENNER, H. A., AXILROD, H. D. and DURLANDER, S. H. Mycotic endocarditis and meningitis. *Yale J. Biol. & Med.*, 18: 259, 1945.

<sup>6</sup> FINLAND, M. The present status of antibiotics in bacterial infections. *Bull. New York Acad. Med.*, 27: 199, 1951.

<sup>7</sup> Personal communication.

<sup>8</sup> Quoted by Finland.

scarlet fever due to hemolytic staphylococcus aureus infections following the treatment of pertussis with the broad spectrum antibiotics, or the use of prophylactic penicillin for the treatment of a burn. These are a few of the new or superimposed infections that have been reported following antibiotic therapy, and they only serve to point up this important problem. It is necessary for all physicians to be alert to the problem and to continue to study this complex phenomenon so that ways and means may be found to prevent new infections as well as treat them when they arise. Bacteriologic studies continue to remain important in the treatment of patients with infectious diseases because new or resistant infections can be recognized only by following the changes in bacterial flora in various regions of the body.

CHESTER S. KEEFER, M.D.

## Notice to Subscribers

THE following excerpt is representative of several letters that have been received requesting that the publishers of the *American Journal of Medicine* make available a complete index of past issues of the *Journal*.

"Since I left \_\_\_\_\_ Hospital in July 1948, the most instructive hours of postgraduate education have been those which I have devoted to reading *The American Journal of Medicine*. Faced with the problem of a comprehensive review for my American Board Examination, with the somewhat limited time at my disposal, I resorted to re-reading the issues of the *Journal* as my formal review exercise. This I found most profitable. Because my office is not in the same building as the local medical library, I frequently go back over the *Journal* for information on some particular topic, usually with productive results. In this regard I have often wondered whether you had ever considered the feasibility and practicality of printing a complete index every two years covering the preceding years' issues. This as I gather it would involve a considerable expense to the publishers. I have discussed the matter with a fair number of my colleagues in this city, all of whom felt they would be willing to pay for such an index in addition to the regular subscription fee.

"I hope that this does not seem like too hare-brained an idea to you. I am sure you can understand it was prompted by the fact that the private practitioner, out of the hospital environment, is no longer able to assimilate by osmosis the many new ideas and facts that he did during his house officer days in the large teaching hospital."

(Signed) L. O. J.  
R. M. W.  
North Carolina

The Editorial Board and the publishers of *The American Journal of Medicine* are most anxious to make the contents of the *Journal* maximally available for teaching and review purposes, and have therefore decided to issue a complete index of articles and abstracts which appeared in the first five years of publication. This will be published separately, virtually at cost. The index will include a complete author and subject index from July, 1946, through June, 1951. Preparation of this publication is almost complete and it is scheduled to appear early in 1952. Announcement of publication will be made in *The American Journal of Medicine* and a notice will be sent to all subscribers.

THE EDITOR



# Clinical Studies

## Clinical and Physiologic Features of Some Types of Pulmonary Diseases with Impairment of Alveolar-capillary Diffusion\*

### *The Syndrome of "Alveolar-capillary Block"*

ROBERT AUSTRIAN, M.D., JOHN H. McCLEMENT, M.D., † ATTILIO D. RENZETTI, JR., M.D.,  
KENNETH W. DONALD, M.D., RICHARD L. RILEY, M.D. and ANDRÉ COURNAND, M.D.

*New York, New York*

IN recent years attempts have been made to isolate patterns of pulmonary dysfunction in a variety of chronic pulmonary diseases.<sup>1</sup> Studies of the physiopathology of the lungs have suggested that alveolar respiratory insufficiency, manifested by arterial oxygen unsaturation and sometimes also by carbon dioxide retention, could result from disturbances of the ventilation-perfusion relationships or from alterations of the pulmonary diffusing surface. The purpose of this paper is to describe a syndrome caused by a variety of pathologic processes and characterized histologically by alterations of the pulmonary diffusing surface, i.e., the alveolar-capillary septa, and physiologically by a reduction in the oxygen diffusing capacity of the lungs. The group of subacute or chronic diseases which leads to this syndrome have common clinical characteristics, prominent among which are the diffuse, finely dispersed pulmonary lesions as revealed by x-ray and those signs and symptoms which result from the physiologic disturbances. The term "alveolar-capillary block" is tentatively offered to identify this syndrome as it describes the essential histologic as well as the principal physiologic features.

The concept that altered permeability of the pulmonary membrane to respiratory gases may cause a reduction in oxygen diffusing capacity is not new. Schjerner<sup>2</sup> in 1922, in attempting to elucidate the early cyanosis of "grippe pneu-

monia" described by Brauer, suggested that alterations might occur in the lungs without microscopically visible change which could lead to disturbances in the diffusion of oxygen. Between 1932 and 1935 several German authors<sup>3-5</sup> wrote of a syndrome termed by them "pneumonosis" which was believed to arise from decreased permeability of the pulmonary diffusing surface to oxygen and to result from a variety of causes. Among the etiologic agents included were "grippe pneumonia," war gases, chronic pulmonary congestion, amyloid deposits and xanthomatosis. The syndrome was characterized by the development of cyanosis, the cause of which could not be attributed to ventilatory or circulatory disturbances. The complexity of the problem of analyzing the oxygen tension gradient across the pulmonary membrane was appreciated fully, and for lack of sufficiently precise means of analysis the diagnosis of "pneumonosis" remained one of exclusion unless material became available for pathologic examination. In 1944 Hamman and Rich<sup>6</sup> reported four cases of a diffuse pulmonary disease which was characterized clinically by dyspnea, cyanosis and a rapidly advancing course, pathologically by a marked thickening of the alveolar-capillary septa due to deposition of fibrous tissue. The arterial blood oxygen saturation was determined in two of these patients and found to be markedly reduced. However, no other physio-

\* From the Department of Medicine, Columbia University, College of Physicians and Surgeons, and the Cardio-Pulmonary Laboratory of the First Medical and Chest Services (Columbia University Division), Bellevue Hospital, New York, N. Y. The work described in this paper was supported by a grant from the United States Public Health Fund, with additional support from the Commonwealth Fund and the Life Insurance Medical Research Fund.

† James Alexander Miller Research Fellow.

logic studies were carried out to explain this finding.

Elucidation of the problem was carried further by Baldwin, Cournand and Richards<sup>1</sup> who reported on fourteen patients with chronic pulmonary disease not associated with emphysema and characterized by alveolar-respiratory insufficiency. They were grouped under the heading of pulmonary fibrosis, a descriptive pathologic term which subsequent studies have shown to be inapplicable in certain instances. The diseases included in this group were scleroderma of the lung, sulfur dioxide poisoning, exposure to asbestos fibers, lymphangitic carcinomatous metastases to the lungs and pulmonary granulomatosis and fibrosis of unknown etiology. The clinical and physiologic findings in these cases were essentially the same as those demonstrated by the patients to be reported here. From the physiologic data it was inferred that the mechanics of breathing were not altered, that the distribution of gas in the lungs was not abnormal, and that "alveolar-respiratory insufficiency . . . results both from perfusion of large areas of fibrotic tissue which cannot be ventilated and impairment to the adequate diffusion of respiratory gases across a greatly thickened alveolar septa, or reduction in the area of alveolar capillary interface."

Since these earlier reports there has been appreciable progress in the understanding of the oxygen tension gradient existing between alveolar air and systemic arterial blood. The work of Lilienthal, Riley et al.<sup>7</sup> and its subsequent expansion by Riley, Cournand et al.<sup>8-11</sup> have made it possible to assess separately alteration of the pulmonary membrane and abnormal ventilation-perfusion relationships as factors contributing to alveolar-respiratory insufficiency.

Making partial use of the newer techniques in a few cases of the pulmonary granulomatosis following exposure to beryllium, Wright and his associates<sup>12</sup> interpreted their observations on alveolar-arterial oxygen tension gradients determined while their patients breathed room air at rest and during exercise as evidence of alterations of the pulmonary diffusing membrane. Bruce and others<sup>13</sup> studying the same type of pulmonary granulomatosis also used some of these newer methods and published figures for oxygen diffusion constants, i.e., oxygen diffusing capacity, which are exceedingly low. However, the significance of the latter data is difficult to assess, not only because the calculated oxygen

diffusion constants are inconsistent with the data given for their calculation but also because data obtained while breathing room air cannot be used for such a calculation.

Twelve additional cases which exemplify the essential features, clinical, physiologic and pathologic, of this syndrome of "alveolar-capillary block" are presented here. In these cases all the newer methods for analysis of alveolar-arterial oxygen tension gradient have been employed and in many instances pathologic confirmation of the clinical diagnosis is available. These cases display great variation in the severity and duration of the process, and thus give a fair account of the physiologic basis of the symptoms and signs encountered in the course of this most interesting group of diseases.

#### METHODS

Measurements of lung volumes, maximum breathing capacity, ventilation and gas exchange during rest, standard exercise and recovery were made according to methods previously described from<sup>1</sup> this laboratory. Arterial blood was obtained through an indwelling needle in the brachial artery. Oxygen content and capacity and carbon dioxide content were determined in the Van Slyke-Neill apparatus. Partial pressures of oxygen and of carbon dioxide in systemic arterial blood were obtained by the direct technic of Riley, Proemmel and Franke,<sup>14</sup> and the carbon dioxide tensions so determined were checked by comparison with values derived by application of blood pH and carbon dioxide content to the line charts of Van Slyke and Sendroy. Sampling of mixed venous blood and determination of cardiac output were carried out by methods previously reported.<sup>15</sup> Measurement of blood pressures in the pulmonary artery, right ventricle and brachial artery were recorded with the use of calibrated Hamilton optical manometers.

Analysis of ventilation-perfusion relationships and of the diffusion characteristics of the lungs was made according to the methods devised by Lilienthal, Riley et al.,<sup>7</sup> and extended by Riley, Cournand and others.<sup>8-10</sup> These methods make possible a quantitative expression, in terms now to be defined briefly, of the effectiveness of ventilation and of perfusion of the lungs and of the permeability of the alveolar-capillary interface.

*The Dead Space Ventilation.* This represents the portion of the total ventilation which takes



no part in gas exchange. It is made up not only of the actual volume of gas contained in the anatomic dead space but also of a virtual volume of gas arising from the contributions of all alveoli where ventilation is maintained but circulation is reduced or absent. In this manner total ventilation is divided into two components: dead space ventilation and effective alveolar ventilation. The calculation of dead space ventilation is based upon the relationship between alveolar carbon dioxide tension and the carbon dioxide tension of expired air and upon the assumption that arterial and alveolar carbon dioxide tensions are equal.<sup>16</sup> The dead space ventilation is expressed as a percentage of total ventilation and its value in the normal subject does not exceed 30 per cent.

*The Venous Admixture in the Arterial Blood.* This quantity is made up of an actual volume of mixed venous blood which reaches the peripheral arterial blood by anatomically distinct right to left shunts (bronchial veins, thebesian veins, etc.), and a virtual volume of mixed venous blood, the calculation of which takes into account all the contributions of capillary blood from alveoli in which ventilation is variably reduced in relation to perfusion. This total volume of mixed venous blood is expressed as a percentage of total pulmonary blood flow (cardiac output). Normally this value does not exceed 6 per cent.

*The Oxygen Diffusing Capacity of the Lungs.* This is a measure of the permeability to oxygen of the alveolar-capillary membrane for the lung as a whole. Diffusion of oxygen across this membrane requires a higher partial pressure of oxygen in the alveolar gas than in the capillary blood. This gradient of oxygen pressure between the alveolus and the capillary is largest at the pulmonary arterial end of the pulmonary capillary, becomes smaller as oxygen is transferred from the alveolus to the capillary and is smallest at the pulmonary venous end of the capillary. By methods that have been described before<sup>10</sup> it is possible to calculate a theoretic mean pressure gradient which, were it present continuously along the alveolar-capillary membrane of the entire lung, would account for the same oxygen transfer that is accomplished by the progressively decreasing alveolar-capillary gradient which actually exists. The oxygen intake per minute divided by this calculated mean pressure gradient is the oxygen diffusing capacity. At rest in normal individuals

this value exceeds 15 cc. of oxygen per minute per millimeter of mercury mean pressure gradient.

These last two quantities can be derived from an analysis of the alveolar-arterial oxygen pressure gradient (A-A gradient) at two levels of inspired oxygen concentration. The theoretic considerations upon which this analysis is based, the fundamental assumptions involved and the details of technics and of calculation have been discussed elsewhere.<sup>9,10</sup>

#### CLINICAL AND PHYSIOLOGIC OBSERVATIONS

##### *Clinical Observations*

The vital statistics and diagnoses of the twelve patients are given in Table 1. A brief summary of each case history is also given.

*Etiology.* A variety of diseases have been found to produce the physiologic and clinical pattern reported here. A definite anatomic or etiologic diagnosis could not be made in four of these twelve patients. In the remaining cases the following diagnoses were established by history, biopsy or autopsy: (1) pulmonary granulomatosis after exposure to beryllium in two cases; (2) pulmonary granulomatosis of unknown etiology in one case; (3) Boeck's sarcoid in one case; (4) scleroderma in one case; (5) pulmonary fibrosis of unknown etiology in three cases (in one case after exposure to beryllium; in two cases associated with granulomas in other organs).

Studies on miliary tuberculosis which will be reported elsewhere suggest that during the early and acute phase of this disease a physiologic defect similar to that seen in these cases is present. One patient (Case III) included in this series had a diffuse miliary lesion which cleared completely while streptomycin was being given. Because tubercle bacilli were not found on careful and repeated search, and because significant fever and other toxic manifestations were lacking, a diagnosis of tuberculosis was considered unlikely. However, it is possible that this was a case of chronic or subacute hematogenous pulmonary tuberculosis.

*Pathology.* Examination of the lungs of some of these patients has shown involvement and thickening of the alveolar-capillary septa to be the predominant pathologic lesion. The alveolar-capillary septa were found to be invaded by fibrous tissue or by the cellular elements of granulomas. Involvement of various parts of the



lung has usually not been uniform, and this inhomogeneity may account for some of the physiologic findings. Some have shown areas of focal emphysema. Involvement of the lymph nodes in those cases with granulomatous diseases has suggested that the pulmonary involvement was only part of a generalized disease.

**Laboratory Findings.** The usual clinical laboratory observations were seldom abnormal. Polycythemia was seen only twice and was associated with severe right-sided failure in both of these patients.

**X-ray.** X-ray films of the chest (Figs. 1 to 12) in every case showed involvement of the entire

TABLE I  
PHYSICAL CHARACTERISTICS AND DIAGNOSIS IN TWELVE CASES WITH "ALVEOLAR-CAPILLARY BLOCK"

Case	Age (yr.)	Sex	Weight (kg.)	Height (cm.)	BSA M <sup>2</sup>	Diagnosis	Duration of Illness (mo.)
1. M. B.	34	F	54	161	1.55	Pulmonary granulomatosis (clinical); exposure to beryllium	60
2. M. M.	22	F	47	153	1.42	Scleroderma with pulmonary involvement; (biopsy of skin)	24
3. L. K.	29	M	61	168	1.69	Undiagnosed disease of the lungs	6
4. J. H.	47	F	54	161	1.55	Pulmonary fibrosis of unknown etiology; granuloma of hilar and mediastinal lymph nodes; cor pulmonale; death; (autopsy)	60
5. E. H.	17	M	45	166	1.47	Pulmonary granulomatosis of unknown etiology; (biopsy of lung)	24
6. L. W.	19	F	43	155	1.38	Boeck's sarcoid with pulmonary involvement; (biopsy of lymph node)	60
7. E. L.	47	F	60	167	1.67	Pulmonary fibrosis; Boeck's sarcoid; death; (autopsy)	15
8. F. L.	39	M	70	167	1.79	Pulmonary fibrosis; exposure to beryllium; acute cor pulmonale; terminal bronchopneumonia; death; (autopsy)	24
9. F. S.	43	M	48	165	1.52	Undiagnosed disease of the lungs	42
10. E. M.	60	F	72	151	1.68	Undiagnosed disease of the lungs	11
11. M. H.	64	F	46	150	1.37	Undiagnosed disease of the lungs; cor pulmonale; death; (no autopsy)	84
12. J. Hi.	66	M	55	169	1.63	Pulmonary granulomatosis; exposure to beryllium; spontaneous pneumothorax; death; (autopsy)	Several years?

**Symptomatology.** In most of the patients the disease was insidious in onset and marked by increasing dyspnea on exertion and later at rest. Cough was present in all instances and was productive usually of small amounts of tenacious mucoid sputum. Significant weight loss occurred in many instances. In the recent history it was not unusual to find febrile episodes with pulmonary symptoms.

**Physical Examinations.** The striking physical findings included tachypnea at rest, limited excursion of the thorax, the presence of persistent fine rales heard usually over the lower lobes, and accentuation of the second pulmonic heart sound. Cyanosis significantly was absent or of minimal degree at rest, except in the terminal stages of the disease, but appeared rapidly on mild exertion. Clubbing of the fingers was noted occasionally.

lung field. There was some variation in the size and appearance of the abnormal x-ray shadows which were of the nodular, reticular or conglomerate type. In one case bullae could be identified on the x-ray film. Hilar and mediastinal lymph nodes were enlarged in other cases.

**Clinical Course.** These twelve patients illustrated all possible courses from complete clinical recovery to fatal outcome. Thus in one instance (Case III), under the influence of antibiotic therapy (streptomycin), x-ray and clinical regression took place and has been maintained over a one-and-a-half year period. In another patient (Case II) who was observed and studied during a one and one-half-year period there were no changes in the clinical manifestations of this syndrome. However, the most common course seems to be a steady progression in the symptoms of pulmonary insufficiency, with in-




FIG. 1. Case I. X-ray of M. B.; pulmonary granulomatosis; exposure to beryllium.  
 FIG. 2. Case II. X-ray of M. M.; scleroderma with pulmonary involvement.  
 FIG. 3. Case III. A, x-ray of L. K.; undiagnosed disease of lungs before streptomycin therapy; B, x-ray after streptomycin therapy.  
 FIG. 4. Case IV. X-ray of J. H.; pulmonary fibrosis of unknown etiology.  
 FIG. 5. Case V. X-ray of E. H.; pulmonary granulomatosis of unknown etiology.




FIG. 6. Case vi. X-ray of L. W.; Boeck's sarcoid with pulmonary involvement.  
FIG. 7. Case vii. X-ray of E. L.; pulmonary fibrosis; Boeck's sarcoid.  
FIG. 8. Case viii. X-ray of F. L.; pulmonary fibrosis; exposure to beryllium.  
FIG. 9. Case ix. X-ray of F. S.; undiagnosed disease of the lungs.  
FIG. 10. Case x. X-ray of E. M.; undiagnosed disease of the lungs.  
FIG. 11. Case xi. X-ray of M. H.; undiagnosed disease of the lungs.



creasing dyspnea, cyanosis and inability to exercise. The rate at which these symptoms progress is quite variable, and the interval from the first appearance of symptoms to profound pulmonary insufficiency has varied from a few months to several years. In some of the cases the appearance of irreversible right heart failure has modified the course and hastened a fatal outcome. Acute respiratory infections occurred in some and were controlled by antibiotics. The course of one patient (Case XII) was complicated by the development of a spontaneous pneumothorax.

#### Physiologic Observations

The physiologic data are recorded in Tables II to VI. The characteristic findings were as follows: Measurements of the *lung volumes* demonstrated a significant reduction in the total pulmonary capacity which in general was associated with a parallel reduction in each of its subdivisions. As a result, the ratio of residual volume to total capacity remained normal in all but three instances, and in these the low concentration of alveolar nitrogen after the patient had breathed 99 per cent oxygen for seven minutes




FIG. 12. Case XII. X-ray of J. H.; pulmonary granulomatosis; exposure to beryllium.

and the appearance of the spiograms indicated that no obstructive emphysema was present.

The *maximum breathing capacity* was close to normal in almost all cases and in fact was greater than the predicted value in seven of the twelve.

TABLE II  
LUNG VOLUMES AND MAXIMUM BREATHING CAPACITY IN TWELVE CASES WITH "ALVEOLAR-CAPILLARY BLOCK"

Case	Vital Capacity % Predicted	Residual Volume % Predicted	Total Lung Capacity % Predicted	$\frac{\text{Residual Volume}}{\text{Total L. Capacity}} \times 100$	Maximum Breathing Capacity % Predicted
Normal	100	100	100	<35	100
1. M. B.	73	70	75	19	116
2a. M. M.	33	47	37	33	86
b. M. M.*	25	43	32	35	75
3a. L. K.	63	115	73	25	120
b. L. K.*	85	90	86	21	127
4a. J. H.	42	43	42	26	107
b. J. H.*	36	30	38	20	109
5. E. H.	51	84	58	28	89
6. L. W.	32†	63	38†	44†	67
7. E. L.	37	44	36	32	82
8a. F. L.	51	64	57	28	106
b. F. L.*	49	36	45	20	95
9. F. S.	51	56	51	27	108
10. E. M.	42	100	52	46	94
11. M. H.	26	72	42	44	76
12. J. Hi.	50	70	58	29	108

\* Refers to a second study; M. M. = 18 months later, no change in clinical status; L. K. = 2½ months later, after streptomycin treatment; J. H. = 18 months later, after development of congestive heart failure; F. L. = 1½ months later, after treatment with cortisone, 12 days before death.

† Poor cooperation was obtained in determining the vital capacity.

The presence of constant and at times extreme hyperventilation was demonstrated by studies during rest, standard exercise and recovery. In spite of a large maximum breathing capacity this hyperventilation resulted in an abnormal reduction of the breathing reserve,

nearly equal magnitude. The *oxygen diffusing capacity* was found to be lower than normal under resting conditions in the eleven patients in whom it was measured.

Analysis of *ventilation-perfusion relationships* revealed a significant increase in the degree of

TABLE III  
VENTILATION AND BREATHING RESERVE DURING THE STANDARD EXERCISE TEST AND  
INDEX OF INTRAPULMONARY MIXING IN TWELVE CASES WITH "ALVEOLAR-CAPILLARY BLOCK"

Case	Ventilation			Breathing Reserve		Index of Intrapulmonary Mixing N <sub>2</sub> %
	Rest (L./min./M <sup>2</sup> )	Exercise † (L./min./M <sup>2</sup> )	1st Minute Recovery (L./min./M <sup>2</sup> )	Exercise † (L./min.)	1st Minute Recovery (L./min.)	
Normal	3.4	10.6	12.4	72	70	2.50
1. M. B.	4.3	17.4	22.6	90	73	1.03
2a. M. M.	5.1	16.1	14.4	51	54	1.01
b. M. M.*	6.1	15.5	18.7	43	24	0.90
3a. L. K.	6.2	25.1	21.7	102	100	1.39
b. L. K.*	3.8	26.2	12.5	100	133	1.41
4a. J. H.	5.0	14.7	22.7	59	43	1.29
b. J. H.*	7.8	....	....	...	...	0.87
5. E. H.	7.8	24.5	23.2	66	60	1.00
6. L. W.	7.1	14.7	13.0	37	28	0.96
7. E. L.	7.9	26.9	15.8	21	36	1.43
8a. F. L.	5.9	13.8	16.5	100	96	1.04
b. F. L.*	6.0	18.7	23.1	82	66	1.68
9. F. S.	9.0	30.7	29.5	58	63	1.05
10. E. M.	5.4	17.0	18.1	39	26	1.23
11. M. M.	10.1	....	....	...	...	0.97
12. J. Hi.	9.1	....	....	...	...	2.58

\* Second study as in Table II.

† Standard exercise = Stepping up and down a platform 20 cm. high thirty times in one minute.

especially during exercise and recovery from exercise in all but three patients.

The *arterial oxygen saturation* fell from a normal or minimally reduced value at rest to a distinctly low value during exercise. In every instance the calculated *alveolar oxygen tension* was normal, and in several of the cases exceeded the normal level. The *alveolar-arterial oxygen tension gradient* was increased above normal in all patients when breathing room air at rest but in three of the cases this increase was small and probably insignificant. Of greater importance is the fact that all members of this group had an abnormally large alveolar-arterial oxygen tension gradient when breathing an atmosphere with an oxygen concentration lower than that of ambient air. In most instances the alveolar-arterial oxygen gradients measured at two levels of inspired oxygen concentration were of

calculated *venous admixture* in nine of the eleven cases in whom it was determined. It should be noted, however, that cardiac output was of sufficient magnitude to supply a normal effective pulmonary blood flow, measured in absolute terms, despite the increased fraction of venous admixture.

An increase in *physiologic dead space* was noted in most of the cases. By constant and often marked hyperventilation, normal or increased values for effective alveolar ventilation were maintained even in those cases with a very large physiologic dead space.

Studies of the *hemodynamics* of the pulmonary circulation revealed a mild to marked degree of hypertension in the pulmonary artery in all patients in whom measurements were made during a steady state of mild exercise. The cardiac output at rest was normal or increased

in all cases without congestive failure and was significantly decreased in two cases when it was measured during cardiac decompensation. The increase in cardiac output during mild exercise was in the expected range.

series. Increasing disability from this syndrome does seem to be related to the appearance of arterial oxygen unsaturation at rest, and in the final stages to a return of the usually low arterial carbon dioxide tension at rest to normal or

TABLE IV  
MEASUREMENTS OF GAS EXCHANGE IN TWELVE CASES WITH "ALVEOLAR-CAPILLARY BLOCK"

Case	Oxygen Consumption		Oxygen Consumption per L. Ventilation		Arterial Blood			
	Rest cc./min./M <sup>2</sup>	Exercise* cc./min./M <sup>2</sup>	Rest cc./L.	Exercise* cc./L.	Oxygen Capacity cc./L.	Oxygen Saturation		pCO <sub>2</sub> Rest mm. Hg
						Rest %	1st Min. Recovery %	
Normal	132	494	45.1	55.8	190	98	98	39
1. M. B.	138	486	39.4	34.4	182	98	84	37
2a. M. M.	138	349	25.3	25.8	158	96	79	37
b. M. M.†	114	389	29.0	30.2	118	93	80	38
3a. L. K.	190	604	37.5	34.8	213	96	88	31
b. L. K.†	136	682	43.7	50.5	206	98	96	35
4a. J. H.	121	333	29.3	27.0	146	96	79	37
b. J. H.†	138	...	20.9	....	199	90	..	33
5. E. H.	186	485	29.0	24.1	189	92	86	38
6. L. W.	153	367	25.9	30.0	185	96	74	40
7. E. L.	151	592	27.9	26.3	171	94	81	34
8a. F. L.	133	331	27.5	29.3	200	96	75	33
b. F. L.†	139	421	28.6	27.3	201	90	68	33
9. F. S.	151	373	18.8	14.7	181	91	72	43
10. E. M.	130	358	28.5	20.5	186	92	67	39
11. M. H.	144	...	17.0	....	198	78	..	43
12. J. Hi.	158	...	21.6	....	170	89	..	42

\* See footnote Table III.

† Second study as in Table II.

Observations on the physiologic course of this syndrome are available not only from studies on several patients at different clinical stages of their disease but also from repeated studies of the same patients.

In the tables and case histories the patients have been arranged in order of increasing severity of their disability, as judged from their clinical findings when first seen. This arrangement makes it apparent that the disability is neither related to the reduction of lung volume nor to the maximum breathing capacity. The residual volume/total capacity ratio was increased in some of the patients with the most severe disease. An increase in this ratio has been reported by others<sup>12</sup> in some cases of pulmonary granulomatosis following exposure to beryllium but it was not observed in such patients in this

slightly increased levels. The more advanced cases have in general a large physiologic dead space and venous admixture, and there appears to be some relationship between the severity of the disease and the degree of reduction of the oxygen diffusing capacity.

Four of the twelve cases were studied twice. In one patient with generalized scleroderma (Case II) there was no significant change during an eighteen-month period. During a similar interval in another patient (Case IV) with a granulomatous disease of the lungs there was marked physiologic change with the appearance of arterial oxygen unsaturation at rest, an increase in the A-A pO<sub>2</sub> gradients at rest both while breathing room air and a low oxygen mixture, a further decrease in the oxygen diffusing capacity, and an increase in the venous ad-



mixture and the dead space. In the latter patient irreversible congestive failure had developed, and a reduction in the cardiac output was observed a short time before death. More rapid progression took place in one of the patients (Case VIII) who had been exposed to beryllium phosphors. During a period of 1½ months

percentage of venous admixture. However, it is of interest that there was no change in the degree of pulmonary hypertension after exercise.

## COMMENTS

From these observations it is possible to identify a syndrome, "alveolar-capillary block,"

TABLE V  
MEASUREMENT OF VENTILATION-PERFUSION RELATIONSHIP AND OXYGEN DIFFUSING CAPACITY IN TWELVE CASES WITH "ALVEOLAR-CAPILLARY BLOCK"

Case	Dead Space Ventilation in % Total Ventilation	Effective Alveolar Ventilation (L./min./M <sup>2</sup> )	Alveolar pO <sub>2</sub> Room Air (mm. Hg)	A-A Gradient at Two Levels of Oxygen		Oxygen Diffusing Capacity of the Lung (DL <sub>O2</sub> )	Venous Admixture in % of Cardiac Output
				Room Air	Low O <sub>2</sub>		
Normal	30	2.2	100	12	12	15	6
1. M. B.	27	3.1	114	27	26	10	13
2a. M. M.	26	3.8	109	13	33	5	3
b. M. M.*	33	3.2	105	19	23	5	5
3a. L. K.	26	4.6	121	30	32	10	12
b. L. K.*	15	3.2	116	16	12	17	5
4a. J. H.	38	3.1	120	23	22	9	8
b. J. H.*	46	4.2	112	44	27	6	14
5. E. H.	42	4.5	110	34	30	8	14
6. L. W.	44	4.0	111	34	29	7	13
7. E. L.	46	4.3	114	33	31	7	15
8a. F. L.	46	3.2	102	16	22	7	6
b. F. L.*	33	4.0	114	45	43	..	..
9. F. S.	67	3.0	107	53	42	6	34
10. E. M.	40	3.2	98	16	32	..	..
11. M. H.	57	4.4	98	61†	49	4	25
12. J. Hi.	58	3.8	106	46	43	6	22

\* Second study as in Table II.

† Inspired oxygen concentration 25 per cent.

it was observed that arterial oxygen unsaturation at rest appeared and that there had been a marked increase in the alveolar-arterial oxygen tension gradient both while breathing room air and a lower inspired oxygen concentration.

The physiologic changes which accompany clinical improvement in this syndrome were observed only once (Case III) in the patients reported here. This improvement occurred, as has been previously mentioned, during treatment with streptomycin and was characterized by a slight increase in lung volume; a marked decrease in hyperventilation at rest and during the first minute of recovery from exercise, a decrease in the resting oxygen consumption, the presence of a normal arterial oxygen saturation after exercise and a return to normal of the oxygen diffusing capacity of the lungs and the

which has distinct clinical, pathologic and physiologic features. Most of the findings characteristic of this condition were described previously by Baldwin, Cournand and Richards.<sup>1</sup> The clinical observations made here extend the list of pathologic processes with which the syndrome is associated to include Boeck's sarcoid and the granulomatosis following exposure to beryllium. In the earlier work five cases of Boeck's sarcoid were reported but in all of these cases the only physiologic abnormalities found were characteristic of isolated ventilatory insufficiency and no evidence of interference with the transfer of oxygen to the arterial blood was discovered. The present study makes it clear, however, that in some cases of Boeck's sarcoid with pulmonary involvement there may be an impedance to the diffusion of oxygen. The

reason for these different physiologic patterns is not clear.

It is possible that the lesions in some cases are located predominantly in the alveolar-capillary septa while in others they are chiefly elsewhere. It may also be that although involvement of this

assumption of Wright and others<sup>12</sup> that there is an interference with the diffusion of oxygen across the alveolar membrane in such patients. With the physiologic data available in the work of Baldwin and her associates referred to, the reasons for the defective gas exchange could not

TABLE VI  
HEMODYNAMIC MEASUREMENTS IN TWELVE CASES WITH "ALVEOLAR-CAPILLARY BLOCK"

Case	Pulse Rate	Oxygen Consumption (cc./min./M <sup>2</sup> )	Arterio-Venous Oxygen Difference (cc./L.)	Cardiac Index (L./min./M <sup>2</sup> )	Pulmonary Artery		Systemic Artery	
					s/d (mm. Hg)	m (mm. Hg)	s/d (mm. Hg)	m (mm. Hg)
Normal								
Rest	...	126	39	3.3	20/9	13	112/65	84
Exercise	...	(406)*	(73)	(5.4)	(13/4)	(11)	(125/78)	(100)
1. M. B.	84	133	34	3.9	21/9	15	130/78	103
	(120)	(301)	(59)	(5.1)	(26/9)	(18)	(174/100)	(139)
2a. M. M.	79	133	37	3.6	32/13	20	95/57	74
	(105)	(313)	(60)	(5.2)	.....	(39)	(140/85)	(104)
b. M. M.†	70	141	38	3.7	29/12	19	134/87	106
3a. L. K.	100	185	38	4.9	30/12	19	126/75	93
	(120)	(317)	(49)	(6.5)	(46/13)	(24)	(124/61)	(83)
b. L. K.†	82	174	46	3.8	28/12	18	145/94	114
	(104)	(376)	(59)	(6.4)	(38/15)	(24)	(156/102)	(122)
4a. J. H.	96	115	31	3.7	35/14	23	123/83	100
b. J. H.†	115	196	73	2.7	.....	..	.....	...
5. E. H.	107	196	41	4.8	35/14	23	96/63	75
6. L. W.	110	165	46	3.6	34/15	24	.....	...
7. E. L.	82	149	44	3.4	24/12	17	112/69	84
	(92)	(280)	(56)	(5.0)	(44/21)	(31)	.....	...
8a. F. L.	80	140	49	2.9	30/14	21	108/80	92
9. F. S.	115	159	45	3.5	27/10	17	.....	71
10. E. M.	90	126	39	3.2	58/20	34	121/69	91
11. M. H.	92	144	56	2.6	69/25	43	120/59	81
12. J. Hi.	80	164	34	4.8	.....	..	125/62	87

\* Figures in parentheses represent exercise data obtained during "steady state" exercise.

† Second study as in Table II.

region is always present it is neither severe enough nor sufficiently widespread to interfere with diffusion in the lung as a whole; or the granulomatous lesions are such that they produce an obliteration of both alveolus and capillary, a condition which would produce impedance to diffusion only if it was very extensive. The finding of another case of pulmonary granulomatosis, in which the granulomas contained numerous foreign body giant cells and bi-refractile crystals confirms the earlier observation<sup>1</sup> that a granulomatous lesion of as yet unknown etiology but, with these particular histologic characteristics, may be among the causes of this syndrome. The inclusion of the beryllium cases in this group confirms the

be precisely defined. Although the results of this study lend strong support to the previous inferences that this syndrome is characterized primarily by a disturbance in the diffusion of oxygen across the alveolar membrane, the new information has also revealed that the distinction between diffusion defects and disturbances in the ventilation-perfusion relationships is not absolutely clear-cut in the late or advanced stages of these diseases. Finally, hemodynamic measurements in all patients and complete studies on two different occasions in four patients have served to give a more comprehensive description of this syndrome.

Analysis of the results of physiologic studies yields information about the derangement of

cardiopulmonary function. Reduction of the total pulmonary capacity is not remarkable when one considers the extensive fibrotic or granulomatous involvement of the lungs. The pathologic process apparently reduces the available air spaces. The normal or nearly normal ratio of residual volume to total capacity in all but three instances, together with the normal spiograms, suggests that generalized emphysema is not a part of this pattern; in fact, when pathologic examination demonstrates any emphysema, it is usually of the focal type.

The persistence of a normal maximum breathing capacity with such widespread pulmonary disease is of interest. It implies that the relationship which exists normally between the motive forces produced by the chest muscles to deform the lung and those resulting from the elastic and non-elastic resistances in the lung has not been greatly modified even under conditions of maximum effort. If the anatomic changes in these lungs tend to increase their elastic resistance, as seems likely, and if continuous hyperventilation is present, as it is, hypertrophy of the respiratory muscles or improvement in their coordination must have taken place to account for this well preserved maximum breathing capacity.

Various mechanisms may be involved in the marked hyperventilation at rest and during exercise. In characterizing this syndrome emphasis has been placed on the lack of arterial anoxia at rest in all but the most abnormal cases. Thus anoxia cannot be implicated in the resting hyperventilation and it must be due either to some abnormal reflex stimulus to the respiratory center or to a respiratory center in which chemosensitive cells are more than normally sensitive to carbon dioxide (supranormal centrogenic drive). The infiltrative process in the lungs may well give rise to excessive stimuli via the stretch reflexes and thus explain the initiation of hyperventilation. Once established, hyperventilation leads to lowering of arterial carbon dioxide tension which in turn is followed by a reduction of the alkaline reserve. Both of these have been observed in many of the patients reported here. As has been shown in altitude studies,<sup>17</sup> the existence of a decreased alkaline reserve lowers the threshold of stimulation of the respiratory center to carbon dioxide. Thus a supranormal centrogenic drive may be present and may well act as an additional factor in maintaining hyperventilation. The marked arterial oxygen unsaturation which occurs during exercise will

produce, by its effect on the chemosensitive cells of the carotid body, an additional strong stimulus to ventilation.

The impairment of oxygen transfer from the atmosphere to the blood constitutes the most interesting and important defect observed in this syndrome. The consistently high alveolar oxygen tensions which are present eliminate the possibility of any defect in the transfer of oxygen from atmosphere to alveolus. Of the two causes for the increased oxygen gradient between the alveolus and the arterial blood, namely, abnormal ventilation-perfusion relationships and a reduced oxygen diffusing capacity, the latter is of much greater significance in this syndrome. In contrast to the sometimes normal physiologic dead space and percentage of venous admixture all of these individuals had a reduction in the oxygen diffusing capacity of the lung. The low diffusing capacity may either be due to a reduction in the total area of alveolar membrane which is available for the diffusion of gases, or to a reduction in the permeability of this membrane per unit of area, or to both. The observation of rather widespread thickening of the alveolar capillary septa suggests that the reduction in permeability per unit area is the major reason for the low diffusing capacity. Whether the area of alveolocapillary interface is also reduced under resting conditions cannot be determined.

The observation that many of these patients have a normal resting arterial oxygen saturation while breathing room air at rest is not surprising, even in the presence of a significant reduction of the resting oxygen diffusing capacity. The theoretic considerations developed by Riley and others<sup>9</sup> indicate that in spite of a marked reduction of the oxygen diffusing capacity, the pulmonary capillary blood will still be almost fully saturated if the alveolar oxygen tension is normal. The marked arterial oxygen unsaturation that occurs in these patients after exercise is probably related to a reduced ability to increase the oxygen diffusing capacity of their lungs. Lilienthal, Riley et al.<sup>7</sup> have shown that the increase in oxygen intake that occurs during severe exercise in normal subjects is associated with a six- to seven-fold increase in their oxygen diffusing capacity. It would seem reasonable to assume that this larger  $O_2$  diffusing capacity results from an increase in the area available for gaseous diffusion. The finding of an increase in the pulmonary artery pressure during mild



exercise in all cases in which it was measured suggests that these patients may be unable to increase the area of their pulmonary capillary bed during exercise. Hence such an inability to increase area would prevent the increase in oxygen diffusing capacity necessary to maintain a normal arterial oxygen saturation.

The dead space and venous admixture measurements were abnormal in some of these patients. Such abnormalities usually indicate defective ventilation-perfusion relationships; yet, by other tests these individuals had efficient intrapulmonary gas mixing and alveolar ventilation and no emphysema. Riley et al.<sup>9</sup> point out that by the analysis employed here the only method of distinguishing between distribution and diffusion effects is based on the assumption that alveolar and capillary carbon dioxide tensions are identical at the end of the capillary. They further state that "a small but significant carbon dioxide diffusion gradient exists in . . . cases in which diffusion is severely impaired in all parts of the lungs, but more severely impaired in some parts than others." In such instances the presence of this gradient contributes to the measurements of dead space and venous admixture.

The distinctive characteristics of the pulmonary circulation in this syndrome have been previously discussed briefly.<sup>18</sup> In those patients without cardiac failure an increase in cardiac output was often observed and was associated with a higher than normal oxygen intake. The consistency of these two findings in this group of patients does not seem to be due to the absence of a "steady state" or to the presence of anxiety. These patients appeared as free of anxiety as other patients with different diagnoses who were studied during the same period and who did not show larger than normal oxygen intake. Furthermore, on repeated measurements, with or without a cardiac catheter in place, the oxygen intake was elevated. It is likely therefore that the disseminated and active pathologic process found in many of these cases caused an increased metabolism. When right heart failure developed in the final stages of two of these patients, the low cardiac output was in distinct contrast to what has been described in cases of chronic pulmonary emphysema with anoxia, but similar to what is observed in cases of pulmonary disease with restriction of the pulmonary vascular bed. The elevated pulmonary artery pressure which was noted in nearly every case must be

due to a reduced vascular bed since anoxia is present only during exercise until late in the course of this syndrome. Finally, it must be emphasized that the absence of a continuous anoxic stimulus to the bone marrow probably explains the absence of polycythemia in most of these patients. It was only in the terminal stages, when arterial anoxia of severe degree develops at rest, that a significant increase in red blood cell volume was observed.

#### CASE REPORTS

**CASE I.** M. B., a thirty-four year old white female was admitted to the hospital on February 8, 1949, complaining of shortness of breath of four to five years' duration. Between the ages of eighteen and twenty-two years she was exposed to beryllium phosphors while working in a factory which made fluorescent light bulbs. At about the age of thirty the patient became aware of mild exertional dyspnea and fatigability. A year later a routine x-ray film of her chest was abnormal. Her exertional dyspnea continued but did not progress.

The only significant physical findings were moderate tachypnea, an accentuated second pulmonic sound and slight clubbing of the fingers. There was no cyanosis. X-ray of the chest showed fine nodular infiltration distributed evenly throughout both lung fields. (Fig. 1.)

There was no change in her status while she was in the hospital and she was discharged to her home.

**CASE II.** M. M. was a twenty-two year old colored woman. Two years before hospitalization depigmentation in the left scapular region occurred. A year later changes characteristic of scleroderma were noted in the skin of the face and upper extremities. At this time the patient developed a cough and mild dyspnea on exertion. She was admitted to the hospital on May 24, 1948, and presented the physical findings of well established scleroderma. In addition, thoracic expansion was limited, fine rales were heard posteriorly over both lungs and the second pulmonic sound was accentuated. There was no tachypnea nor clubbing of the fingers.

X-ray of the chest (Fig. 2) showed diffuse fine nodular infiltrates in both lungs. Barium swallow demonstrated delayed emptying of the esophagus.

She was readmitted to the hospital for physiologic studies eighteen months after her first admission. During this interval there had been

no change in her clinical findings. The effect of treatment with cortisone in this patient has been reported elsewhere.<sup>20</sup>

CASE III. L. K. was a twenty-nine year old white Polish man admitted to the hospital on September 20, 1948, complaining of weakness, shortness of breath on exertion and night sweats of four months' duration. Six years prior to admission he had bilateral pneumonia complicated by pleural effusion and pericarditis from which he had recovered. His present illness began insidiously with the development of vague thoracic discomfort, fatigability and occasional elevations of temperature to 99°F. An x-ray of the chest taken four months after the onset of symptoms demonstrated bilateral fine nodular infiltration of the lungs.

Physical examination revealed a well developed, thin patient who did not appear seriously ill. He showed mild tachypnea, slight limitation of chest expansion and fine rales over the lower half of each lung posteriorly. The tuberculin reaction was positive. Tubercle bacilli were never recovered from his sputum. An electrocardiogram shortly after admission showed a prolonged P-R interval. X-ray of the chest showed fine nodular shadows scattered throughout both lung fields. (Fig. 3A.)

The patient was treated with streptomycin, receiving 1 gm. daily for forty-two days. Coincident with therapy the respiratory symptoms decreased, his slightly productive cough disappeared and he gained 18 pounds. These clinical findings were associated with marked clearing of the pulmonary infiltration (Fig. 3B) and with a return of the electrocardiogram to normal. The patient was discharged and has remained well for eighteen months.

CASE IV. J. H., a forty-seven year old colored female, had had four respiratory illnesses in the eight years preceding her first admission, on March 21, 1949. Each was characterized by fever, cough productive of small amounts of mucoid sputum and weight loss. Her chest x-ray was known to have been abnormal for at least four years before her first admission, and dyspnea on exertion had been present for about two years. On physical examination she appeared chronically ill. There was tachypnea, moderate axillary lymph node enlargement, limited chest expansion, rales at the base of both lungs, a greatly accentuated second pulmonary sound and clubbing of the fingers.

X-ray of the chest (Fig. 4) revealed irregular

and coarse infiltration of both lungs. The electrocardiogram was normal. Hematocrit was 39 and blood volume was normal.

Sections of an axillary lymph node showed replacement of lymphoid tissue by numerous discrete and confluent epithelioid tubercles containing large numbers of foreign body type giant cells. Within the giant cells were noted large fragments of doubly refractile crystalline material.

The patient had another admission one and a half years later because of the abrupt onset of fever and an increase in her cough and dyspnea. On physical examination she was acutely ill. Her temperature was 104°F., pulse 120 and respirations 60 per minute. Cyanosis, severe dyspnea, ankle edema and enlargement of the liver were now present in addition to her previous findings. The x-ray of the chest showed no significant change from that of her previous admission but the electrocardiogram now showed evidence of right ventricular hypertrophy.<sup>19</sup>

The white blood count was 14,250, venous pressure 190 mm. saline, decholin circulation time 25 seconds and hematocrit 47 per cent. She was given penicillin, aureomycin, digitalis and mercurial diuretics with a dramatic temporary improvement in her dyspnea and a rapid fall in her temperature. For a brief period after admission her edema cleared, but her dyspnea remained severe enough to confine her to bed. During the four months that she remained in the hospital on this second admission there was a steady increase in her dyspnea, cyanosis and signs of right-sided cardiac failure. For a brief period she also received cortisone, without definite benefit. Progressive cardiorespiratory insufficiency resulted in death.

Autopsy showed widespread pulmonary fibrosis (Fig. 13) involving the alveolar-capillary septa. There was evidence of bronchiectasis at both bases. No granulomatous lesions were found in the lungs but granulomatous lesions containing epithelioid cells, foreign body giant cells and bi-refractile crystals were present in the hilar and mediastinal lymph nodes. These lesions were similar to those found in Case v. Marked right ventricular hypertrophy was present.

CASE V. E. H., a seventeen year old white boy was admitted to the hospital on November 11, 1949, with the chief complaint of weakness. Two years previously he had had "virus pneumonia" marked by a prolonged convalescence. Thereafter the patient was never free of cough.






FIG. 13. Case iv. Microscopic section of lung of J. H.; pulmonary fibrosis of unknown etiology; hematoxylin and eosin stain;  $\times 160$ .



FIG. 14. Case v. Biopsy of lung of E. H.; pulmonary granulomatosis of unknown etiology; hematoxylin and eosin stain;  $\times 160$ .

For a year he had noted increasing exertional dyspnea. Two weeks before admission he had an increase in dyspnea, intermittent pleuritic pain in the left chest, anorexia and low-grade fever. On physical examination the patient appeared poorly nourished. There was tachypnea, fine rales at the lung base, slightly enlarged supraclavicular and axillary lymph nodes and clubbing of the fingers.

X-ray of the chest (Fig. 5) showed a bilateral diffuse granular infiltration. There was also bilateral hilar and mediastinal lymph node enlargement. Biopsies of the lung (Fig. 14), mediastinal and axillary lymph nodes were obtained. The lung biopsy revealed disruption of pulmonary architecture by granulomas in the peribronchial and perivascular connective tissue and in the interalveolar septa. The lesions were discrete, consisting of epithelioid and giant cells of the foreign body type, the latter containing laminated, crystalline, doubly refractile bodies. Mediastinal and axillary lymph nodes showed almost complete replacement of the normal gland structure by similar granulomas. The effect of treatment with cortisone upon the clinical, physiologic and pathologic picture in the patient has been reported separately.<sup>20</sup>

CASE VI. L. W., a nineteen year old colored woman, was admitted to the hospital on September 11, 1950. About five years before admission she became aware of shortness of breath on exertion which slowly but steadily progressed in severity. At that time she also noted the appearance of nodular swellings on her nose. For about two years preceding admission she had a cough productive of small amounts of whitish

sputum and a progressive weight loss which amounted to 17 pounds. On physical examination she appeared chronically ill. There were several discrete, non-ulcerated nodules about the tip of the nose. The lungs were normal. There was an accentuated second pulmonic sound.

X-ray of the chest (Fig. 6) showed a broadened mediastinum and finely nodular densities, closely packed, scattered through both lung fields. X-ray of the hands showed multiple cystic changes in the phalanges. Two cultures and four smears of her sputa failed to show tubercle bacilli. Her skin test to old tuberculin was positive. The albumin/globulin ratio was 4.2/5.0. Biopsies of one of the skin nodules and of a cervical lymph node showed a granulomatous lesion consistent with the diagnosis of Boeck's sarcoid. The electrocardiogram showed evidence of right ventricular hypertrophy.<sup>19</sup>

The effect of treatment with cortisone upon the clinical and physiologic picture in the patient is to be reported at a later date.

CASE VII. E. L., a forty-seven year old colored woman, was admitted to the hospital on December 4, 1948, complaining of cough and weight loss. In the past she had worked as a dispenser of cosmetics. In the year prior to admission she gradually developed a respiratory illness characterized by cough which was at first non-productive, increasing dyspnea and orthopnea and a weight loss of 30 pounds. Six months before admission a chest x-ray was found to be abnormal.

Physical examination revealed a well developed, somewhat undernourished woman who appeared both acutely and chronically ill.






FIG. 15. Case VII. Microscopic section of lung of E. L.; pulmonary fibrosis; hematoxylin and eosin stain;  $\times 160$ .




FIG. 16. Case VIII. Microscopic section of lung of F. L.; pulmonary fibrosis; exposure to beryllium; hematoxylin and eosin stain;  $\times 160$ .

Respirations were rapid and panting in character and bouts of dyspnea were occasioned by paroxysms of cough productive of scant amounts of mucoid, slightly purulent sputum. The anteroposterior diameter of the thorax was increased slightly and expansion of the chest was limited. Breath sounds were everywhere diminished and accompanied by showers of fine rales. There was marked accentuation of the second pulmonic sound.

X-ray of the chest (Fig. 7) showed diffuse finely nodular infiltration of both lungs. During nineteen months in the hospital her dyspnea increased and her cough became productive of sputum which was purulent and at times foul. Therapy with oxygen and antibiotics produced no significant change in her condition. She was transferred to another hospital for chronic care. Death from progressive respiratory failure complicated by hemoptysis occurred about seven months after this transfer. Autopsy showed widespread pulmonary fibrosis involving the alveolar-capillary septa. (Fig. 15.) This fibrosis was of non-specific character, but a few tiny islands of granulomatous lesions characteristic of Boeck's sarcoid were found within it. There was widespread bronchiolectasis in both lower lobes. Granulomatous lesions characteristic of Boeck's sarcoid were present in the liver and spleen.

CASE VIII. F. L. was a thirty-nine year old electrician admitted to the hospital on October 30, 1950. In the five years preceding admission he had broken many fluorescent bulbs and was thus exposed to beryllium phosphors. Two years before admission a chest x-ray was said to show a diffuse pulmonary infiltration. Nine months prior to admission he developed an

acute respiratory illness and a diagnosis of "virus pneumonia" was made. Following this he had a persistent dry cough and exertional dyspnea. Physical examination showed a healthy looking man who was breathing rapidly. There was cyanosis, rales over the lower half of the chest posteriorly and marked clubbing of fingers.

X-ray of the chest revealed a uniform nodular infiltration involving both lung fields. (Fig. 8.) Treatment with cortisone was started. The details of this therapy will be reported at a later date. During seven weeks of hospitalization there was a steady increase in exertional dyspnea. About ten days before his death he complained of chest pain. An increase in his sputum, more cyanosis, fever to as high as  $103^{\circ}\text{F}$ , an elevated white blood count and an increase in the number of rales were noted. Bronchopneumonia was believed to be present and therapy with antibiotics and oxygen was started. Two days before his death a rapid cardiac rate and a diastolic gallop were present. Electrocardiogram showed evidence of acute cor pulmonale. His condition deteriorated rapidly and he died of cardio-respiratory failure about ten months after his first significant symptom and two months after leaving full-time work.

Autopsy showed diffuse pulmonary fibrosis with a marked increase in the thickness of the alveolar-capillary septa. (Fig. 16.) No granulomatous lesions were found in the lungs or lymph nodes. Beryllium could not be identified spectrographically in the lungs but was identified in the hilar lymph nodes. The lungs showed bronchiolectasis in both lower lobes and there was evidence of terminal bronchopneumonia.

CASE IX. F. S. was a forty-three year old

white man who entered the hospital on September 13, 1948, complaining of cough and shortness of breath. Except for a ten-year history of "sinusitis" he had enjoyed good health. In the three years before admission he had developed an increasing and productive cough as well as progressive dyspnea. He had also lost 25 pounds. He gave no history of exposure to toxic agents. Physical examination showed a tachypneic, chronically ill man. The anteroposterior diameter of the chest was slightly increased and expansion of the thorax was limited. Fine rales were present over both lungs. The second pulmonic sound was accentuated. There was moderate clubbing of the fingers and toes.

X-ray of the chest (Fig. 9) showed extensive diffuse reticular infiltration throughout both lungs, more marked on the right. There was no clinical change during hospitalization of two months and he was discharged to his home.

**CASE X.** E. M. was a sixty year old colored woman admitted to the hospital on November 13, 1947. She had been well until a year before hospitalization when she developed exertional dyspnea and a non-productive cough. Physical examination showed only persistent rales at both lung bases.

X-ray of the chest (Fig. 10) showed a linear and patchy infiltration throughout both lung fields but greater on the right where there was also evidence of some pleural fibrosis. There was no significant change in her clinical status during hospitalization for three months and she was discharged to her home.

**CASE XI.** M. H., a sixty-four year old white woman, had had for the twenty years preceding admission a chronic non-productive cough and periodic episodes of acute respiratory illness. Dyspnea on exertion, which increased in its severity progressively, was first noted about seven years before admission. Two months before entering Bellevue Hospital, on March 28, 1950, ankle edema was noted at another hospital. Physiologic studies there revealed pulmonary hypertension, an elevated right ventricular end diastolic pressure, polycythemia and arterial oxygen unsaturation. Digitalization and phlebotomies were followed by loss of edema. Another acute episode led to her entry to Bellevue. On physical examination the positive findings were fever of 101°F., tachypnea, cyanosis, rales over the lower portions of both lungs, an accentuated pulmonic second sound, an enlarged liver and marked clubbing of the fingers. The electro-




FIG. 17. Case XII. Microscopic section of lung of J. Hi.; pulmonary granulomatosis after exposure to beryllium; hematoxylin and eosin stain;  $\times 200$ .

cardiogram showed peaked P waves in lead II and digitalis effect.

X-ray of the chest revealed a widespread nodular infiltration (Fig. 11) throughout both lung fields. On penicillin and aureomycin her temperature returned to normal by the fourth hospital day. Digitalis and mercurial diuretics were also given, with considerable improvement in her dyspnea. She left the hospital after a few weeks and is known to have died at home four months later with symptoms of severe respiratory insufficiency.

**CASE XII.** J. Hi., a sixty-six year old white male electrician, was admitted to the hospital on June 6, 1949, complaining of shortness of breath, weakness and loss of appetite of six weeks' duration. Twenty years previously he had been shot in the left chest and the bullet remained embedded in the chest wall. For nine years, as a result of breaking fluorescent light bulbs, he had been exposed to beryllium phosphors and for several years he had had cough productive of 1 to 3 ounces of clear mucoid sputum daily. This symptom became somewhat worse before admission and was accompanied by slight dyspnea at rest and marked dyspnea on mild exertion. In addition, the patient lost 30 pounds in the six months before hospitalization. On physical examination the patient showed poor nutrition, tachypnea, slight cyanosis, limited chest expansion with an increased anteroposterior diameter, rales at both bases, an accentuated pulmonic second sound and clubbing of the fingers. The electrocardiogram revealed alternating left and right bundle branch block.

X-ray of the chest showed dense homogeneous infiltrations of a reticular type involving the



lower two-thirds of both lung fields, more marked on the right. (Fig. 12.) Two months after study he developed a right spontaneous pneumothorax. Despite re-expansion of the right lung he grew progressively more dyspneic and dependent upon oxygen therapy. He died of respiratory failure five months after study.

At autopsy the lungs were nodular and lacked normal crepitation. There was evidence of marked interstitial thickening with preservation of the air spaces. In addition the upper lobes showed confluent areas of fibrosis obliterating the normal pulmonary architecture. Sections of the lungs showed two types of pathologic processes. At the apices the parenchyma was replaced by proliferating granulation tissue infiltrated with lymphocytes, plasma cells and mononuclear cells and contained Langhans and foreign body giant cells. The few remaining air spaces were reduced to narrow slits. In sections from the basal portions of the lungs the outlines of pulmonary architecture were preserved but the alveolar walls were thickened markedly by similar granulation tissue with resultant reduction of the alveolar air spaces. (Fig. 17.) An additional finding of interest was the presence in all sections of the lungs of marked encroachment upon the lumen of pulmonary arteries by intimal proliferation.

#### SUMMARY

1. Twelve additional cases with various diffuse diseases of the lungs characterized physiologically principally by interference with the diffusion of oxygen across the alveolar-capillary septum have been studied.

2. The patients in this group included two with pulmonary granulomatosis following exposure to beryllium; one with pulmonary granulomatosis of the Boeck's sarcoid type; one with pulmonary granulomatosis of undetermined etiology, in which the granulomatous lesion contained unusually large numbers of foreign body-type giant cells and bi-refractile crystals; one patient with scleroderma; three patients with pulmonary fibrosis of unknown etiology (in one case after exposure to beryllium, in two cases associated with granulomas in other organs), and four cases in which a diagnosis could not be made.

3. The pattern of pulmonary dysfunction consisted of (1) reduced lung volumes, (2) maintenance of a large maximum breathing capacity, (3) hyperventilation at rest and during exer-

cise, (4) normal or nearly normal arterial oxygen saturation at rest but a marked reduction of the arterial oxygen saturation after exercise, (5) normal alveolar oxygen tension, (6) a reduced oxygen diffusing capacity and (7) pulmonary artery hypertension.

4. In some severe cases the dead space-like ventilation and the venous admixture-like perfusion was increased. These findings have been interpreted as an indication of the inhomogeneous nature of the pathologic process.

5. The clinical findings have been analyzed in the light of the physiologic data and the evolutionary trends, both clinical and physiologic, have been described.

6. Because the major pathologic changes are localized in the alveolar capillary septa and because the major physiologic defect is a reduction of the permeability of the alveolar capillary membrane for oxygen, the name "alveolar-capillary block" has been tentatively offered to describe this syndrome.

#### REFERENCES

1. BALDWIN, E. DEF., COURNAND, A. and RICHARDS, D. W., JR. Pulmonary insufficiency. I. Physiological classification, clinical methods of analysis, standard values in normal subjects. *Medicine*, 27: 243, 1948. II. A Study of thirty-nine cases of pulmonary fibrosis. *Medicine*, 28: 1, 1949. III. A study of 122 cases of chronic pulmonary emphysema. *Medicine*, 28: 201, 1949.
2. SCHJERNING, J. Ueber das Problem der Zyanose und den Begriff der Pneumose. *Beitr. z. Klin. Tuberk.*, 50: 96, 1922.
3. BRAUER, L. Die respiratorische Insuffizienz. *Verhandl. d. deutsch. Gesellsch. f. inn. Med.*, 44 Kong., p. 120, 1932.
4. JANSEN, K., KNIPPING, H. W. and STROMBERGER, K. Klinische Untersuchungen ueber Atmung und Blutgase. *Beitr. z. Klin. Tuberk.*, 80: 304, 1932.
5. KNIPPING, H. W. Die Pneumose. *Ergebn. d. inn. Med. u. Kinderh.*, 48: 249, 1935.
6. HAMMAN, L. and RICH, A. R. Acute diffuse interstitial fibrosis of the lungs. *Bull. Johns Hopkins Hosp.*, 74: 177, 1944.
7. LILIENTHAL, J. L., JR., RILEY, R. L., PROEMMEL, D. D. and FRANKE, R. E. An experimental analysis in man of the oxygen pressure gradient from alveolar air to arterial blood during rest and exercise at sea level and at altitude. *Am. J. Physiol.*, 147: 199, 1946.
8. RILEY, R. L. and COURNAND, A. "Ideal" alveolar air and the analysis of ventilation-perfusion relationships in the lungs. *J. Applied Physiol.*, 1: 825, 1948-49.
9. RILEY, R. L. and COURNAND, A. Analysis of factors affecting the concentration of oxygen and carbon dioxide in the gas and blood of the lungs. I. Theory. *J. Applied Physiol.*, in press.



10. RILEY, R. L., DONALD, K. W. and Cournand, A. Analysis of factors affecting the concentration of oxygen and carbon dioxide in the gas and blood of the lungs. II. Methods. *J. Applied Physiol.*, in press.
11. DONALD, K. W., RILEY, R. L., RENZETTI, A. and Cournand, A. Analysis of factors affecting the concentration of oxygen and carbon dioxide in the gas and blood of the lungs. III. Results in patients with pulmonary disease. *J. Applied Physiol.*, in press.
12. WRIGHT, G. W., FILLEY, G. F., PROEMMEL, D. D., PRINCI, F., PLACE, I. and GRINNELL, R. Observations concerning the pathological physiology underlying the disease granulomatosis occurring in beryllium workers. Symposium on Military Physiology. Digest Series No. 4; p. 135, 1947.
13. BRUCE, R. A., LOVEJOY, F. W., JR., BROTHERS, G. B. and VELASQUEZ, T. Observations on the course of dyspnea in chronic pulmonary disease, in chronic pulmonary granulomatosis in beryllium workers. *Am. Rev. Tuberc.*, 59: 364, 1949.
14. RILEY, R. L., PROEMMEL, D. D. and FRANKE, R. E. A direct method for determination of oxygen and carbon dioxide tensions in blood. *J. Biol. Chem.*, 161: 621, 1945.
15. Cournand, A., Baldwin, J. S. and Himmelstein, A. Cardiac Catheterization in Congenital Heart Disease, chap. 1, p. 6. New York, 1949. The Commonwealth Fund.
16. ROSSIER, P. H. and BLICKENSTORFER, E. Espace mort et hyperventilation. *Helvet. med. acta*, 13: 328, 1946.
17. RAHN, H. and OTIS, A. R. Man's respiratory response during and after acclimatization to high altitude. *Am. J. Physiol.*, 157: 445, 1949.
18. HARVEY, R. M., FERRER, M. I., RICHARDS, D. W., JR. and Cournand, A. The influence of chronic pulmonary disease on the heart and circulation. *Am. J. Med.*, 10: 719, 1951.
19. JOHNSON, J. B., FERRER, I. M., WEST, J. R. and Cournand, A. The relation between electrocardiographic evidence of right ventricular hypertrophy and pulmonary arterial pressure in patients with chronic pulmonary disease. *Circulation*, 1: 536, 1950.
20. WEST, J. R., McCLEMENT, J., CARROLL, D. G., BLISS, H. A., KUSCHNER, M., RICHARDS, D. W., JR. and Cournand, A. Effects of cortisone and ACTH in cases of chronic pulmonary disease with impairment of alveolar-capillary diffusion. *Am. J. Med.*, 10: 156, 1951.

# Primary Pulmonary Hypertension\*

## *I. Clinical and Hemodynamic Study*

DAVID T. DRESDALE, M.D., MARTIN SCHULTZ, M.D. and ROBERT J. MIGHTOM, M.D.

*Brooklyn, New York*

PULMONARY hypertension is known to occur in association with left heart failure, chronic pulmonary disease, certain types of congenital heart disease (especially those with increased pulmonary blood flow), diffuse pulmonary embolism, kyphoscoliotic heart disease and specific affections of the pulmonary vascular bed. There is also a primary form of pulmonary hypertension, which is defined as that condition in which an elevated pulmonary artery pressure exists without demonstrable cause. We believe that primary pulmonary hypertension, which has emerged as a distinct entity since the advent of right heart catheterization,<sup>1</sup> is the cause of isolated right ventricular hypertrophy<sup>2-23</sup> whether or not associated with pulmonary vascular sclerosis.

Hypertrophy of the right ventricle in the absence of the previously mentioned disorders was reported by Romberg<sup>24</sup> in 1891. There has been considerable speculation concerning the primary role of pulmonary hypertension<sup>12,16-19</sup> in this condition. Since pulmonary vascular sclerosis was not constantly found, doubt was cast on hypertension in the lesser circulation as the cause of the right ventricular hypertrophy. De Navasquez et al.,<sup>17</sup> for example, preferred to designate their cases as "idiopathic right ventricular hypertrophy" rather than introduce the concept of pulmonary hypertension since no significant changes were found in the pulmonary blood vessels in their cases at necropsy.

Autopsied cases which we consider to fall into the category of primary pulmonary hypertension have been reported under various titles, such as "primary pulmonary vascular sclerosis,"<sup>13,18,21</sup> "right ventricular hypertrophy of unknown origin: so-called pulmonary hypertension,"<sup>17</sup> "isolated hypertrophy of the right ventricle of the heart of unknown cause."<sup>20</sup> These titles

stressed different facets of the pathophysiologic features of this disease. All cases showed marked right ventricular hypertrophy, but the nature, distribution and severity of the pulmonary vascular changes proximal to the capillaries showed considerable variation. The predominant changes were atheromatous lesions of the stem and large elastic arteries, alone or in conjunction with fibrous intimal thickening and narrowing of the smaller arteries and arterioles. Medial hypertrophy was occasionally noted. In about half of the reported cases occlusive lesions of the smaller arteries and arterioles were described. Most of these changes have been ascribed to intimal sclerosis and some to thrombi in various stages of organization. A small group remains in which the pathogenesis of the occlusive lesions cannot be definitely established and may conceivably represent organized pulmonary emboli or healed inflammatory lesions. The last group, of course, would not fit into the category of primary pulmonary hypertension. Whether the obliterative sclerosis or thrombosis, when seen, is the cause or the effect of the hypertension has evoked considerable difference of opinion. Arguments against the primary role of the vascular changes are (1) the absence of these findings in some cases of isolated right ventricular hypertrophy,<sup>6,12,14,16,17,20,22</sup> one of which was proven to have pulmonary hypertension by heart catheterization,<sup>22</sup> and (2) the finding of pulmonary arteriosclerosis unassociated with right ventricular hypertrophy.<sup>13</sup>

Ten<sup>22,25-27</sup> catheterized cases of pulmonary hypertension without demonstrable cause have been mentioned in the literature. Although Nellen<sup>28</sup> reported two cases with hemodynamic studies as "idiopathic pulmonary hypertension," they are not included because of extensive thromboses involving the major pulmonary

\* From the Medical Service, Maimonides Hospital of Brooklyn and the Department of Medicine, State University of New York at New York City, College of Medicine. This investigation was supported in part by a research grant from the National Heart Institute of the National Institutes of Health, Public Health Service.

arteries. In only three of the ten cases were hemodynamic studies reported. The data were limited to pressure determinations in the pulmonary artery or right ventricle or both.

Prior to the introduction of heart catheterization technics this disease was considered to be rare. In 1931 MacCallum<sup>11</sup> reported one case in 12,000 autopsies at the Johns Hopkins Hospital. In 1939 Killingsworth<sup>15</sup> reported one case in 2,707 necropsies at the Children's Memorial Hospital in Chicago. Since heart catheterization has become an accepted procedure, the diagnosis of primary pulmonary hypertension is being made more frequently than such statistics would imply. Wood,<sup>26</sup> in England, studying a group of 233 unselected cases of suspected congenital heart disease, 152 of whom were catheterized, found six cases which he classified as "idiopathic pulmonary hypertension." Chapman<sup>25</sup> in a similar study so diagnosed two of seventy-two patients in whom right heart catheterization was performed. In our experience four cases were encountered over a two-year period at a 500-bed general hospital. Heart catheterization was performed in three of these four patients and the details of these studies will be presented. The clinical course and pathologic findings of the other patient were characteristic of the disease under discussion.

#### CLINICAL PICTURE

Primary pulmonary hypertension has been described in a patient as young as twenty months<sup>21</sup> and in one as old as seventy-four years,<sup>11</sup> the majority of the cases falling in the age group between twenty and forty years. Sixty per cent of the cases reviewed, in which the sex was stated, were females.

Brenner<sup>13</sup> in reviewing this subject stated that the disease appeared to run a course of from five months to five years, with an average course of two years. MacCallum<sup>11</sup> reported a patient who died fifteen years after onset of symptoms. The disease usually runs a progressive downhill course characterized by right heart failure, not infrequently terminating in sudden death.<sup>7,13,15,18,19</sup> The heart failure apparently is not favorably influenced by digitalis.

The salient clinical features in a series of thirty-nine reported cases<sup>2-23</sup> were, in order of frequency, exertional dyspnea and weakness, substernal and left chest pain with exertion resembling angina, syncope on exertion, palpitations, orthopnea and hemoptysis. On physical

examination these patients were described as showing variable degrees of cyanosis without clubbing of the fingers. Orthopnea was rarely noted. The lungs were generally described as clear, except terminally when basal rales have been reported. Percussion of the heart revealed increased substernal dullness and widening of the conus area. The rhythm was regular in all but one case<sup>16</sup> in which transient auricular fibrillation was noted terminally. Systolic murmurs have usually been noted at the apex or pulmonic area. Diastolic murmurs at the pulmonic area have also been mentioned. The second pulmonic sound was greatly accentuated in all cases. Protracted neck-vein distention, elevated venous pressure and hepatomegaly were frequently observed, unassociated with peripheral edema which was usually a late finding.

Except for cyanosis, orthopnea and hemoptysis, the findings in our cases conform to those described in the literature. An impressive feature in two of our cases was the contrast between the appearance of good health when at rest and the striking discomfort evoked by even mild exertion.

Roentgenograms in our patients and in those reviewed show certain characteristic features which, although not specific, were highly suggestive of this disease. There was evidence of right ventricular enlargement. The conus and the pulmonary arterial trunk including its major branches were prominent, whereas the intrapulmonary vascular markings were normal or decreased. There was no evidence of left atrial enlargement. Electrocardiograms were consistent with right ventricular hypertrophy.

Routine examinations of the blood and urine were generally helpful only in that they were within normal limits. Although a slight degree of polycythemia<sup>9,16</sup> has been noted, it was not present in our cases.

#### CASE REPORTS

CASE I. G. N., a twenty-five year old married Czechoslovakian female, was admitted to the Maimonides Hospital on January 10, 1950, because of exertional weakness, dyspnea and effort syncope of six years' duration. Substernal and epigastric pain was provoked by mild exertion which, if continued, would result in syncope lasting two to five minutes. There was no associated incontinence or tongue biting. There was no history of dependent edema, orthopnea, cyanosis, pulmonary disease, systemic hyper-






FIG. 1. Case 1, G. N. (February 16, 1950.) The pattern of right ventricular hypertrophy is shown by the delayed intrinsicoid deflection in  $V_1$  and the tall R waves, depressed S-T segments and inverted T waves in  $V_1$ - $V_4$ .

obese, robust young female who appeared somewhat younger than her stated age. The systemic blood pressure was 120/70, pulse 99, temperature 99°F. She seemed comfortable at rest and there was no cyanosis, clubbing or dependent edema. Fundoscopic examination was normal. The neck veins were not distended and the thyroid was not enlarged. The lungs were clear to percussion and auscultation. The heart was not enlarged to percussion and the heart action was regular. A diastolic shock could be felt over the pulmonic area. The second pulmonic sound was split and greatly accentuated. No murmurs were heard. The liver and spleen were not palpably enlarged. There were good arterial pulsations in the lower extremities.

Laboratory studies: urinalysis, negative; hemoglobin, 13 gm. per cent; red blood cell count, 4.55 million; white blood cell count, 8,000 with a normal differential. Erythrocyte sedimentation rate (Wintrobe) was 14 mm. in one hour. The Mazzini test was negative. Fasting blood sugar was 60 mg. per cent; blood urea nitrogen, 20 mg. per cent. Total proteins were 8.2 gm. per cent, albumin, 4.7 gm. per cent and globulin, 3.5 gm. per cent. The blood cholesterol was 212 mg. per cent with 24 per cent free; blood




FIG. 2. Case 1, G. N. A, showing a prominent pulmonary artery segment and enlarged hilar vascular shadows with clear lung fields; B, the barium-filled esophagus in the right anterior oblique view shows no posterior displacement; C, diffuse dilatation of the pulmonary artery is noted in the lateral view.

tension or thromboembolic phenomena. She had been treated at various clinics, European and American, for these complaints without improvement. There was no past history of rheumatic fever, chorea, scarlet fever or nephritis. The family history was non-contributory, except for a twin sister who died in infancy of a "cold."

Physical examination revealed a short, slightly

calcium, 7.0 mg. per cent; phosphorus, 3.6 mg. per cent. The cephalin flocculation test was negative. Venous pressure on two separate occasions was 100 and 110 mm. of saline and the decholin circulation time was nineteen seconds. The electrocardiogram (Fig. 1) was consistent with right ventricular and right atrial hypertrophy. Chest x-rays are shown in Figure 2A to 2C.

The patient's course in the hospital was uneventful. Exercise tolerance was self-limited to three to four successive stair steps. The diagnosis of primary pulmonary hypertension was suggested. Because of the x-ray findings and limited exercise tolerance, heart catheterization was performed.

CASE II. M. R., a 35 year old Puerto Rican housewife, was admitted for the first time to the Maimonides Hospital on April 17, 1950, because of pharyngitis. She was apparently well until the delivery of her fifth child four years prior to this admission. As far as could be determined the first record of her ante-partum status was made during the seventh month of this pregnancy. Blood pressure was 120/88 and no heart murmurs were described. Urinalysis was negative. The patient complained of some exertional dyspnea at this time. Two months later prior to the onset of labor blood pressure was 138/98 and a "to and fro" murmur was heard at the apex. Following the spontaneous delivery of a normal male infant and before the expulsion of the placenta she had a generalized convulsive seizure which lasted one and one-half minutes. The sensorium was clouded for the ensuing fifteen minutes. Blood pressure immediately following the convulsion was 90/70 and the pulse was 160 beats per minute. There was no chest pain or hemoptysis. Recovery was rapid and uneventful. About two weeks after delivery the patient noted swelling of the face, abdomen and legs without respiratory difficulties. She was admitted to another hospital five months later for these complaints. She was digitalized and given weekly mercurial diuretics. Two and one-half years later progressive exertional dyspnea and orthopnea appeared. Fleeting chest pain on exertion was noted. She had no syncope, chronic cough, hemoptysis or thromboembolic phenomena. There was no past history of rheumatic fever, chorea, scarlet fever, nephritis or toxemia of pregnancy.

Physical examination disclosed a well developed and well nourished female in no distress. The systemic blood pressure was 165/85, pulse 80, temperature 99°F. There was evidence of mild pharyngitis. The retinal vessels showed minimal sclerotic changes. The neck veins were distended and pulsating. The lungs were clear to percussion and auscultation. The point of maximal impact of the heart was in the sixth intercostal space, 2 cm. outside the mid-clavicular line. There was a pre-systolic apical thrill

and a grade III, harsh, apical diastolic blow which was widely transmitted and well heard at the lower sternum. A grade III, apical systolic murmur was widely transmitted and heard equally well over the lower sternum. The pulmonic second sound was accentuated and was louder than the aortic second sound. A slightly tender liver edge was felt 6 cm. below the costal margin. There was no cyanosis, clubbing or edema. The heart action was regular.

Laboratory studies revealed: urinalysis, negative except for an occasional trace of albumin; hemoglobin, 14.5 gm. per cent; red blood cell count, 4.69 million; white blood cell count, 16,000 with a shift to the left initially, returning to normal. The erythrocyte sedimentation rate (Wintrobe) was 4 mm. per hour. Blood chemistries showed: fasting blood sugar, 65 mg. per cent; blood urea nitrogen, 13 mg. per cent; total proteins, 6.9 gm. per cent, albumin, 3.6 gm. per cent, globulin, 3.3 gm. per cent; total cholesterol, 161 mg. per cent with 25 per cent free; blood calcium, 9.8 mg. per cent; cephalin flocculation test, negative; thymol turbidity, 2.8 units; blood sodium, 142.8 mEq./L.; chlorides, 105.5 mEq./L. and potassium, 5.1 mEq./L. The blood Mazzini test was negative. The venous pressure was 145 mm. of saline with a positive hepatojugular reflux. The decholin circulation time was thirty-one seconds and the ether circulation time was eleven seconds. The electrocardiogram (Fig. 3) was consistent with right ventricular hypertrophy. Chest x-rays are shown in Figures 4A to 4C.

The pharyngitis responded promptly to penicillin. The patient improved on bed rest, salt poor diet, digitoxin and mercurial diuretics. Nine days after admission the venous pressure was 110 mm. of saline. The diastolic murmur heard initially became inaudible during the third week of hospitalization. Because of the unusual x-ray findings, changing murmurs and lack of left atrial enlargement she was referred to the cardiopulmonary laboratory for heart catheterization. As a result of these studies and the clinical findings this patient was thought to have rheumatic heart disease with mitral stenosis and insufficiency and a pulmonary endarteritis. She was then discharged on a cardiac regimen.

The patient was readmitted to the Maimonides Hospital three months later because of increasing heart failure responding poorly to digitoxin and mercurial diuretics. Dyspnea became more marked during the two days prior to admission.






FIG. 3. Case II, M. R. (May 12, 1950.) There is a Q wave and delay of the intrinsicoid deflection in  $V_1$  and an R wave in  $aVr$ .

hepatojugular reflux. The lungs were clear to percussion and auscultation. The heart action was irregular with a pulse deficit, the radial rate being 56 beats per minute. No diastolic murmurs were heard. The liver edge was felt four finger breadths below the costal margin. There was three plus ankle edema extending to the mid-calf. The nail beds were dusky.

Laboratory studies were essentially unchanged. The white blood cell count was 12,200. The electrocardiogram showed ventricular premature contractions in addition to the signs of right ventricular hypertrophy.

The patient was placed on bed rest, a salt-free diet, mercurial diuretics and digitoxin. Digitoxin was discontinued after the second dose because of the onset of bigeminal rhythm and vomiting. The venous pressure was 242 mm. of saline with a positive hepatojugular reflux. The decholin circulation time was thirty-seven seconds. Following estimation of the ether circulation time the patient developed nausea. The following day the patient's condition deteriorated with the onset of retching, feeble pulse and a sinus tachycardia of 120 beats per minute. There was no dyspnea or orthopnea. Five hours later she was pulseless, cool, cyanotic and the blood pressure




FIG. 4. Case II, M. R. A, showing enlargement of the heart to the left and inferiorly with rounding of the cardiac apex above the diaphragm and increased convexity of the pulmonary artery segment. The hilar vessels are enlarged whereas the intrapulmonary vascular markings are diminished; B, enlargement of the right ventricle but there is no displacement of the barium-filled esophagus in the right anterior oblique view; C, the left ventricle overlaps the spine in the left anterior oblique view. The left ventricle was normal at necropsy.

There was no cough, chest pain or hemoptysis. On physical examination the patient appeared in no acute distress. Blood pressure was 130/95 and the temperature 99.2°F. The neck veins were distended and pulsating with a positive

was unobtainable. A portable chest film showed the lung fields to be clear. She was placed in an oxygen tent and given an infusion containing neosynephrine without demonstrable effect. The cyanosis was relieved by oxygen adminis-






FIG. 5. Case II, M. R. The right atrium is dilated. Marked dilatation and hypertrophy of the right ventricle with scattered patches of endocardial thickening is seen.

FIG. 6. Case II, M. R. The left atrium, left ventricle and mitral valve are normal.

tered either by tent or mask. She failed to respond to therapy and died sixteen hours after the onset of cardiovascular collapse.

Autopsy\* was performed sixteen hours after death through a limited abdominal incision. The pericardial cavity and its lining were normal. The heart weighed 330 gm. The right atrium and ventricle were dilated. The right atrium was thin whereas the right ventricle was markedly hypertrophied (Fig. 5) and measured 0.9 cm. in thickness. The left atrium and ventricle (Fig. 6) were normal. The left ventricle measured 1.4 cm. in thickness. The interventricular septum was hypertrophied and bulged somewhat into the left ventricle. The valvular circumferences were as follows: tricuspid, 11.5 cm.; pulmonic, 7.0 cm.; mitral, 8.0 cm. and aortic, 6.0 cm. There were no abnormalities of the valve leaflets or cusps. The coronary arteries including their ostia were normal. The pulmonary artery and its major branches were dilated and showed widespread atheromatous plaque formation. The aorta was normal. No congenital malformations were found.

A small amount of fluid was present in each pleural space. The pleural surfaces were smooth and glistening. The lungs appeared normal except for some atelectasis of the left lower lobe. The right lung weighed 310 gm. and the left lung weighed 335 gm. No thrombi or emboli were noted in the pulmonary vessels.

The liver weighed 1,240 gm. and aside from

\* We are indebted to Dr. A. Kantrowitz, Director of the Pathological Laboratories, Maimonides Hospital of Brooklyn, for his kind assistance in the preparation of the autopsy report.




FIG. 7. Case II, M. R. Marked intimal sclerosis of a small intrapulmonary artery is noted;  $\times 300$ .

a characteristic "nutmeg" appearance presented no other abnormalities. No other essential organ changes were noted.

Histologically, sections of the heart showed hypertrophy of the muscle fibers of the right ventricle. There was no evidence of myocardial fibrosis or valvular disease.

Multiple sections from various portions of the lungs were examined. Widespread sclerotic changes were noted in the pulmonary arterial tree, varying in degree even in vessels of similar caliber. Plaque formation with intimal thickening as well as increase in the width of the media were noted in the largest vessels. Subintimal thickening and proliferation were the prominent changes in the vessels of microscopic size. (Fig. 7.) In many such vessels the lumen was extremely narrow, at times slit-like. In addition to the above changes there were thrombotic lesions in various stages of organization. They ranged from recent thrombi (Fig. 8A) to re-




FIG. 8. Case II, M. R. A, a thrombus of recent origin in a pulmonary vessel of microscopic size is seen;  $\times 150$ . B, a recanalized thrombus in a pulmonary vessel of microscopic size is seen;  $\times 250$ .

canalized thrombi (Fig. 8B) in the precapillary and post-capillary vessels of microscopic size. These latter changes were much less numerous than the sclerotic changes previously described. All sections revealed normal vessels although those affected seemed to be present in preponderant number.

The bronchi and peribronchial tissues showed no evidence of chronic inflammatory changes. The alveoli revealed no morphologic changes and no "heart failure" cells were seen. The alveolar septal interstitial tissue was not thickened. Sections from the basilar portion of the right lower lobe revealed a slight amount of intra-alveolar edema fluid. Slight atelectasis was noted in sections from the left lower lobe.

Sections of the liver and spleen revealed only marked chronic congestive changes. The remainder of the microscopic examination was normal.

The anatomic diagnoses were: (1) Severe pulmonary arteriosclerosis and arteriolarsclerosis; (2) hypertrophy and dilatation of the right ventricle; (3) old and recent thromboses of the pulmonary arterioles and venules; (4) chronic passive congestion of the liver and spleen; (5) moderate edema of lower lobe of right lung and (6), atelectasis of lower lobe of left lung.

In summary, the autopsy revealed right ventricular hypertrophy and widespread pulmonary vascular sclerosis unassociated with intrinsic lung disease. A preponderance of sclerotic changes over thrombotic lesions in the small pulmonary vessels was evident.

CASE III. M. K., a thirty-four year old white housewife, was admitted to the Maimonides

Hospital on July 17, 1950, because of weakness, exertional dyspnea and numerous syncopal episodes of two years' duration. Syncope occurred only during exertion and was frequently preceded by great weakness. It was never accompanied by convulsions or tongue biting. Hyperventilation was denied. There was no history of orthopnea or cyanosis. A complete medical study including chest x-ray, electrocardiogram, electroencephalogram and blood studies at another institution was reported as normal. The patient was then advised to seek psychiatric aid. Intermittent ankle edema, epigastric pressure on exertion, transient nausea, vomiting and diarrhea appeared two months before admission. One day prior to admission she lost consciousness and incurred a contusion of the right eye. Her past history was non-contributory. The patient has two normal children, ages three and eight years, respectively. Pregnancies were uneventful.

Physical examination disclosed a well developed young woman in no distress. The systemic blood pressure was 120/95, pulse 80, temperature 98°F. There was an ecchymosis over the right eye. Dyspnea, orthopnea, cyanosis, clubbing and edema were absent. The neck veins were moderately distended. The lungs were clear to percussion and auscultation. The point of maximal impact of the heart was in the fifth intercostal space at the mid-clavicular line and a pulsatory, heaving impulse was noted in the third intercostal space at the left of the sternum. Retrocardiac dullness was increased. The heart action was regular and the heart sounds were of good quality. The pulmonic second sound was greatly accentuated and was



louder than the aortic second sound. No murmurs were heard. A firm, non-tender liver edge was palpable at the level of the umbilicus.

Laboratory studies showed: urinalysis, negative; hemoglobin, 14.5 gm. per cent; red cell count, 5.1 million; white cell count, 10,100 with a normal differential. Blood chemistries revealed: fasting blood sugar, 75 mg. per cent; blood urea nitrogen, 13 mg. per cent; serum albumin, 4.0 gm. per cent; serum globulin, 2.3 gm. per cent; alkaline phosphatase, 5.8 King-Armstrong units; thymol turbidity, 0.2 units; icteric index, 7.9. The blood Mazzini test was negative. The venous pressure was 200 mm. of saline with a positive hepatjugular reflux. The ether circulation time was four seconds and the decholin circulation time was twenty-five seconds. The electrocardiogram (Fig. 9) revealed a pattern consistent with right atrial and right ventricular hypertrophy. Chest x-rays are shown in Figures 10A to 10c.

In view of the presenting symptoms and physical findings the diagnosis of primary pulmonary hypertension was made. The patient was then referred to the cardiopulmonary laboratory for confirmation of this diagnosis and for hemodynamic studies. During cardiac catheterization the patient was digitalized with lanatoside C. Subsequently she was placed on a maintenance dose of digitoxin and discharged on August 1, 1950. Despite a strict cardiac regimen there was no obvious improvement in her clinical status. She was readmitted two




FIG. 9. Case III, M. K. (July 25, 1950.) There are peaked P waves in leads II and III. The R wave in aVr is broad and the R wave in V<sub>1</sub> is high. The T waves are inverted in V<sub>1-3</sub>. This pattern is consistent with right atrial and right ventricular hypertrophy.

months later for re-evaluation and pharmacodynamic studies. Physical examination was essentially as on the previous admission.




FIG. 10. Case III, M. K. A, showing enlargement of the heart to the left with rounding and elevation of the cardiac apex above the diaphragm. The pulmonary artery segment and hilar shadows are prominent whereas the intrapulmonary vascular markings are normal; B, there is no displacement of the barium-filled esophagus in the right anterior oblique view. The outflow tract of the right ventricle is prominent; C, the left ventricle overlaps the spine in the left anterior oblique view.



Laboratory studies showed: albumin, 4.7 gm. per cent; globulin, 3.9 gm. per cent; total cholesterol, 195 mg. per cent with 20 per cent free; serum calcium, 9.8 mg. per cent; serum phosphorus, 4.3 mg. per cent; serum potassium, 5.8 mEq./L.; serum chlorides, 101 mEq./L.; serum sodium, 145.3 mEq./L.; serum carbon

maintenance dose of digitoxin and oral prisco-line. She is being followed at the cardiopulmonary laboratory.

## HEMODYNAMIC STUDIES

The physiologic data presented in Tables I and II were obtained at the time of right heart

TABLE I  
DATA OBTAINED DURING RIGHT HEART CATHETERIZATION

Case No.	Name	Sex	Age	Date	Surface Area (M <sup>2</sup> )	Arterial Blood			O <sub>2</sub> Consumption (cc./min.)	BMR (%)	Heart Rate (beat/min.)	Arteriovenous O <sub>2</sub> Difference (vol. %)	Cardiac Output (L./min.)	Cardiac Index (L./min./M <sup>2</sup> )	Stroke Volume (cc.)	Blood Volume		
						O <sub>2</sub> Content (vol. %)	O <sub>2</sub> Capacity (vol. %)	Saturation (%)								TBV* (cc./M <sup>2</sup> BSA)	PV† (cc./M <sup>2</sup> BSA)	Hematocrit (%)
I	G. N.	F	25	1-20-50	1.52	18.0	18.4	97.8	217	+12	78	5.9	3.72	2.44	48	2,710	1,615	40
II	M. R.	F	35	5-12-50	1.41	17.7	18.0	98.4	179	+3	94	8.3	2.16	1.53	23	3,550	1,882	47
III	M. K.	F	34	10-10-50	1.71	18.9	19.6	96.4	189	-14	93	6.9	2.72	1.59	29	2,615	1,555	40.5

\* TBV = Total blood volume

† PV = Plasma volume

Predicted values (female):

Total blood volume	Plasma volume	Hematocrit
2,670	1,600	40

TABLE II  
INTRACARDIAC AND PERIPHERAL BLOOD PRESSURES, RESISTANCES IN PULMONARY AND PERIPHERAL CIRCULATION

Case	Cardiac Output (L./min.)	Peripheral Artery			Systemic Resistance (dynes/cm. <sup>5</sup> /sec.)		Pulmonary Artery			Pulmonary Resistance (dynes/cm. <sup>5</sup> /sec.)		Right Ventricle		Right Atrium
		Systolic	Diastolic	Mean	Predicted †	Observed	Systolic	Diastolic	Mean	Predicted †	Observed	Systolic	Diastolic	Mean
		(mm. Hg)					(mm. Hg)					(mm. Hg)		
G. N.	3.72	117*	79	98	1,440	2,100	121	47	72	221	1,550	121	8	7
M. R.	2.16	135†	91	106	1,560	3,930	73	41	53	239	1,965	75	18	12
M. K.	2.72	113*	80	92	1,280	2,710	90	46	62	197	1,825	84	14	13

\* Brachial artery.

† Femoral artery.

‡ See formulas on page 695.

dioxide, 31.5 mEq./L. The bromsulfalein test revealed 15 per cent retention after forty-five minutes. The congo red test was negative.

Following heart catheterization the patient was discharged on October 23, 1950, on a

catheterization. The patients were studied in a resting state after fasting for at least twelve hours. They received oral phenobarbital 0.06 gm. and nembutal 0.1 gm. the evening before the procedure and phenobarbital 0.06 gm. two

to three hours before any observations were made. Two of the patients, M. K. and M. R., were fully digitalized at the time of the reported studies. M. K. had been digitalized for three months and M. R. for over three years because of earlier clinical findings consistent with congestive heart failure.

The blood oxygen content and capacity were determined by the method of Van Slyke and Neill.<sup>29</sup> The blood volumes were determined by the direct T-1824 (Evans blue) dye method, as modified by Noble and Gregersen.<sup>30</sup> The cardiac outputs and the cardiac indices were determined by the direct Fick principle.<sup>31</sup> Peripheral and intracardiac pressures were recorded using Sanborn electromanometers and a direct writing Sanborn Poly-Viso recorder. The pressures in the pulmonary artery and right ventricle were not recorded simultaneously. Vascular resistances in the pulmonary and systemic circuits were calculated according to the formulas:

$$R. (\text{pulmonary}) = \frac{P.A.m \times 1.332 \times 60}{C.O.}$$

$$R. (\text{peripheral}) = \frac{B.A.m \times 1.332 \times 60}{C.O.}$$

where R. = vascular resistance in dynes/cm.<sup>5</sup>/sec.

P.A.m = mean pressure in the pulmonary artery in mm. Hg

B.A.m = mean pressure in the brachial artery in mm. Hg

C.O. = cardiac output in L. per minute

The predicted resistances were obtained using normal mean pressures and blood flows, as reported by Cournand,<sup>32</sup> and corrected for body surface area. The mean pressures were determined by planimetry of the recorded tracings using one or more complete respiratory cycles. The point of zero reference for measurement of intracardiac pressures was 5 cm. below the angle of Louis. The expired air was collected in a Tissot spirometer and was analyzed in the Scholander<sup>33</sup> gas analyzer. The basal metabolic rates were calculated from the observed gas exchange at the time of cardiac output determinations. Pulmonary function studies were performed at a time separate from cardiac catheterization using the technics described by Baldwin et al.<sup>34</sup>

**Arterial Blood Oxygen Saturation.** The resting arterial blood oxygen saturation was normal in all of the patients. Immediately after maximal




FIG. 11. Case 1, G. N., primary pulmonary hypertension. The brachial and pulmonary artery pressures were taken simultaneously. Continuous tracings during withdrawal of the catheter from the pulmonary artery to the right ventricle showed no change in systolic pressure.

exercise there was no change in G. N. and a very slight decrease in M. K. An exercise sample was not obtained in the third patient.

**Blood Volume.** The total blood volume and hematocrits were normal in G. N. and M. K. and increased in M. R. The total blood volume in M. R. was increased 33 per cent and the hematocrit was 47.

**Blood Flow.** The cardiac index was low in all of the patients. It was highest but still below the average normal in G. N. The arteriovenous oxygen difference was increased and the stroke volume was decreased in all of the patients.

**Arterial and Right Heart Pressures.** The peripheral arterial pressures were normal at the time of cardiac catheterization. The pulmonary arterial pressures were markedly elevated. In G. N. the pulmonary systolic pressure was slightly higher than the brachial artery systolic pressure during a simultaneous recording. (Fig. 11.) The right ventricular end diastolic pressures and right atrial pressures were elevated above the accepted normal<sup>35</sup> in all of the patients.

**Systemic and Pulmonary Resistances.** Pulmonary resistances, calculated from the observed values, averaged eight times the predicted values

whereas the systemic resistances similarly calculated were only twice those predicted. (Table II and Fig. 12.)

**Pulmonary Function Studies.** Pulmonary function studies were carried out on two of the three patients. (Table III.) The ventilatory capacity

was normal in M. K. and reduced in G. N. Oxygen consumption was reduced at rest in M. K. and the rise during exercise was less than expected in both. There was no rise in the oxygen consumption per liter of ventilation during exercise.

TABLE III  
PULMONARY FUNCTION AND ARTERIAL OXYGEN SATURATION STUDIES IN TWO PATIENTS WITH PRIMARY PULMONARY HYPERTENSION

Case	Vital Capacity (% of predicted)	Maximal Breathing Capacity (% of predicted)	Ventilation (L./min./M <sup>2</sup> BSA)		Breathing Reserve (%)			Oxygen Consumption (cc./min./M <sup>2</sup> BSA)		Respiratory Quotient		Oxygen Intake (cc./L. vent.)		Arterial Oxygen Saturation (%)	
			Rest	Exercise†	Rest	Exercise	1st Min. Recovery	Rest	Exercise	Rest	Exercise	Rest	Exercise	Rest	Exercise
G. N.	62	72	4.0	9.2	91	77	74	136	352	.704	.774	34	34	96.8	97.5
M. K.	102	101	2.8	22.1	95	61	78	97	399	.765	.941	35	18	96.2	93.2‡
Predicted*	100	100	3.2	9.0	93	..	79	126	463	.776	.678	45	60	96.2	95.8

\* Baldwin et al.<sup>34</sup>

† Exercise—G. N., 23 steps in 1.5 min.

M. K., 24 steps in 1.1 min.

‡ Maximal exercise, supine, during cardiac catheterization.




FIG. 12. Peripheral and pulmonary resistance in primary pulmonary hypertension. Calculated pulmonary resistances are seven to nine times those predicted (see procedure).






FIG. 13. Case II, M. R. Note the pulmonary hypertension and the elevated right ventricular end diastolic pressure. (The tracings were inked over for photographic purposes.)

#### COMMENTS

Pulmonary hypertension of considerable magnitude was observed in our three patients. (Table II, Figs. 11, 13 and 14.) Secondary pulmonary hypertension as defined in the introduction was ruled out by clinical and laboratory studies in G. N. and M. K. and by postmortem examination in M. R.

Although the clinical picture is fairly characteristic in patients with primary pulmonary hypertension, the diagnosis has rarely been made during life. Brill and Krygier<sup>18</sup> in their review of the cases up to 1941 noted that the ante-mortem diagnosis was made in only two of twenty cases. Mention has been made of the current increasing frequency of diagnosis. However, in those patients who do not present exertional syncope, as emphasized by Dressler,<sup>40</sup> or angina, or both, the diagnosis may be obscure. It becomes even more difficult in this group when murmurs are heard ordinarily associated with mitral stenosis. Such was the case in M. R. in whom the diagnosis was not made until necropsy despite cardiac catheterization.

Primary pulmonary hypertension should not be confused with so-called Ayerza's disease, which has never been clearly defined but which is often thought of in connection with pulmonary vascular sclerosis. The patients frequently put into Ayerza's group<sup>36-38</sup> are deeply cyanotic because of the underlying lung disturbance. Although many of the cases falling into the category of primary pulmonary hypertension or primary pulmonary vascular sclerosis have been described as cyanotic<sup>13,16-18</sup> no arterial blood oxygen studies were mentioned. Brill and Kry-




FIG. 14. Case III, M. K. Pulmonary hypertension and elevated right ventricular end diastolic and mean right atrial pressures are observed.

gier<sup>18</sup> went so far as to state that the disproportion of cyanosis to dyspnea differentiated this syndrome from cor pulmonale secondary to lung disease. Our patients were not clinically cyanotic and arterial blood oxygen studies showed normal saturations. Stagnant anoxia due to decreased blood flow could explain the cyanosis so frequently described, especially during the terminal phase of this disease. Another possible cause for cyanosis when seen in primary pulmonary hypertension could be a complicating patent foramen ovale. This could have contributed to the cyanosis in the patient reported by Brill and Krygier. The mechanism of cyanosis in primary pulmonary hypertension with patent foramen ovale would be similar to that described by Engle and Taussig<sup>39</sup> in pulmonic stenosis with patent foramen ovale.

Unexplained dyspnea and weakness, especially if associated with exertional syncope<sup>40</sup> or angina in an otherwise healthy-looking individual with an accentuated pulmonic second sound, should suggest primary pulmonary hypertension. Syncopal attacks, described in 20 per cent<sup>11,13,15,17,18,22,26</sup> of the thirty-nine cases reviewed,<sup>2-23</sup> was an early and presenting complaint in two of our patients. It is interesting to note that the syncopal attacks were precipitated




FIG. 15. Case III, M. K. (July 25, 1950.) Effect of exercise on right ventricular pressures; (A) before lanatoside C; (B) seventy-five minutes after the intravenous administration of 1.6 mg. of lanatoside C. There may be a slight error in the pressure tracings during exercise because of possible artifacts introduced during vigorous physical exertion.

by exertion or sudden movements, especially if associated with emotional tension. In M. K. the syncope was preceded by a feeling of weakness while in G. N. the syncope was preceded by marked pressure under the lower part of her sternum. Syncope is not generally associated with secondary pulmonary hypertension.

The mechanism for syncope in primary pulmonary hypertension has not been established. Our patients did not have sensitive carotid sinus reflexes. Arterial anoxemia was not observed with exercise and therefore can be eliminated as a possible cause for exertional syncope. M. K., who had numerous syncopal episodes, was exercised in the supine position during right heart catheterization,\* both prior to digitalization and seventy-five minutes after the intravenous administration of 1.6 mg. of lanatoside C. Syncope was not precipitated. During exercise at both times the right ventricular end diastolic pressure increased considerably above the control values as seen in Figures 15A and B, while the peripheral pressure remained essentially unchanged. What effect changing intrapleural pressure during exercise had on the right ventricular end diastolic pressure cannot be stated since the intrapleural pressure was not measured. A grunting type of respiration was not observed during exercise. One can but speculate whether additional exercise would have precipitated a further rise in the right ventricular end diastolic pressure and acute

\* Studies obtained during first cardiac catheterization, July 25, 1950.

right heart insufficiency with resultant inadequate filling of the left ventricle.

Since, as is well known, coronary blood flow is greatest during diastole, it is possible that the marked rise observed in right ventricular end diastolic pressure might impede the coronary circulation. This could account for the symptom of angina and could precipitate ventricular fibrillation or cardiac standstill<sup>41</sup> with resultant syncope. Although hemodynamic studies were not performed, Brill and Krygier<sup>18</sup> suggested that relative coronary insufficiency accounted for the chest pain and was the cause of death in those patients with this syndrome who died suddenly with but slight evidence of congestive heart failure. Diminished coronary flow might explain the sequence of events in patient G. N. who had a syncopal attack while performing a standard step-up exercise test<sup>34</sup> during pulmonary function studies. She first experienced excruciating substernal pain which made her stop exercising. About thirty seconds later she became unconscious, apneic, pulseless and finally cyanotic. No heart sounds were audible for the next two and one-half minutes during which time artificial respiration and oxygen were given. The heart sounds, when heard again, were slow and totally irregular. Soon coupled beats were present and, after several minutes, a regular sinus rhythm was established. On another occasion an exercise tolerance test in G. N., with electrocardiographic studies, showed depression of S-T segments in leads I and II and further downward displacement of the S-T segment in



lead V<sub>3</sub>. (Fig. 16.) Exercise was stopped with the onset of chest pain.

In addition to our cases G. N. and M. K., angina was present in 23 per cent of the cases reviewed in the literature. Brenner's case,<sup>13</sup> an eleven year old boy, died during an attack of chest pain. Killingsworth<sup>15</sup> reported the case of a ten year old boy who died following an episode of severe epigastric pain radiating to the arms. In neither these nor the remainder of the thirty-nine cases accepted as instances of primary pulmonary hypertension was coronary sclerosis or pulmonary embolism described at autopsy.

Angina and syncope have been reported in approximately 20 per cent of the cases reviewed. These symptoms may occur together or separately. It is of interest that angina and syncope do not occur in secondary pulmonary hypertension. An increased pulmonary artery pressure and right ventricular hypertrophy are common denominators in both primary and secondary pulmonary hypertension. The apparently unique effect of prisoline\* (2-benzyl-4,5-imidazoline hydrochloride), a sympatholytic and adrenolytic agent,<sup>42</sup> in reducing the elevated pulmonary artery pressure with a concomitant increase of blood flow in primary pulmonary hypertension, suggests that an overactive autonomic nervous system may play a role in this disorder. Whether there is any direct association between this autonomic imbalance and the distinctive symptoms of angina and syncope is a matter for further investigation. Indeed, the mechanism for the production of angina and syncope based on our limited observations is probably more complex than suggested.

In only 8 per cent of the cases reviewed was hemoptysis noted. Evidence of pulmonary infarction was not found at postmortem. Orthopnea, noted in only 10 per cent of the cases, cannot be ascribed to pulmonary congestion in view of the physical, radiologic and postmortem findings. The lungs at necropsy were described grossly as well aerated except for occasional basal congestion. Microscopic studies in these cases revealed normal alveolar membranes and capillaries.

Roentgenograms of the chest should suggest the diagnosis. The characteristic x-ray findings are (1) right ventricular hypertrophy, (2) a bulging pulmonary artery segment, (3) promi-




FIG. 16. Case 1, G. N. Immediately following exercise there is depression of the S-T segments in leads I and II and further downward displacement of the S-T segment in V<sub>3</sub>. The patient complained of substernal pressure at this time.

nent hilar vessels and (4) normal or decreased intrapulmonary vascular markings. There may be overlapping of the spine by the cardiac silhouette in the left anterior oblique view. This finding was present in two of our patients. (Fig. 4c and 10c.) In M. R. the left ventricle was normal in size at necropsy. Pulmonic stenosis<sup>43</sup> and idiopathic dilatation of the pulmonary artery<sup>44</sup> may present similar x-ray features but can be differentiated definitively from primary pulmonary hypertension by right heart catheterization. Patients with pulmonic stenosis rarely have syncopal episodes<sup>26</sup> and may be cyanotic when there is an associated patent foramen ovale.<sup>39</sup> Physical examination in these patients always reveals a loud systolic murmur, occasionally associated with a thrill over the pulmonic area, and the pulmonic second sound is frequently diminished in intensity rather than accentuated. The pulmonic second sound may be accentuated in patients with idiopathic dilatation of the pulmonary artery. However, these patients are generally asymptomatic and the electrocardiogram is usually normal.

The pulmonary hypertension in the three patients herein reported appears to originate proximal to the capillary bed. Arterial blood oxygen unsaturation is associated with pulmonary hypertension secondary to extensive capillary fibrosis. Arterial unsaturation was not

\* Prisoline was generously supplied by Ciba Pharmaceutical Products, Inc., through Dr. Jock L. Graeme.






FIG. 17. Case III, M. K. There is a striking fall in the pulmonary artery pressure after parenteral priscoline, most marked after the second dose. Second dose of priscoline was administered eighteen minutes after the first dose.

observed in our patients. Necropsy in M. R. showed no changes in the pulmonary capillary bed. Furthermore, the ability simultaneously to lower the blood pressure in the lesser circulation (Fig. 17) and increase the blood flow significantly with priscoline is further evidence for a precapillary etiology of the observed hypertension. In the absence of clinical and laboratory evidence of left heart failure it would seem unlikely that the effects of priscoline on the pulmonary circulation could be ascribed to better emptying of the left ventricle. These studies are the subject of a separate report.<sup>45</sup> An attempt to measure capillary pressure was made in M. R. according to the method of Hellems et al.<sup>46</sup> Although our value was in the high normal range, 13 mm. Hg, due to our inability to withdraw blood through the catheter at the site of this pressure measurement, this result cannot be unequivocally accepted.

Intrinsic lung disease is unlikely in view of the pulmonary function and arterial blood oxygen studies as noted in Table III. The decreased ventilatory capacity in G. N. was probably the result of weakness and chest pain caused by the exertion of the test. However, these studies do reveal a significant abnormality in the ratio of oxygen consumption to ventilation during exercise. There was no increase in this ratio as normally expected.<sup>34</sup> This phenomenon is associated with diminished pulmonary blood flow as seen in patients with heart failure<sup>47</sup> and certain types of congenital heart disease.<sup>48-50</sup>

The results of the hemodynamic studies, namely, the high right ventricular end diastolic pressures, the increased arteriovenous oxygen differences and the low cardiac indices, place these patients in the "low output failure" category. Although there is some correlation between

the hemodynamic measurements and the clinical status of the patients, it is of interest that none of them had dependent edema at the time of the studies and one of them, G. N., had no clinical signs of right heart failure.

The presence of normal blood volumes in M. K. and G. N., patients with hemodynamic evidence of right heart failure, is unusual since increased blood volumes ordinarily are associated with right heart failure.<sup>51-53</sup> M. K. had an enlarged, firm liver when the hemodynamic observations were made but there was no dependent edema. G. N. had no edema or enlargement of the liver.

Digitalis did not appear to influence the course of the heart failure in most of the cases reported in the literature that we considered to be instances of primary pulmonary hypertension. The hemodynamic effects of acute and chronic digitalization in M. K. and G. N. confirm this clinical impression. A detailed report of these studies will be made.<sup>54</sup>

The increase in systemic resistance (Fig. 12) in our patients is probably a compensatory mechanism to maintain effective systemic blood pressure in the presence of a decreased cardiac output.<sup>55</sup> However, the magnitude of increase in the pulmonary resistance in these patients is much greater than can be explained by the diminished cardiac output. It is postulated that the pulmonary capillary pressure is not elevated in patients with primary pulmonary hypertension. The necropsy findings of a normal sized left atrium in our case and in those reported support this idea. Therefore, it is believed that the failure to correct for pulmonary capillary pressure in calculating these resistances would not significantly alter the results.

Anoxia, known to increase the pulmonary

resistance in man in acute studies,<sup>56</sup> is not a factor here as demonstrated by blood oxygen studies. Cournand<sup>32</sup> in his latest review of the dynamics of the lesser circulation stated that there was little conclusive evidence for vasomotor control of the pulmonary circulation in normal man. That the autonomic nervous system may be a factor in primary pulmonary hypertension is suggested by the lowering of pulmonary bed resistance in the two patients with this disease who were given priscoline in an acute study.<sup>45</sup> In one patient the fall in resistance was much greater in the pulmonary than in the systemic circulation, whereas tetraethylammonium chloride, an autonomic blocking agent, had a greater effect on the systemic resistance in this patient.<sup>45</sup> (Fig. 18.) Friske *et al.*<sup>57</sup> demonstrated a relatively greater fall in the pulmonary resistance using tetraethylammonium chloride in patients with essential hypertension and normal pulmonary pressures. In reviewing the study of Fowler *et al.*<sup>58</sup> it is noted that tetraethylammonium chloride had a similar effect in some patients with secondary pulmonary hypertension. That other factors may be responsible for increasing the pulmonary artery resistance in man is further suggested by a preliminary observation showing a lack of effect of priscoline on anoxia-induced pulmonary hypertension in man.<sup>59</sup>

Without attempting to draw any conclusions, one is tempted to note, as has previously been pointed out,<sup>18,20</sup> certain similarities between essential systemic hypertension and primary pulmonary hypertension. In both conditions the etiology is unknown. The elevated blood pressure in essential systemic hypertension is considered to indicate a state of increased tonus of the pre-capillary arterioles while in primary pulmonary hypertension there would seem to be a state of increased tonus of the small pulmonary arteries. Usually there is ventricular hypertrophy of the particular circuit involved. The respective vascular beds, pulmonary or peripheral, show similar histologic patterns ranging from normal vessels to obliterative arteriolar sclerosis. In either disease hypertension may exist without significant arteriolar changes.<sup>22,61,62</sup> Pharmacodynamic studies<sup>45,63</sup> in both diseases suggest that the autonomic nervous system may be a factor in increasing the peripheral vascular resistance.

Of a group of autopsied cases reviewed by Greene *et al.* and cited as idiopathic dilatation

of the pulmonary artery,<sup>44</sup> but in which heart catheterization was not done, four showed right ventricular hypertrophy without pulmonary vascular sclerosis. Two of the latter died in congestive heart failure. It may be that these cases as well as those reported by De Navasquez *et*




FIG. 18. Case III, M. K. The predominant effect of TEAC was on the peripheral resistance whereas priscoline exerted a maximal effect on the pulmonary resistance.

*al.*<sup>17</sup> were instances of primary pulmonary hypertension.

Thrombosis of the smallest vessels of the pulmonary circuit has been described in the entity under discussion and was found in the autopsied cases in this series. Rich<sup>60</sup> described similar lesions in 90 per cent of cases of tetralogy of Fallot that he studied. He believed that the predisposing factors for the development of thrombosis were polycythemia and sluggish pulmonary blood flow. The data which we have presented show that patients with primary pulmonary hypertension have a decreased pulmonary blood flow without polycythemia. There may be other predisposing factors in the pulmonary vascular bed in this disease to account for this unusual finding not described in other conditions with diminished pulmonary blood flow. In view of the preponderance of the sclerotic changes over the number of thrombotic lesions in the autopsied case reported it seems unlikely that sclerosis was the result of thrombosis.

Since it is generally accepted that the hallmark of prolonged hypertension of the greater circulation, essential or secondary, is found in the arteriolar bed, many of the observed changes in the pulmonary vasculature in primary pulmonary hypertension are similarly interpreted.

Undoubtedly, these sclerotic vessels add a significant increment to an already excessive vascular resistance further burdening the right heart but the deduction that these changes initiate the hypertension seems unwarranted. The findings of a normal pulmonary arteriolar bed in several cases of isolated right ventricular hypertrophy,<sup>6,12,14,16,17,20,22</sup> assumed by us to be instances of primary pulmonary hypertension, one proven to have marked pulmonary hypertension by right heart catheterization<sup>22</sup> and the effect of certain drugs in lowering the pulmonary resistance, suggest that the sclerotic changes, when seen, are of a secondary nature. Conceivably, involvement of the pulmonary vasculature by pulmonary emboli, parasitic infestations, lymphangitic carcinomatosis or pulmonary arteritis, if sufficiently severe and diffuse, might produce an identical clinical picture.

The material herewith presented indicates that primary pulmonary hypertension is a formidable disease with a grave prognosis. Heart catheterization not only enables the clinical investigator to make this diagnosis during life but also permits him to evaluate quantitatively the effects of various therapeutic measures and to obtain some additional information about the dynamics of the pulmonary circulation. The clinician is often faced with the problem of treating the effects of a disease while the etiology is still obscure. A logical approach to therapy in primary pulmonary hypertension would be to relieve the markedly increased resistance in the pulmonary circulation. Priscoline administered parenterally in acute studies has proven effective in this regard.<sup>46</sup> In addition, clinical improvement as manifested by increased exercise tolerance has also been noted after parenteral priscoline. However, studies thus far would indicate that the beneficial effects in primary pulmonary hypertension of parenterally administered priscoline are of short duration, i.e., twenty to thirty minutes; the response to chronic oral priscoline is inconclusive. A similar approach might be indicated in certain cases of pulmonary embolism with occlusion of but a relatively small segment of the pulmonary arterial tree. In view of the known reserve of the pulmonary circulation the pathologic findings in such cases seem inadequate to explain the fatal outcome. It is possible that reflex sympathetic overactivity might add a significant increment to the pulmonary resistance, sufficient to cause acute cor pulmonale and death. Prisco-

line or similarly acting drugs would then seem indicated in this situation. However, the conclusions drawn by various investigators in evaluating the role of vasomotor activity in induced pulmonary embolization in animals are contradictory.<sup>64-67</sup> Studies of a similar nature are in progress in this laboratory.

Since primary pulmonary hypertension presents certain unique features pointing toward isolated sympathetic overactivity, surgical thoracic sympathectomy must be considered. This, of course, is a subject for further investigation.

#### SUMMARY

The concept of primary pulmonary hypertension is defined and the clinical and hemodynamic features in three patients with this disease are presented. Such cases present a clinical syndrome that has been previously described under various names, e.g., "primary pulmonary vascular sclerosis," "right ventricular hypertrophy of obscure origin" and "idiopathic pulmonary hypertension."

The salient clinical features in this syndrome are exertional weakness and dyspnea in patients who subsequently develop right heart failure without antecedent cardiac or pulmonary disease. Effort syncope and angina are most significant when present. Sudden death is not infrequent. The pertinent physical findings are a normal systemic blood pressure, clear lungs, an accentuated pulmonic second sound and the variability or absence of heart murmurs. Venous distention and hepatomegaly occur unassociated with peripheral edema or ascites except late in the course of the disease. Cyanosis, when encountered, is a terminal feature unless the disease is complicated by a patent foramen ovale. This syndrome has been seen in both sexes at all ages but is most common between the ages of twenty and forty years.

The electrocardiographic tracings are consistent with right ventricular hypertrophy. The characteristic x-ray findings are (1) right ventricular enlargement, (2) a bulging pulmonary artery segment, (3) prominent hilar vessels and (4) normal or diminished intrapulmonary vascular markings.

Marked right ventricular hypertrophy is a constant pathologic finding in primary pulmonary hypertension but the nature, severity and distribution of the pulmonary vascular changes proximal to the capillary bed show considerable



variation. In some cases no significant pulmonary vascular changes are noted.

Pulmonary function studies were performed in two of the patients and ruled out intrinsic lung disease. Hemodynamic studies, utilizing the technic of right heart catheterization, were performed in the three patients presented in this series. All had greatly elevated pulmonary artery pressures, elevated right ventricular end diastolic pressures, diminished cardiac outputs, increased arteriovenous oxygen differences and normal arterial blood oxygen saturations. The blood volumes and hematocrits were normal in two of the patients and slightly elevated in the third. One patient had no clinical evidence of right heart failure and none had edema.

An attempt was made to correlate the hemodynamics at rest and during exercise with some of the clinical findings. A seven- to ninefold increase in pulmonary resistance was calculated at rest. Physiologic studies and postmortem findings were presented to support the view that the locus of the increased resistance is in the small pulmonary arteries.

Isolated overactivity of the sympathetic nervous system was suggested by the pronounced effect of prisolone, an adrenolytic and sympatholytic agent, in lowering the pulmonary artery blood pressure. On the basis of the material presented the problem of sympathectomy in primary pulmonary hypertension was raised.

**Acknowledgments:** The authors wish to express their appreciation to Drs. Dickinson W. Richards, Jr., André Cournand and David Seegal for reviewing this paper and for their valuable suggestions concerning the material presented. Technical assistance by Miss Leila Israel and Mrs. Dorothy Bowen is gratefully acknowledged.

## REFERENCES

1. Cournand, A. and Ranges, H. A. Catheterization of the right auricle in man. *Proc. Soc. Exper. Biol. & Med.*, 46: 462, 1941.
2. Mönckeberg, J. G. Ueber die genuine Arteriosklerose der Lungenarterie. *Deutsche med. Wchnschr.*, 33: 1243, 1907.
3. Rössle, R. Ueber Hypertrophie und Organkorrelation. *München. med. Wchnschr.*, 55: 377, 1908.
4. Sanders, E. Primary pulmonary arteriosclerosis with hypertrophy of the right ventricle. *Arch. Int. Med.*, 3: 257, 1909.
5. Schütte, H. Rechtsseitige Herzhypertrophie hervorgerufen durch eine entzündliche Veränderung der kleinen Lungenarterien. *Centralbl. f. allg. Path. u. path. Anat.*, 25: 483, 1914.
6. Ljungdahl, M. Untersuchungen über die Arteriosklerose des kleinen Kreislaufs. Weisbaden, 1915. J. F. Bergmann.
7. Hart, C. Ueber die isolierte Sklerose der Pulmonalarterie. *Berl. klin. Wchnschr.*, 53: 304, 1916.
8. Krutsch, G. Ueber rechtsseitige Herzhypertrophie durch Einengung des Gesamtquerschnittes der kleineren und kleinsten Lungenarterien. *Frankfurt. Ztschr. f. Path.*, 23: 163, 1920.
9. Eppinger, H. and Wagner, R. Zur Pathologie der Lunge. *Wien. Arch. f. inn. Med.*, 1: 83, 1920.
10. Tschistowitsch, T. Thrombo-endarterite pulmonaire. *Compt. rend. Soc. de biol.*, 89: 627, 1923.
11. MacCallum, W. G. Obliterative pulmonary arteriosclerosis. *Bull. Johns Hopkins Hosp.*, 49: 37, 1931.
12. Ulrich, H. L. The clinical diagnosis of pulmonary arteriosclerosis. *Ann. Int. Med.*, 6: 632, 1932.
13. Brenner, O. Pathology of the vessels of the pulmonary circulation. *Arch. Int. Med.*, 56: 211, 457, 724, 976, 1189, 1935.
14. Kaump, D. H. and Dry, T. J. Pulmonary arteriolar sclerosis. *Arch. Int. Med.*, 61: 1, 1938.
15. Killingsworth, W. P. and Gibson, S. Primary proliferative pulmonary arteriolar sclerosis. *Am. J. Dis. Child.*, 57: 1099, 1939.
16. East, T. Pulmonary hypertension. *Brit. Heart J.*, 2: 189, 1940.
17. De Navasquez, S., Forbes, J. R. and Holling, H. E. Right ventricular hypertrophy of unknown origin: so-called pulmonary hypertension. *Brit. Heart J.*, 2: 177, 1940.
18. Brill, I. C. and Krygier, J. J. Primary pulmonary vascular sclerosis. *Arch. Int. Med.*, 68: 560, 1941.
19. Gilmour, J. R. and Evans, W. Primary pulmonary hypertension. *J. Path. & Bact.*, 58: 687, 1946.
20. Raaschou, F. and Samuelsson, S. Isolated hypertrophy of the right ventricle of the heart of unknown cause. *Acta med. Scandinav.*, Suppl., 130: 206, 102, 1946.
21. Cross, K. R. and Kobayshi, C. K. Primary pulmonary vascular sclerosis. *Am. J. Clin. Path.*, 17: 155, 1947.
22. Whittenberg, S. J. Primary pulmonary hypertension. Reported in the Clinical Society Conference of the Beth David Hospital in New York, April 10, 1950; (to be published).
23. Jernell, O. Pulmonary sclerosis. *Acta med. Scandinav.*, 138: 430, 1950.
24. Romberg, E. Sklerose der Lungenarterie. *Deutsches Arch. klin. Med.*, 48: 197, 1891.
25. Chapman, Don W., Earle, D. M., Gugle, L. J., Huggins, R. A. and ZimdaHL, W. Intravenous catheterization of the heart in suspected congenital heart disease. *Arch. Int. Med.*, 84: 644, 1949.
26. Wood, P. Congenital heart disease. A review of its clinical aspects in the light of experience gained by means of modern techniques. *Brit. M. J.*, 2: 639, 693, 1950.
27. Johnson, J. B., Ferrer, I. M., West, J. R. and Cournand, A. The relation between electrocardiographic evidence of right ventricular hypertrophy and pulmonary arterial pressure in patients with chronic pulmonary disease. *Circulation*, 1: 536, 1950.
28. Nellen, M. Idiopathic pulmonary hypertension. *South Africa M. J.*, 21: 682, 1947.

29. VAN SLYKE, D. D. and NEILL, J. M. The determination of gases in blood and other solutions by vacuum extraction and manometric measurement. *J. Biol. Chem.*, 61: 524, 1924.
30. NOBLE, R. P. and GREGERSEN, M. I. Blood volume in clinical shock. i. Mixing time and disappearance rate of T-1824 in normal subjects and in patients in shock, determination of plasma volume in man from ten minute sample. *J. Clin. Investigation*, 25: 158, 1946.
31. Cournand, A., RILEY, R. L., BREED, E. S., BALDWIN, E. DE F. and RICHARDS, D. W., JR. Measurement of cardiac output in man using the technique of catheterization of right auricle or ventricle. *J. Clin. Investigation*, 24: 106, 1945.
32. Cournand, A. Some aspects of the pulmonary circulation in normal man and in chronic cardiopulmonary diseases. *Circulation*, 2: 641, 1950.
33. SCHOLANDER, P. F. Analyzer for accurate estimation of respiratory gases in one-half cubic centimeter samples. *J. Biol. Chem.*, 167: 235, 1947.
34. BALDWIN, E. DE F., Cournand, A. and RICHARDS, D. W., JR. Pulmonary insufficiency. i. Physiological classification, clinical methods of analysis, standard values in normal subjects. *Medicine*, 27: 243, 1948.
35. HARVEY, R. M., FERRER, M. I., CATHCART, R. T., RICHARDS, D. W., JR. and Cournand, A. Some effects of digoxin upon the heart and circulation in man. Digoxin in left ventricular failure. *Am. J. Med.*, 7: 439, 1949.
36. ARILLAGA, F. C. Sclérose de l'artère pulmonaire secondaire à certains états pulmonaires chroniques (cardiaques noirs). *Arch. d. mal. du cœur*, 6: 518, 1913.
37. CASTEX, M. R., CAPDEHOURAT, E. L. and MAZZEI, E. S. El factor bronquial en los cardiacos negros de Ayerza. *Brasil méd-cir.*, 10: 139, 1948.
38. LEGRAND, R. Black cardiac disease. *Echo méd. du nord*, 3: 284, 1935.
39. ENGLE, M. E. and TAUSSIG, H. B. Valvular pulmonic stenosis with intact ventricular septum and patent foramen ovale. Report of illustrative cases and analysis of clinical syndrome. *Circulation*, 2: 481, 1950.
40. DRESSLER, W. Effort syncope as an early manifestation of primary pulmonary hypertension; (to be published).
41. SCHWARTZ, S. P., LEINER, G. C. and MICHOTOM, R. J. Transient ventricular fibrillation. ii. The effects of gradually induced oxygen deficiency on patients with transient ventricular fibrillation and on patients with periodic standstill of the ventricle. *Am. Heart J.*, 37: 918, 1949.
42. CHES, D. and YONKMAN, F. F. Adrenolytic and sympatholytic actions of priscor (benzyl-imidazole). *Proc. Soc. Exper. Biol. & Med.*, 61: 127, 1946.
43. DOW, J. W., LEVINE, H. D., ELKIN, M., HAYNES, F. W., HELLEMS, H. K., WHITTENBERGER, J. W., FERRIS, B. C., GOODALE, W. T., HARVEY, W. P., EPPINGER, E. C. and DEXTER, L. Studies of congenital heart disease. iv. Uncomplicated pulmonic stenosis. *Circulation*, 1: 267, 1950.
44. GREENE, D. G., BALDWIN, E. DE F., BALDWIN, J. S., HIMMELSTEIN, A., ROH, C. E. and Cournand, A. Pure congenital pulmonary stenosis and idiopathic congenital dilatation of the pulmonary artery. *Am. J. Med.*, 6: 24, 1949.
45. DRESDALE, D. T., SCHULTZ, M., FALSONE, J. J. and NOBLE, W. J. Primary pulmonary hypertension. ii. The effect of priscoline and tetraethylammonium chloride on the pulmonic and systemic resistances; (to be published).
46. HELLEMS, H. K., HAYNES, F. W. and DEXTER, L. Pulmonary "capillary" pressure in man. *J. Applied Physiol.*, 2: 24, 1949.
47. HICKAM, J. B. and CARGILL, W. H. Effect of exercise on cardiac output and pulmonary arterial pressure in normal persons and in patients with cardiovascular disease and pulmonary emphysema. *J. Clin. Investigation*, 27: 10, 1948.
48. BING, R. J., VANDAM, L. D. and GRAY, F. D., JR. Physiological studies in congenital heart disease. ii. Results of preoperative studies in patients with tetralogy of Fallot. *Bull. Johns Hopkins Hosp.*, 80: 121, 1947.
49. SCHULTZ, M., DRESDALE, D. T. and MICHOTOM, R. J. Hemodynamic studies in congenital heart disease before and after operation. i. Pulmonary valvulotomy in pulmonic stenosis with atrial septal defect. ii. Ligation in patent ductus arteriosus; (to be published).
50. VANDAM, L. D., BING, R. J. and GRAY, F. D., JR. Physiological studies in congenital heart disease. iv. Measurement of the circulation in five selected cases. *Bull. Johns Hopkins Hosp.*, 81: 192, 1947.
51. GIBSON, J. G. and EVANS, W. A., JR. Clinical studies of the blood volume. iii. Changes in blood volume, venous pressure and blood velocity rate in chronic congestive heart failure. *J. Clin. Investigation*, 16: 851, 1937.
52. MENEELY, G. R. and KALTREIDER, N. L. A study of the volume of the blood in congestive heart failure. Relation to other measurements in 15 patients. *J. Clin. Investigation*, 22: 521, 1943.
53. ALTSCHULE, M. D. Physiology in Diseases of the Heart and Lungs, p. 107. Cambridge, 1949. Harvard University Press.
54. DRESDALE, D. T., SCHULTZ, M. and MICHOTOM, R. J. Hemodynamic effects of lanatoside 'C' in i. Normal patients. ii. Patients with primary pulmonary hypertension; (to be published).
55. ALTSCHULE, M. D. P. 33.<sup>53</sup>
56. MOTLEY, H. L., Cournand, A., WERKO, L., HIMMELSTEIN, A. and DRESDALE, D. T. The influence of short periods of induced anoxia upon pulmonary artery pressure in man. *Am. J. Physiol.*, 150: 315, 1947.
57. FRISKE, R. A., HAMMERSTROM, S., LAGERLOF, H., WERKO, L., BJORKHEIM, G., HOLMGREN, A. and LARSSON, Y. The effect of tetraethylammonium chloride in arterial hypertension. *Am. J. Med.*, 5: 807, 1948.
58. FOWLER, N. O., WESTCOTT, R. N., HAUENSTEIN, V. D., SCOTT, R. C. and McGUIRE, J. Observations on autonomic participation in pulmonary arteriolar resistance in man. *J. Clin. Investigation*, 29: 1387, 1950.
59. DRESDALE, D. T. and SCHULTZ, M. The effects of

- prisco-line and tetraethylammonium chloride on the pulmonary circulation in patients with anoxia induced pulmonary hypertension and in patients with chronic pulmonary disease; (to be published).
60. RICH, A. R. A hitherto unrecognized tendency to the development of widespread pulmonary vascular obstruction in patients with congenital pulmonary stenosis (tetralogy of Fallot). *Bull. Johns Hopkins Hosp.*, 82: 389, 1948.
61. FISHBERG, A. M. Hypertension and Nephritis, 3rd ed., p. 489. Philadelphia, 1934. Lea & Febiger.
62. CASTLEMAN, B. and SMITHWICK, R. H. The relation of vascular disease to the hypertensive state. *J. A. M. A.*, 121: 1256, 1943.
63. LYONS, R. H., MOE, G. K., NELIGH, R. B., HOOBLER, S. W., CAMPBELL, K. N., BERRY, R. L. and RENNICK, B. R. The effects of blockade of the autonomic ganglia in man with tetraethylammonium. *Am. J. M. Sc.*, 213: 315, 1947.
64. HAYNES, F. W., KINNY, T. D., HELLEMS, H. K. and DEXTER, L. Circulatory changes in experimental pulmonary embolism. *Federation Proc.*, 6: 125, 1947.
65. HARA, M. and SMITH, J. R. Experimental observations on embolism of pulmonary lobar arteries. *J. Thoracic Surg.*, 18: 536, 1949.
66. DOTTER, C. T. and LUKAS, D. S. Acute cor pulmonale. An experimental study utilizing a special cardiac catheter. *Am. J. Physiol.*, 164: 254, 1951.
67. DALEY, R., WADE, J. D., MARAIST, F. and BING, R. J. Pulmonary hypertension in dogs induced by injection of lycopodium spores into the pulmonary artery, with special reference to the absence of vasomotor reflexes. *Am. J. Physiol.*, 164: 380, 1951.



# Pulmonary Function Studies in Polycythemia Vera\*

## Results in Five Probable Cases

WALTER NEWMAN, M.D., JAMES A. FELTMAN, M.D. and BLANCHE DEVLIN  
New York, New York

THERE have been numerous studies of pulmonary function in polycythemia vera. The results have been variable and inconsistent and many of the older technics

which were used are open to criticism at the present time. Therefore, the results of previous investigators measuring cardiac output and arteriovenous oxygen differences will not be listed. Of particular interest are reported results of vital capacity, residual air, total capacity, ratio of residual air to total capacity, and arterial oxygen saturation at rest and exercise. (Table 1.)

Examination of this table would indicate that in general the vital capacity is reduced, residual air increased, total capacity normal or decreased and ratio of residual air to total capacity is increased. The arterial oxygen saturation is normal or slightly reduced at rest and decreased with exercise. In Brooks' case<sup>2</sup> the lung volumes after treatment improved. Other investigators found an increase in blood volume<sup>10</sup> and viscosity<sup>1</sup> in polycythemia.

We have made detailed studies of pulmonary function in five cases of polycythemia. All of these were studied before and after treatment with phlebotomy.

### METHODS

The measurement of residual air volume was made using an open circuit method in which the nitrogen of the lungs is washed out by continuous inhalation of pure oxygen and collected over a period of seven minutes. An index of intrapulmonary mixing was obtained by sampling of the alveolar air at the end of this period.<sup>11</sup>

The measurements of lung volumes and maximum breathing capacity were obtained using the spirographic technic.<sup>11</sup>

Arterial blood samples were obtained from the brachial artery using an indwelling Cournand-

TABLE I  
RESUME OF RESULTS REPORTED IN THE LITERATURE

	Vital Capacity
Isaacs <sup>1</sup>	Decreased, 1 case; average vital capacity 2115; predicted value 4700; average % predicted value 45 %
Brooks <sup>2</sup>	Decreased, 1* case; vital capacity 1.86 L. before treatment; after treatment 2.36 L. (phenylhydrazine)
Altschule <sup>3</sup>	3 cases, normal in 2; slightly decreased in 1; no change in 2 cases treated with phlebotomy
Stewart <sup>4</sup>	6 cases, reduced in 3; one case treated; vital capacity 3.3 L. before treatment to 3.9 L. after treatment
	Residual Air
Harrop <sup>5</sup>	Increased, 7 cases
Brooks <sup>2</sup>	Increased, 1* case; residual air 1.19 L. before treatment; after treatment 0.99 L.
	Total Capacity
Harrop <sup>5</sup>	Normal, 7 cases
Brooks <sup>2</sup>	Decreased, 1* case; 3.07 L. before treatment; 3.35 L. after treatment
	Ratio of Residual Air to Total Capacity
Harrop <sup>5</sup>	Increased, 7 cases
Brooks <sup>2</sup>	Increased, 1* case; before treatment 38.7 %; after treatment 29.5 %
	Arterial O <sub>2</sub> Saturation at Rest
Harrop <sup>5</sup>	Decreased
Wasserman <sup>6</sup>	17 cases with arterial O <sub>2</sub> saturation determinations; 8 below range 94% to 98% and one case showed 88.2% saturation
von Bergmann, <sup>7</sup> Brooks, <sup>2</sup> Hitzinger, <sup>8</sup> and Rover	Normal
	Arterial O <sub>2</sub> Saturation After Exercise
Harrop <sup>5</sup>	3 cases: 91%→89%; 93%→87%; 95%→90%

\*Second case reported was secondary polycythemia.

\* From the Medical Service of the Veterans Administration Hospital, Bronx, N. Y. Reviewed in the Veterans Administration and published with the approval of the Chief Medical Director. The statements and conclusions published by the authors are the result of their own study and do not necessarily reflect the opinion or policy of the Veterans Administration.

type needle. The blood was collected by the technic described by Riley,<sup>12</sup> using heparin to prevent clotting and a small globule of mercury in the syringe to facilitate mixing. The oxygen content and capacity and the carbon dioxide content were determined on duplicate samples

of Riley.<sup>12</sup> Duplicate measurements were made using two Roughton-Scholander syringes and all samples checked within 2 mm. of mercury. Expired air samples were collected in duplicate simultaneously with the blood specimens. Oxygen intake and carbon dioxide output were

TABLE II A  
PULMONARY FUNCTION STUDIES BEFORE AND AFTER PHLEBOTOMIES IN FIVE CASES OF POLYCYTHEMIA VERA

		Case I Before After (2.85 L. in 21 days)		Case II Before After (2.5 L. in 21 days)		Case III Before After (3.7 L. in 30 days)		Case IV Before After (5.4 L. in 32 days)		Case V Before After (4.8 L. in 60 days)	
Red blood count, million per cu. mm.		7.4	4.8	6.8	4.8	7.7	4.7	7.5	4.96	7.25	5.2
Hemoglobin, gm. per 100 cc.		22	14.6	20.1	14.5	23.6	15	23	15.7	21.8	16.3
Hematocrit, %		67	46	54	47	73	46	67	43	77	48
Blood volume cc.; p <sub>32</sub>	Actual	10,000	7500	6300	5750	7930	5360	-----	-----	8450	7900
	Predicted	3960	3960	4800	4800	3890	3890	-----	-----	5020	5020
Vital capacity in cc.	Actual	4130	4430	4540	4525	3838	3780	2070	3930	2950	4500
	Predicted	4130	4130	4560	4560	4180	4180	4070	4070	4940	4940
Residual air cc.	Actual	1430	1582	1286	1005	1144	863	1626	1700	1378	1307
	Predicted	1320	1320	1150	1150	1030	1030	1320	1320	1220	1220
Total capacity in cc.	Actual	6040	6292	6071	5627	4807	4530	3937	5970	4435	5926
	Predicted	5450	5450	5810	5810	5210	5210	5390	5390	6180	6180
Residual air Total capacity X 100	Actual	23.7	25.2	21.2	17.9	23.8	19.1	41.4	28.5	31.1	22.1
	Predicted	24.4	24.4	19.8	19.8	19.8	19.8	24.4	24.4	19.8	19.8
Alveolar N <sub>2</sub> , (normal 2.5 %)		1.6	2.02	2.18	1.7	1.54	1.26	2.5	2.02	1.88	1.53
Maximum breathing capacity (% of normal)		93.5	102.0	70.3	103	133	173	48.8	95.3	33.3	60.5
Maximum breathing capacity L./min.	Actual	92.0	124.5	111	163	157	264	62	121	55	99
	Predicted	98.4	98.4	158	158	118	118	127	127	164	164
Arterial O <sub>2</sub> saturation (normal 94-98 %)		94.4	95.0	95.5	97.0	95.4	95.8	69	95.4	65	80.3
Arterial pO <sub>2</sub> mm. Hg (normal 95 mm. Hg)		87	92	89	101	104	92	34	92	38	51
Arterial pCO <sub>2</sub> mm. Hg (normal 40 mm. Hg)		43	43	47	40	45	47	78	43	73	66
Arterial CO <sub>2</sub> content (normal 44-53 vol. %)		41.3	50.3	44.7	47.7	43.4	52.4	63.1	51.8	59.3	60.4
pH (calculated) (normal 7.4)		7.35	7.39	7.33	7.42	7.38	7.38	7.3	7.42	7.28	7.34
Ventilation L./min./M <sup>2</sup> At rest	Normal 3.20 ± .65	4.68	4.39	3.58	3.61	3.57	3.64	3.01	3.64	3.31	3.20
	After exercise Normal 8.70 ± .81	10.8	13.6	12.5	9.56	4.33	9.05	-----	9.35	9.48	9.56
Oxygen consumption in cc./min./M <sup>2</sup> At rest	Normal 129 ± 13	152	154	180	163	150	137	144	140	166	124
	After exercise Normal 480 ± 74	522	602	775	634	318	486	-----	443	454	458
Oxygen removal in cc./L. of ventilation At rest	Normal 46.8 ± 7.1	38.9	42.3	61.0	54.5	51.7	46.2	58.2	46.5	61.8	52.0
	After exercise Normal 54.8 ± 6.2	57.6	53.1	74.7	79.9	90.4	65.7	-----	56.8	58.4	58.1
Arterial O <sub>2</sub> saturation after exercise (normal 94-98 %)		94.4	94.0	98.0	96.0	75.0	95.0	-----	93.7	60.6	78.6

using the Van Slyke-Neill apparatus.<sup>13</sup> All samples checked within 0.2 volumes per cent.

The arterial blood tensions of oxygen and of carbon dioxide were determined directly in millimeters of mercury, using the direct method

calculated from the percentage of oxygen and carbon dioxide found in these expired air samples, as determined in a Scholander gas analyzer.<sup>14</sup> The duplicate measurements checked within 0.04 per cent.

Arterial blood samples and expired air were collected under basal conditions with the patients breathing room air, high and low oxygen concentrations each for twenty-minute periods when indicated. Finally oxygen saturation was

due primarily to a decrease in the blood viscosity and to some extent a decrease in total blood volume. This increase in maximum breathing capacity was the only change noted in Cases I and II.

TABLE II B

PULMONARY FUNCTION STUDIES BEFORE AND AFTER PHEBOTOMIES IN FIVE CASES OF POLYCYTHEMIA VERA

		Case I		Case II		Case III		Case IV		Case V	
		Before	After	Before	After	Before	After	Before	After	Before	After
		(2.85 L. In 21 days)		(2.5 L. In 21 days)		(3.7 L. In 30 days)		(5.4 L. In 32 days)		(4.8 L. In 60 days)	
Room air breathing	Alveolar pO <sub>2</sub> mm. Hg	100	101	88	100	92	91	48	92	58	69
	Arterial pO <sub>2</sub> mm. Hg	87	92	89	101	104	92	34	92	38	54
	Arterial pCO <sub>2</sub> mm. Hg	43	43	47	40	45	47	78	43	73	58
	Alveolar-arterial gradient	13	9	-1	-1	-12	-1	14	0	20	15
	Dead space ml. % of tidal air	35	31	24	18	22	31	42	31	27	31
High oxygen breathing	Alveolar pO <sub>2</sub>							257*		97†	288*
	Arterial pO <sub>2</sub>							95		54	91
	Arterial pCO <sub>2</sub>							84		69	64
	Alveolar-arterial gradient							162		43	197
	Dead space							48		27	31
Low oxygen breathing (12.4%)	Alveolar pO <sub>2</sub>	47	42	41	47	37	43				
	Arterial pO <sub>2</sub>	44	38	40	52	37	35				
	Arterial pCO <sub>2</sub>	38	41	41	36	46	42				
	Alveolar-arterial gradient	3	4	1	-5	0	8				
	Dead space	28	31	22	19	32	31				

\*51.7 %

†27.35 %

determined after one minute of standard exercise and the expired gas during the exercise was collected in a Douglas bag and analyzed. During the five-minute recovery period the expired air was collected and analyzed. The expired air, except during exercise, was collected in the Tissot apparatus and the spirogram attached permitted calculation of the respiratory rate and tidal air under the various conditions. The alveolar-arterial gradients were calculated from the data obtained according to the method of Riley and Cournand.<sup>15</sup>

pH determinations were calculated from line charts of the Henderson-Hasselbalch equation.<sup>16</sup>

The blood volumes were determined by the use of P<sup>32</sup> labeled red cells.

## COMMENTS

Five cases of polycythemia vera were subjected to detailed pulmonary function studies before and after phlebotomies. The results of these studies are listed in Table II.

A basic change in all five cases was an increase in the maximum breathing capacity after phlebotomy. This is believed to be due to an increase in the elasticity and a decrease in the viscous resistance of the lung tissue to changes in shape, and when present, to a decrease in the anoxemia. These former changes are probably

Case III, in addition to the maximum breathing capacity change, demonstrated an unusual aberration of pulmonary function. During exercise there was a failure to increase ventilation significantly, the actual value being approximately half the predicted value. This in turn was reflected by an inadequate oxygen consumption during exercise and consequently a decrease in oxygen saturation with exercise. This phenomenon is especially interesting in view of the fact that the patient had the ability to hyperventilate, considering that the maximum breathing capacity was 133 per cent of the predicted normal. He also had the stimuli to hyperventilate with exercise, viz., low pH, high arterial pCO<sub>2</sub> and anoxia. However, there was marked hypoventilation. In view of these facts we feel justified in concluding that this patient's respiratory center was insensitive to these stimuli. Since this patient did manual labor, it is reasonable to assume that he was anoxic during most of his working day. This was substantiated by the fatigue, of which the patient was aware. Following phlebotomies this patient's ventilation and oxygen consumption during exercise and his oxygen saturation after exercise returned to normal. However, his effective alveolar ventilation at rest, as reflected by the arterial pCO<sub>2</sub>, remained elevated. These



changes with phlebotomy would suggest that the respiratory center damage was at least partially reversible.

Case iv demonstrated a marked decrease in vital capacity, total capacity and maximum breathing capacity. The mechanisms for these changes are believed to be the same as postulated before, namely, increased blood viscosity and volume causing a decrease in elasticity, and an increase in the viscous resistance of the lung tissue. Before phlebotomy this patient had a definite distribution gradient indicating perfusion of poorly ventilated areas.<sup>15,18</sup> One explanation for this phenomenon is that in the normal individual at rest all the alveoli are not being well ventilated or perfused. In this patient with polycythemia and increased blood volume there was an opening up of the capillary beds of some of these poorly ventilated alveoli. This would result in a poor correlation of perfusion and ventilation and thus give rise to a high distribution gradient. A similar mechanism has been postulated by Riley in patients with mitral stenosis and pulmonary hypertension.<sup>19</sup> This patient was strikingly unsaturated at rest, with a very poor effective alveolar ventilation in view of the arterial  $p\text{CO}_2$  of 78 mm. of mercury. Thus there appear to be two factors contributing to the arterial oxygen unsaturation at rest, namely, inadequate ventilation and venous admixture. In addition, we believe that a third factor was playing a part, namely, respiratory center damage. This is substantiated by the fact that this patient had the ability to ventilate 62 L. per minute, at least for short periods of time. He certainly had stimuli for hyperventilation at rest with an arterial pH of 7.3, arterial  $p\text{CO}_2$  of 78 mm. of mercury and arterial  $p\text{O}_2$  of 34 mm. of mercury. Despite these strong stimuli this patient's ventilation at rest, while breathing room air, was only 7.83 L. per mm., 13 per cent of his maximum breathing capacity.

One of the remarkable features of this case was the complete return to essentially normal values for ventilation and gas exchange following adequate phlebotomy. This again would indicate the reversibility of the lung volume changes, the distribution gradient and the respiratory center damage.

We do not believe that these changes were due to any significant underlying chronic pulmonary disease such as emphysema since the alveolar nitrogen after breathing pure oxygen was normal and the spirogram showed no

evidence of obstruction. Furthermore, except for a slight increase of the residual air, the lung volume and arterial gas studies all became normal after phlebotomy. This would seem to preclude the presence of any significant pulmonary parenchymal disease. It is of interest to note that the alveolar nitrogen was within normal limits before and after phlebotomy even though tension studies indicated a high venous admixture component. This is in striking contrast to the findings in pulmonary emphysema in which a high alveolar-arterial gradient exists with high oxygen and a high alveolar nitrogen after breathing pure oxygen for seven minutes. Evidently, when there are permanent intrinsic tissue changes in the lung, nitrogen is not washed out as easily as in this patient with polycythemia.

Case v demonstrated essentially the same pulmonary function abnormalities as Case iv except that in addition there was a high diffusion gradient. This possibly was due to the fact that the diffusing surface, i.e., the capillary bed, was reduced by thrombosis. Following phlebotomies this patient's lung volumes returned to normal. This is consistent with the hypothesis that these changes were due to decreased elasticity and increased viscous resistance of the lung as a result of the large volume of viscous blood these patients have. Failure of the maximum breathing capacity to return completely to normal is believed to be due to persistence of the anoxia. This would interfere with the muscular effort necessary for performing the maximum breathing capacity.

The persistence of anoxia was largely due to the continued poor correlation between ventilation and perfusion. This gas exchange abnormality was then primarily due to poor effective alveolar ventilation (arterial  $p\text{CO}_2$  66 mm. of mercury) and less to opening up of the capillary beds of alveoli which normally are poorly ventilated. The poor effective alveolar ventilation in turn is thought to reflect respiratory center damage. At this stage the patient had a maximum breathing capacity of 99 L. per minute and was ventilating only 7.55 L. per minute despite the presence of strong stimuli. Thus he was utilizing only 7.6 per cent of his maximum breathing capacity at a time when his arterial  $p\text{O}_2$  was 51 mm. of mercury and  $p\text{CO}_2$  66 mm. of mercury. In order to eliminate the possibility of anoxic depression of the respiratory center the patient was placed in a Drinker respirator for forty-eight hours. With

this procedure after twenty-four hours the arterial  $p\text{CO}_2$  dropped to 53 mm. of mercury,  $\text{CO}_2$  content went from 62.1 volumes per cent to 51.6 volumes per cent, and the oxygen saturation rose to 93 per cent. However, one hour after removal from the respirator the arterial  $p\text{CO}_2$

This anoxia would tend to augment the decrease in maximum breathing capacity. In addition, the increased blood viscosity and volume would tend to open the capillary bed of alveoli which are normally poorly ventilated and thus increase the distribution gradient and resultant anoxia.




FIG. 1. Sequence of events which may occur in some cases of polycythemia vera.

rose to 61 mm. of mercury and the oxygen saturation dropped to 82 per cent. This would seem to indicate respiratory center damage due to causes other than anoxia; however, a more prolonged period in the respirator might be necessary before this could be definitely ruled out.

We, therefore, believe that this patient demonstrates not only the possible irreversibility of respiratory center damage but also possible progression of this disease along lines not previously indicated.

#### OBSERVATIONS

We have presented pulmonary function studies in five patients with polycythemia vera. As a result of these studies we believe that the following sequence of events may occur in some patients with this disease. (Fig. 1.) The increased blood viscosity and volume causes a decrease in the elasticity and increase in the viscous resistance of the lungs. This in turn leads to a decrease in the maximum breathing capacity and the vital capacity. These changes result in impairment of ventilation which gives rise to a poor correlation between ventilation and perfusion and thereby creates a high distribution gradient with resultant arterial oxygen unsaturation, increased arterial  $p\text{CO}_2$  and respiratory acidosis.

Since polycythemia vera predisposes to thromboses, it is not unreasonable to assume that these may occur in the alveolar capillaries and thereby give rise to a diffusion gradient which would further contribute to the anoxia. Finally, since three of our five cases demonstrated evidence of respiratory center damage, this would be another mechanism further to impair ventilation and thereby contribute to the anoxia. Undoubtedly, this respiratory center damage is in part due to anoxia. However, this would not account for all of it since Case III showed evidence of respiratory center damage while at rest when he had a high arterial  $p\text{CO}_2$  but normal oxygen saturation, and in Case V relief of the anoxia did not improve the function of the respiratory center.

We did not take measurements of the pulmonary artery pressure. However, in view of the development of pulmonary hypertension in normal individuals breathing low oxygen tensions, as demonstrated by Motley et al.,<sup>20</sup> and in view of the accentuated second pulmonic sound in two of our patients (Cases III and V), it is not unreasonable to assume that these patients may go on to pulmonary hypertension and finally to right ventricular failure.

Objective analysis of our findings would admit another possibility. Cases of polycythemia vera,

as we know them now, may represent a heterogeneous group. Some of these patients may actually represent a polycythemic response to medullary center damage of unknown etiology. This could certainly be true of Cases III, IV and V if one would admit that our present technics of studying pulmonary function are not sensitive enough to discern small quantitative changes resulting from such impairment.

#### CASE REPORTS

**CASE I.** A fifty-five year old hat salesman was well until approximately four weeks before admission to the hospital. At that time he noted the gradual onset of painless gangrenous changes of the tips of the distal phalanges of the four medial fingers of the left hand. Because of these changes the patient was hospitalized.

Physical examination on admission revealed a slim, plethoric white male in no acute distress. There was injection of the conjunctivas and slight engorgement of the veins of the fundi. The lungs were clear to percussion and auscultation with normal chest expansion and diaphragmatic excursion. The heart was not enlarged to percussion and there was a regular sinus rhythm with a rate of 82. The blood pressure was 122/88 mm. of mercury in the right arm and 130/85 mm. of mercury in the left arm.  $A_2$  equaled  $P_2$ . No murmurs were heard. The abdomen was negative except for a firm, non-tender spleen palpable two fingerbreadths below the costal margin. Examination of the extremities revealed no clubbing. There was coldness and cyanosis of the distal phalanges of the four medial fingers of the left hand with approximately 5 mm. areas of almost black discoloration at the tips of these fingers.

The hemogram and related data are tabulated. Platelet count was 400,000 and the white blood count 14,700. The remainder of the laboratory findings, including chest x-ray and fluoroscopy, were within normal limits.

Following phlebotomy of 2.85 L. in twenty-one days there was marked subjective improvement and the appearance of the hands returned to normal.

This patient fulfilled all the criteria for the diagnosis of polycythemia vera. He had a polycythemia with leukocytosis, increase in platelets and splenomegaly. In addition, he presented venous thromboses, one of the frequent complications of polycythemia.

**CASE II.** A thirty-three year old white male

had noted lethargy, irritability, frequent epistaxis and a reddish purple color of the skin for three years preceding admission. Physical examination revealed a slightly obese, well developed, young male with plethoric facies. There was no respiratory distress. The eyes showed injection of the conjunctivas and fundi revealed 1 diopter of papilledema with marked distention of the veins. The heart and lungs were within normal limits. The spleen was palpable two fingerbreadths below the costal margin. The remainder of the physical examination was non-contributory.

The hemogram and related data are tabulated. The platelet count was 280,000 and white blood count 9,600. The blood urea nitrogen, fasting blood sugar, urine and urinary 17-ketosteroid excretion were within normal limits. Blood serologic tests were negative. The electrocardiograph was normal.

X-ray examination of the chest revealed the lung fields to be clear and the heart normal in size and configuration. Fluoroscopic examination of the chest demonstrated good diaphragmatic excursion. A bone survey by x-ray was negative and did not demonstrate any osteoporosis.

Following phlebotomies, 2.5 L. in twenty-one days, the patient noted increased energy, marked decrease in his irritability and the plethora was less striking.

This patient fulfilled many of the criteria for the diagnosis of polycythemia vera. He had the typical history and facial appearance of patients with this disease. In addition there was an enlarged spleen and increase in the red blood count, platelet count and plasma volume. One of the features which was missing was the leukocytosis which 50 per cent to 70 per cent of these patients demonstrate.<sup>17</sup> Other causes for polycythemia were sought but none could be found.

**CASE III.** A thirty year old Negro male was admitted to the hospital complaining of abdominal pain of two years' duration. The pain was described as transient, moderately sharp in the right para-umbilical region radiating occasionally to the left upper quadrant. There was no relation to meals, position or bowel movements. Examination for this complaint at its onset two years before admission failed to reveal any significant pathologic disorder. Two months before admission the patient was admitted to another institution, polycythemia was discov-



ered and he was transferred to this hospital for further investigation and therapy. Past history and system review were non-contributory.

Physical examination on admission revealed a well developed and well nourished Negro male in no acute distress. There was an underlying redness in the skin but no cyanosis. The scleras were suffused. The chest revealed no bony deformity and a normal P-A diameter. Respirations were primarily diaphragmatic but with forced breathing there was adequate movement of the chest cage. The breath sounds were vesicular and there were no rhonchi or rales. Examination of the cardiovascular system was within normal limits except for an accentuated  $P_2$ . The blood pressure was 120/80. Abdominal examination revealed no pathologic findings. There was no splenomegaly.

Urine analyses, serology, stool examinations for occult blood, ova and parasites and serum protein determinations were within normal limits. Intravenous pyelogram, gastrointestinal series and barium enema were within normal limits. The hemogram and related data are tabulated. The white blood count was 9,300 with a normal differential, platelets 325,000. Sternal marrow examinations revealed essentially normal findings. Fluoroscopic examination of the chest revealed good diaphragmatic excursions, clear lung fields and a normal cardiovascular silhouette. The electrocardiogram was within normal limits and the electroencephalogram was reported as a normal record in a drowsy patient.

Pulmonary function studies were performed on admission and following phlebotomies totaling 3.7 L. in thirty days there was a striking subjective improvement with a sense of well-being and an increase in work capacity. On examination the plethora disappeared and the previously accentuated  $P_2$  was markedly diminished. One of the striking features before phlebotomy was a failure to increase ventilation following exercise. After completion of the phlebotomies the patient demonstrated a normal ventilatory response to exercise. This point is well demonstrated quantitatively in the tabulated pulmonary function studies.

Polycythemia was noted in this young colored male as an accidental finding. Despite the absence of some of the criteria for the diagnosis of polycythemia vera, viz., splenomegaly and leukocytosis, none of the usual causes for secondary polycythemia could be found. Thus

the diagnosis of polycythemia vera was made by exclusion. The cause for the patient's abdominal complaints was not apparent.

CASE IV. A fifty-four year old white male was admitted to the hospital because of somnolence which had begun four years previously and gradually increased. Prior to hospitalization he slept twenty hours out of twenty-four. He was lethargic, apathetic and his concentration was poor. During gatherings with his friends he would fall asleep. He described his speech as thick and for many months his vision was blurred. In 1920 the patient weighed 145 pounds. He progressively gained weight from this time until admission. During the year before admission he was dyspneic with effort and tired easily. There was no history of chest pain, orthopnea, or nocturnal paroxysmal dyspnea. Five years before admission he had been hospitalized at another institution and at that time his weight was 350 pounds, hemoglobin 18.5 gm. per cent, red blood count 6,500,000 and the hematocrit was 62 per cent.

On examination he was found to be an extremely obese male who was cyanotic and very lethargic. He was comfortable lying flat. The blood pressure was 145/95. There was a bilateral proptosis. The conjunctivas were markedly injected and examination of the fundi disclosed 1 diopter of papilledema with venous engorgement. The breath sounds were diminished throughout and there were a few rales at both lung bases. The heart was normal. Examination of the abdomen was impaired by the marked obesity. The genitalia were normal. There was slight pitting edema and varicosities of both legs. The skin was dusky and there were abdominal striae. There was a stasis dermatitis of the lower limbs. The remainder of the examination was normal.

The hemogram and related data are tabulated. The platelet count was 306,510, and the white blood count 6,550. Urine analysis was negative. The blood urea nitrogen was 18 mg. per cent, Mazzini test negative, fasting blood sugar 97 mg. per cent, serum cholesterol 152 mg. per cent and chlorides 90 mEq./L. The  $CO_2$  combining power was 69 volumes per cent, serum bilirubin 1.0 mg. per cent and urinary ketosteroid excretion was normal. X-ray examination of the chest disclosed normal lung fields and cardiac silhouette. Fluoroscopy confirmed this as well as demonstrating the presence of good diaphragmatic motion. X-ray of the

thoracic and lumbar spine showed marked hypertrophic arthritis. The sella turcica was normal. The electrocardiograph showed a sinus arrhythmia with frequent supraventricular premature contractions.

The patient was initially digitalized because of presumed cardiac failure. However, after a red blood count and hemoglobin were obtained, digitalis was discontinued. Repeated phlebotomies totaling 5.4 L. in thirty-two days were performed with marked subjective and objective improvement. His lethargy lessened considerably. He became much more active and alert. His hours of sleep decreased from approximately twenty hours a day to eight. Electroencephalograms were obtained, one early during the patient's hospitalization and one after treatment. The initial record disclosed evidence of focal vascular lesions. The follow-up electroencephalograph was mildly abnormal due to scattered slow waves in the occipital and central areas but showed a definite improvement.

The history and physical findings of this patient were consistent with progressive polycythemia and obesity. The spleen was not palpable but this examination was impaired by his pendulous abdomen. The hematocrit, hemoglobin and red blood count were high. The blood volume was not determined because of technical difficulties. It was thought that this patient probably represented a case of polycythemia vera although there was no leukocytosis and the presence or absence of splenomegaly could not be ascertained.

**CASE V.** A thirty-two year old male was first admitted in April, 1948, complaining of fatigue and lassitude. At that time the history revealed that in 1943, while in the army, the patient noted reddish blue discoloration of the face. He was hospitalized in an army hospital in which red blood counts of eight to nine million were recorded and the patient was given a medical discharge in 1945 with a diagnosis of polycythemia vera. During his first admission to the hospital he was studied for possible Cushing's syndrome, brain tumor, congenital heart disease and pulmonary disease with negative results. Two arterial oxygen saturation tests at rest done at that time showed 96 per cent saturation. The final discharge diagnosis was polycythemia vera. The patient was readmitted in June, 1950, because of increasing discoloration of the face and an incapacitating degree of lethargy.

On examination the patient was moderately obese and cyanotic but was in no respiratory distress. The fundi showed distended, tortuous veins and blurring of the disc margins. The chest was thick and obese with good mobility and examination of the lungs disclosed a few basilar fine moist rales bilaterally. The area of cardiac dullness was not percussible, the rhythm was regular,  $P_2$  was markedly accentuated and there were no murmurs. The abdomen was obese and no organs or masses could be felt.

The hemogram and related data are tabulated. The white blood cell and platelet counts ranged about 7,000 and 200,000, respectively. Blood serologic tests were negative. The urine contained negative to 2 plus protein with occasional hyaline and granular casts. A glucose tolerance test and 17-ketosteroid excretion were normal. The blood sodium, blood urea nitrogen and serum proteins were within normal limits. A bone survey by x-ray failed to demonstrate any osseous abnormalities. The chest x-ray revealed normal lung fields and cardiovascular silhouette. Fluoroscopy demonstrated good diaphragmatic excursion. An electrocardiogram showed non-specific T wave changes. The venous pressure was 115 mm. of  $H_2O$  and circulation time arm to tongue, using decholin, was fourteen seconds.

The hospital course was uneventful. Following phlebotomy, of 4.8 L. in sixty days, there was a moderate but not striking decrease in the patient's symptomatology.

This patient represented the most complicated case which was studied. Here again, polycythemia vera was diagnosed by exclusion since the patient never demonstrated leukocytosis or thrombocytosis and the obesity precluded adequate examination for splenomegaly. In addition, before phlebotomy, although the red blood cell mass was increased the plasma volume was decreased. In view of the work of Berlin et al.<sup>10</sup> the values for plasma volume cannot be used for differentiating primary from secondary polycythemia. An adequate explanation for the polycythemia could not be found and in view of the normal oxygen saturation three years before the present admission at a time when the patient was polycythemic, we felt justified in considering him a case of polycythemia vera.

#### SUMMARY

Detailed pulmonary function studies are reported in five cases of polycythemia, probably

polycythemia vera. The results would indicate that some of these patients may show incapacitating changes in directions not previously indicated. Thus definite changes in ventilation and gas exchange may occur, with the development of severe degrees of anoxia. The anoxia and other unknown factors result in respiratory center damage and a vicious cycle may be initiated. Since these changes were not reversible in all our cases, early and adequate therapy would seem to be imperative in patients with polycythemia vera.

The possibility is pointed out that polycythemia vera may be a heterogeneous disease with some of the cases representing a polycythemic response to primary respiratory center damage of undetermined etiology.

*Acknowledgment:* The authors wish to acknowledge their appreciation to Drs. Robert F. Loeb and Richard L. Riley for their many helpful comments in the preparation of this manuscript.

## REFERENCES

1. ISAACS, R. Pathologic physiology of polycythemia vera. *Arch. Int. Med.*, 31: 289, 1923.
2. BROOKS, W. D. W. Circulatory adjustments in polycythemia rubra vera. *Proc. Roy. Soc. Med.*, 29: 1379, 1936.
3. ALTSCHULE, M. D., VOLK, M. C. and HENSTELL, H. Cardiac and respiratory function at rest in patients with uncomplicated polycythemia vera. *Am. J. M. Sc.*, 200: 478-483, 1940.
4. STEWART, H. J., WHEELER, C. H. and CRANE, N. F. Circulatory adjustments in polycythemia vera. *Am. Heart J.*, 21: 511, 1941.
5. HARROP, G. A., JR. and HEATH, E. H. Pulmonary gas diffusion in polycythemia vera. *J. Clin. Investigation*, 4: 53-70, 1927.
6. WASSERMAN, L. R., DOBSON, R. L. and LAWRENCE, J. H. Blood oxygen studies in patients with polycythemia and in normal subjects. *J. Clin. Investigation*, 28: 60-65, 1949.
7. VON BERGMANN, G. and PLESCH, J. Über Hyperglobulie. *München. med. Wchnschr.*, 58: 1849, 1911.
8. HITZENBERGER, K. Die Sauerstoffsättigung des arteriellen Blutes bei Polycythemien. *Ztschr. f. klin. Med.*, 126: 495, 1934.
9. ROVER, W. Über Hyperglobulie. *München. med. Wchnschr.*, 58: 2791, 1911.
10. BERLIN, N. I., LAWRENCE, J. H. and GARTLAND, J. Blood volume in polycythemia as determined by  $P^{32}$  labeled red blood cells. *Am. J. Med.*, 9: 747, 1950.
11. BALDWIN, E., Cournand, A., and RICHARDS, D. W., JR. Pulmonary insufficiency. I. Physiological classification, clinical methods of analysis, standard values in normal subjects. *Medicine*, 27: 243, 1948.
12. RILEY, R. L., PROEMMEL, D. D. and FRANKE, R. E. A direct method for determination of oxygen and carbon dioxide tensions in blood. *J. Biol. Chem.*, 161: 621, 1945.
13. VAN SLYKE, D. D. and NEILL, J. M. The determination of gases in blood and other solutions by vacuum extraction and manometric measurements. *J. Biol. Chem.*, 61: 523, 1924.
14. SCHOLANDER, P. F. Analyzer for accurate estimation of respiratory gases in one half cubic centimeter samples. *J. Biol. Chem.*, 167: 235, 1947.
15. RILEY, R. L. and Cournand, A. Analysis of ventilation perfusion relationships in the lungs. *Federation Proc.*, 8: 132, 1949.
16. VAN SLYKE, D. D. and SENDROY, J., JR. Line charts for graphic calculations by the Henderson-Hasselbalch equation, and for calculatory plasma carbon dioxide content from whole blood content. *J. Biol. Chem.*, 79: 781, 1928.
17. WINTROBE, M. M. *Clinical Hematology*, p. 595. Philadelphia, 1946. Lea & Febiger.
18. RAHN, H. Effect of unequal blood flow and ventilation upon the alveolar-arterial gradient. *Federation Proc.*, 8: 129, 1949.
19. RILEY, R. L. Personal communication.
20. MOTLEY, H. L., Cournand, A., WEEKS, L. and DRESDALE, D. The influence of short periods of induced acute anoxia upon pulmonary artery pressure in man. *Am. J. Physiol.*, 150: 315, 1947.



# Effect of Cortisone and ACTH on Fluid and Electrolyte Distribution in Man\*

MARVIN F. LEVITT, M.D.† and MORTIMER E. BADER, M.D.

*New York, New York*

PREVIOUS evidence indicates that the adrenal hormones may influence the distribution of fluid between cells and the extracellular compartment.<sup>1-8</sup> It is not clear whether this redistribution is limited to water and is achieved solely by a primary change in exogenous sodium balance with its attendant osmotic effect or whether it is partly a consequence of an endogenous shift of salt as well as water between the cells and the extracellular compartment. Swingle and his co-workers deduced that adrenal cortical extract increased the isotonic extracellular volume in adrenalectomized, fasted dogs and suggested, therefore, that the adrenal hormone evoked an endogenous shift of salt and water into the extracellular compartment.<sup>1-3</sup> Gaudino and Levitt demonstrated that synthetic desoxycorticosterone produced large but reversible shifts of fluid from within the cells outward whereas adrenalectomy induced the opposite effect.<sup>7</sup> The fact that total body water and serum electrolyte concentrations remained essentially unchanged in their experiments likewise suggested that an endogenous shift of salt and water had occurred. Harrison and Darrow, calculating tissue water compartments upon the assumption that chloride is limited to extracellular distribution, demonstrated that during progressive adrenal insufficiency water shifted from the extracellular to the intracellular compartment but concluded that these water shifts were entirely due to urinary sodium loss and decreased extracellular sodium concentration.<sup>5,6</sup>

The availability of cortisone and ACTH has further stimulated interest in the influence of adrenal activity on electrolyte metabolism. It has been well demonstrated that these agents frequently produce salt retention,<sup>9,10</sup> but it has also been noted that edema and circulatory

failure may occur in patients maintained on salt-free diets.<sup>11</sup> As in the case of the other adrenal hormones, it is possible that such effects may result from endogenous salt and water shifts. It seemed desirable, therefore, to study the influence of cortisone and ACTH on fluid and electrolyte distribution in patients maintained on salt-free diets so that primary retention of sodium chloride is minimized if not excluded.

## METHODS

Serial measurements of plasma volume, inulin space, serum electrolyte concentrations, and inulin and PAH clearances were performed in six patients before and during the course of cortisone or ACTH therapy. Three of the patients were treated with 100 to 150 mg. of cortisone and three with 100 mg. of ACTH daily for periods of ten to twenty days. Throughout the control period (four to seven days) and the period of hormone treatment the patients were maintained on a rice-fruit diet.† The first set of measurements was performed on the last day of the control period and was repeated serially every three to four days throughout treatment. In five of these patients total daily urinary sodium, chloride and potassium outputs were determined. In one patient mercurhydrin was injected daily for three days starting at the sixth day of ACTH treatment but no other phase of the protocol was altered. The patients were weighed daily and adherence to the constant diet was carefully supervised.

Inulin space was measured as previously described.<sup>14-16</sup> Constant priming doses and

† Analyses of the rice diet revealed a daily electrolyte intake of 3, 6 and 40 mEq. of sodium, chloride and potassium, respectively, results which compare favorably with previous analyses of the rice diet.<sup>12,13</sup>

\* From the Departments of Medicine and Pediatrics, Mt. Sinai Hospital, New York, N.Y. Part of these results have been presented in abstract form: LEVITT, M. F. and BADER, M. E. Influence of cortisone and ACTH on water and electrolyte distribution and renal function. *Federation Proc.*, 10: 82, 1951.

† Rosenstock Fellow, Sarah Welt Fellow in Medicine.

identical infusion rates and concentrations were used in an individual patient throughout the course of a study. The infusion was maintained at a constant rate of  $0.8 \pm 0.016$  cc./min. for five and a half hours. At the end of that time the bladder was washed and the infusion discontinued simultaneously. Thereafter the urine was collected until all the inulin had been excreted (eighteen hours). Three plasma samples were drawn in the ninety-minute interval directly preceding cessation of the infusion. The total amount of inulin excreted after stopping the infusion, divided by the average equilibrium plasma concentration, was taken as the volume of distribution of inulin. Prior to the inulin infusion a plasma blank and a timed urinary specimen were collected for appropriate blank corrections. Virtually all the injected inulin was recovered in the urine (94 to 102 per cent). The quantity of infused sodium chloride that was retained during each space measurement did not exceed 3 per cent of the original store of extracellular sodium and chloride and has been included in the balance data. Inulin and p-aminohippuric acid (PAH) clearances were determined during the final one to two hours of the equilibrating infusion according to the method of Smith and his co-workers.<sup>17</sup>

Analyses for inulin were performed by the Schreiner modification of the resorcinol method<sup>18</sup> and blood glucose and urinary-reducing activity were measured separately in each experiment. No significant hyperglycemia or glycosuria was noted during the course of these observations. Furthermore, the serum and urinary blank corrections preclude any error from this source. PAH determinations were made by the method of Bratton and Marshall.<sup>19</sup> Plasma volume was measured by the T-1824 technic of Gregerson and Stewart with determinations based on a single ten-minute sample.<sup>20</sup> Plasma sodium and potassium concentrations were measured on fasting pre-infusion samples with an internally compensated Perkin-Elmer flame photometer. All such samples from a single patient were analyzed simultaneously at the completion of therapy so that relative changes might be apparent. Urinary sodium and potassium concentrations were likewise measured photometrically. Chloride analyses were performed by the method of Schales and Schales.<sup>21</sup> Total extracellular electrolyte was calculated as the product of the inulin space and plasma concentration. No

corrections were made for the water content of serum or for Donnan's equilibrium.

#### RESULTS

Cortisone and ACTH induced in six patients a progressive increase in the inulin space of distribution. This increase reached its peak after eight to nine days of therapy and averaged 30 per cent (range of 20 to 38 per cent) above control values in the three patients treated with cortisone and 23 per cent (21 to 25 per cent) in the three treated with ACTH. (Table 1, Figs. 1 and 2.) The average increase for all six patients equalled 27 per cent. The expansion of inulin space in the cortisone-treated patients was apparent as early as two or three days after the onset of treatment (Cases II and VI) and in all cases was coexistent with a weight loss of 1 to 2 kg. Changes in plasma sodium and chloride were not remarkable although there seemed to be a slight but consistent increase in sodium concentration and either no change or a slight decrease in chloride. The increase in total extracellular sodium and chloride, therefore, approximated the proportionate increase in inulin space. As noted in Table 1 and Figure 3, the positive sodium chloride balance accounted for only 10 to 20 per cent of the total increase in extracellular electrolyte. If the sodium chloride accumulated from the slight positive balance (including the retained portions of the 50 mEq. of the sodium chloride infused during each inulin space determination) is subtracted from the absolute extracellular electrolyte increase, a substantial moiety remains. This increase in extracellular sodium in the five patients in whom balance data are available averaged 25 per cent, with specific percentages increases for Cases I, II, III, V and VI, respectively, equal to 34, 33, 22, 20 and 20 per cent. Similar increase in extracellular chloride averaged 19 per cent with the individual increments for the same patients, respectively, 28, 21, 22, 11 and 12 per cent. In four patients in whom treatment was continued beyond eight or nine days the inulin space reverted to control values by the twelfth to fifteenth day despite continued therapy (Cases II, III, V and VI). Thereafter, continuation of treatment did not further influence the volume of distribution of inulin. (Table 1.) The decrease in extracellular electrolyte which accompanied the decrease in inulin space was not associated with a comparable negative electrolyte balance. The unaccounted for decrease in extracellular

sodium or chloride, therefore, approximated in magnitude the increase detected earlier in treatment. (Figs. 1, 2 and 3.)

Mercuhydrin was injected daily during the sixth to ninth days of ACTH therapy in Case I.




FIG. 1. The influence of cortisone on body weight, inulin space, plasma volume and renal function in Case VI. All indices except weight are expressed as percentile deviation from control.




FIG. 2. The influence of ACTH on body weight, inulin space, plasma volume and renal function in Case III. All indices except weight are expressed as percentile deviation from control.

Despite a moderate sodium and chloride diuresis little change was noted in the inulin space.

In three of the five patients (Cases I, II and III) in whom balance studies were performed a negative potassium balance of about 20 mEq./




FIG. 3. The average absolute changes in total extracellular electrolyte during cortisone and ACTH therapy compared to simultaneous changes in exogenous balance.

day was noted during the first two or three days of hormone treatment. Thereafter no demonstrable negative or positive potassium balance was present except in the patient who received mercurial diuretics. Here a negative balance of 125 mEq. was produced during the three-day period of electrolyte diuresis.

The changes in plasma volume were conspicuously small and probably within the range of experimental error. There was an increment in plasma volume which averaged 6 per cent, with a range from 2 to 10 per cent. (Table I.)

Renal function studies revealed a progressive increase of filtration rate in six of seven patients. (Table I.)\* Simultaneously, PAH clearances showed variable changes with slight increase in some and decrease in others. Cortisone and ACTH therapy, therefore, produced an increase in filtration fraction in each patient. The average increase in glomerular filtration rate was 31 per cent, in filtration fraction 23 per cent. Peak increases in renal function generally coincided with maximal expansion of the inulin space. Renal function reverted to control or subcontrol levels as adrenal therapy was continued beyond eight or nine days.

\* The renal function data obtained without coincident space measurements are not included in the tables.



TABLE I  
EFFECTS OF ACTH AND CORTISONE ON THE DISTRIBUTION OF WATER AND ELECTROLYTES IN THE BODY FLUIDS

Observation	Weight (kg.)	Inulin Space		Serum Electrolytes (mEq./L.)			Total Extracellular Na (mEq.)	Exogenous Na Balance between Observations (mEq.)	Total Extracellular Cl (mEq.)	Exogenous Cl Balance between Observations (mEq.)	Unaccounted for Change in Extracellular Electrolytes (mEq.)		Plasma Volume (cc.)	Renal Function		
		(L.)	% of Body Wt.	Na	K	Cl					Na ( $\Delta a-b$ )	Cl ( $\Delta c-d$ )		$C_{In}$	$C_{PAH}$	F. F. (%)
Case I; female; age 52; non-toxic adenoma of the thyroid																
Control	57.7	10.0	17.3	147	3.9	107	1,470		1,070				3,170	97	492	19.7
6 days of ACTH, 100 mg./day	56.8	12.3	21.6	149	4.2	106	1,830	+38	1,300	+22	+322	+208	3,290	110	502	21.9
9 days of ACTH, 100 mg./day, 3 days of daily mer- captopril, 2 cc. i.m.	55.0	12.1	22.0	149	4.1	94	1,800	-213	1,140	-252	+183	+92	3,350	149	620	24.0
										Total	+305	+300				
Case II; female; age 38; bronchial asthma																
Control	58.0	13.3	22.9	146	4.2	109	1,940		1,450				2,270	138	440	31.4
2 days of corti- sone, 150 mg./ day	57.4	15.7	27.4	146	3.8	107	2,290	+38	1,680	+41	+312	+189		141		
6 days of corti- sone, 150 mg./ day	57.2	16.4	28.7	149	3.6	103	2,440	+48	1,690	+45	+102	-35	2,560	142	393	36.2
9 days of corti- sone, 150 mg./ day	56.5	17.9	31.6	151	3.2	104	2,700	+39	1,860	+18	+221	+152	2,440	194	450	43.2
										Total	+635	+305				
15 days of corti- sone, 150 mg./ day	56.0	12.0	21.4	148	3.7	107	1,770	+20	1,280	-12	-950	-568	2,520	90	370	24.3
20 days of corti- sone, 150 mg./ day	55.7	12.3	22.0	147	3.7	105	1,810	+35	1,290	+14	+5	-4	2,440	104	355	29.3

Case III; female; age 26; no disease												
Control	54.3	11.0	20.2	151	4.1	110	1,660	1,210	2,570	103	500	20.6
6 days of ACTH, 100 mg./day	53.2	12.1	22.8	154	3.9	110	1,860	+36	+164	122	592	20.6
9 days of ACTH, 100 mg./day	52.5	13.7	26.1	152	3.8	110	2,080	+19	+201 +268	142	550	25.8
15 days of ACTH, 100 mg./day	51.8	11.4	22.0	150	3.7	108	1,710	-54	-316	135	560	24.1
20 days of ACTH, 100 mg./day	51.2	11.4	22.2	146	3.2	105	1,670	-10	- 30	116	490	23.7
Case IV; * female; age 21; bronchial asthma												
Control	56.0	8.600	15.4									
(cc.)												
9 days of corti- sone, 100 mg./ day orally	54.0	11,900	22.0									
(cc.)												
Case V; female; age 45; mild hyperthyroidism												
Control	61.0	145	4.1	107								
3 days of ACTH, 100 mg./day	60.5	9.5	15.7	144	105	1,370	998			128	480	26.7
6 days of ACTH, 100 mg./day	59.4	10.5	17.7	144	3.9	105	1,510	+24	+116	128	515	24.9
8 days of ACTH, 100 mg./day	58.8	11.5	19.6	147	4.0	102	1,690	+20	+160 +108	147	500	29.4
12 days of ACTH, 100 mg./day	57.8	8.7	15.0	141	4.1	109	1,230	-52	-408	110	502	21.9
Case VI; female; age 21; no disease												
Control	63.0	9.5	15.1	146	4.4	110	1,370	1,050		120	665	18.1
3 days of corti- sone, 150 mg./day orally	62.0	10.6	17.1	152	4.0	108	1,610	+10	+210	115	552	20.8
8 days of corti- sone, 150 mg./day	60.7	11.4	18.8	148	4.0	106	1,690	+26	+ 54 +264	121	639	19.0
13 days of corti- sone, 150 mg./day	59.6	9.3	15.6	145	4.1	107	1,350	-40	-200	93	558	16.7

$C_{IN}$  = Plasma clearance of inulin in cc./min.  $C_{PAH}$  = Plasma clearance of para-aminohippuric acid in cc./min. F. F. = Filtration fraction.  
 \*No balance studies were performed in this patient.

The changes produced by cortisone and ACTH were qualitatively similar. Because of variation in dosage, the undetermined factor of dosage equivalence and the inherent limitations of the methods it is impossible to conclude that these agents produce quantitatively different responses. Consequently, the data from the two groups of patients have been treated as essentially one group of experimental observations.

#### COMMENTS

Available evidence suggests that the space of distribution of inulin affords the most reliable estimate yet made of the true extracellular volume.<sup>14-16,22,47</sup> This conclusion is deduced from two separate lines of investigation; one purporting to demonstrate that inulin is completely excluded from the cells, and the other that it is freely diffusible throughout the extracellular compartment. That inulin does not enter cells seems probable *a priori* in view of its average molecular weight (5,000),<sup>23</sup> its elongate structural configuration<sup>24</sup> and its lipoid insolubility. The inulin molecule has been shown to be excluded from erythrocytes,<sup>25</sup> bile,<sup>26</sup> gastric juice<sup>27</sup> and renal tubular cells.<sup>25</sup> No enzymes are known to be present in man which are capable of hydrolyzing it,<sup>25</sup> and its rapid and quantitative recovery in the urine<sup>14,16,28</sup> argues against any significant storage or metabolism in the living organism. Its space of distribution is smaller than simultaneous volumes of other substances (thiocyanate, sodium, chloride, mannitol) used for the measurement of extracellular fluid, most of which are now known to enter cells or undergo partial metabolism.<sup>22,29-31</sup> The volume of distribution of inulin, as a measure of extracellular fluid, is therefore more readily compatible with recent estimates of total body water.<sup>32,33</sup>

The alternate possibility of incomplete diffusion into the extracellular compartment is made improbable by two sets of experimental observations: (1) Once the minimum infusion time necessary to establish equilibrium distribution of inulin has been established (two hours in the dog, five hours in man), further prolongation of the infusion does not increase its space of distribution.<sup>15,16</sup> (2) The volumes of distribution of sucrose and inulin in the nephrectomized rabbit and rat,<sup>26,34</sup> and mannitol (with corrections for metabolism) and inulin in the human are identical.<sup>22,31,35</sup> The equality of the spaces of distribution of molecules of such great variation in size offers strong evi-

dence against the possibility of extracellular compartmentalization.

In the experiments herein described cortisone and ACTH produced significant increases in the inulin space. That these changes were not attributable to the metabolism of inulin is confirmed by the virtually complete recovery of the infused inulin which was achieved in each experiment. Inasmuch as these alterations were not associated with comparable weight changes or exogenous sodium or chloride balance only two alternative explanations are possible: (1) Cortisone and ACTH induce an internal redistribution of salt and water. (2) These agents produce an alteration in cellular permeability so that the inulin molecule is temporarily accessible to a small proportion of body cells (10 to 15 per cent). The second alternative is rendered unlikely by the evidence cited to support the hypothesis that inulin is excluded from cells. Furthermore, it seems improbable that a carbohydrate molecule of such size may enter cells without undergoing breakdown or metabolism. This alternate explanation, however, cannot be completely excluded at this time.

If it is accepted that the inulin volume of distribution provides a reliable estimate of extracellular volume, it follows as an immediate corollary that cortisone and ACTH induce an internal redistribution of salt and water. The following considerations are based on this premise:

The endogenous sodium and chloride emptied into the extracellular compartment during the first eight to nine days of cortisone or ACTH therapy increased the total amount of extracellular electrolyte by 25 and 19 per cent, respectively. These average figures define the proportion of sodium and chloride originally contained within the extracellular compartment as approximately 80 and 84 per cent, respectively, of the amounts contained therein at the point of peak response. Previous observations regarding the proportion of extracellular sodium to total body sodium (simultaneous inulin/sodium space ratios) have yielded average ratios of 0.62 in man and dog.<sup>15,16,36</sup> Similar inulin/chloride space ratios in the dog (not performed simultaneously) have averaged 0.76,<sup>15,37</sup> or if, as previously suggested, bromide be used as a measure of total body chloride, 0.67.<sup>14,38</sup> No simultaneous over-all inulin/chloride space ratios are available in man. Recently, however,



it has been shown that the inulin/chloride space ratio in human muscle tissue averages 0.82<sup>39</sup> despite the circumstance that muscle cells are relatively chloride-free.<sup>29</sup> Simultaneous inulin/bromide space ratios in man approximate 0.60 to 0.70.<sup>16,33</sup> Consequently, the quantity of sodium and chloride contained outside of the inulin space appears entirely adequate to account for the average changes described. The magnitude of the chloride shift was proportionately smaller than the simultaneous sodium movement, possibly as a result of the disparity between the proportion of non-extracellular chloride and sodium.

The artificial diuresis produced by mercurhydrin in Case 1 did not obscure the hormone-induced changes. The mercurials were administered before the expected peak increase in inulin space and, although a moderate sodium and chloride diuresis was induced, the fall in total extracellular electrolyte was disproportionately small. The artificial diuresis, therefore, did not prevent the movement of sodium and chloride into the extracellular compartment.

In three patients a mild and transient negative potassium balance occurred during the early phase of the sodium and chloride shift into the extracellular compartment. Otherwise no clear-cut relation was observed between the redistribution of sodium and chloride and exogenous potassium balance.

These experiments do not reveal the precise source of the endogenous water, sodium and chloride which are diverted into the extracellular compartment. The known calcium and phosphorus mobilization which is induced by the adrenal steroids<sup>10,40</sup> suggests that some of the sodium may derive from bone, which is known to contain appreciable stores of sodium. The quantity of salt-containing fluid present in the small intestine is another potential source and, of course, some tissue cells may contribute a significant proportion. The mechanism of reversal of this redistribution with continued cortisone or ACTH therapy is also unknown.

The changes in plasma volume are extremely small and possibly within the range of experimental error. What may be of significance is that each patient showed an increase, albeit small, coincident with the expansion of the extracellular volume. Certainly, as has been previously noted, plasma volume does not offer a sensitive index of changes in the extracellular volume.<sup>7,41</sup>

In six of the seven patients studied there was a significant increase in filtration rate and, in all seven, an increase in filtration fraction similar to observations previously recorded.<sup>42-44</sup> It has been shown that the rice diet itself may produce alterations in renal function but in opposite direction from those observed here.<sup>45,46</sup> This circumstance may account for the sub-control values noted after twelve to fifteen days of cortisone or ACTH therapy. Although the peak changes in renal function and inulin space generally coincided, it does not follow that they are causally related. In Case 1 further expansion of the extracellular volume was inhibited by mercurial diuretics but a progressive change in renal function occurred nevertheless. This observation suggests that the adrenal hormones may influence body water and electrolyte distribution and renal function simultaneously but independently.

#### SUMMARY

1. Cortisone and ACTH induce a transient shift of water, sodium and chloride into the measurable extracellular volume (inulin space).
2. This redistribution reaches its peak after eight to nine days of cortisone or ACTH therapy and then regresses despite continuation of therapy.
3. Cortisone and ACTH produce a progressive increase in glomerular filtration rate and filtration fraction. The maximal changes in body fluid distribution and renal function coincide.

*Acknowledgment:* The authors wish to express their appreciation to Dr. Jonas H. Sirota for his assistance in these studies and to Dr. W. P. Anslow, Jr., for his advice in analysis of the data.

#### REFERENCES

1. SWINGLE, W. W., PFIFFNER, J. J., VARS, H. M. and PARKINS, W. M. The effect of fluid deprivation and fluid intake upon the revival of dogs from adrenal insufficiency. *Am. J. Physiol.*, 108: 144, 1934.
2. SWINGLE, W. W., PARKINS, W. M., TAYLOR, A. R. and HAYS, H. W. Relation of serum sodium and chloride levels to alterations of body water in the intact and adrenalectomized dog, and the influence of adrenal cortical hormone upon fluid distribution. *Am. J. Physiol.*, 116: 438, 1936.
3. SWINGLE, W. W., PARKINS, W. M., TAYLOR, A. R. and HAYS, H. W. A study of water intoxication in the intact and adrenalectomized dog and the influence of adrenal cortical hormone upon fluid and electrolyte distribution. *Am. J. Physiol.*, 119: 557, 1937.
4. HARROP, G. A. The influence of the adrenal cortex

- upon the distribution of body water. *Bull. Johns Hopkins Hosp.*, 59: 11, 1936.
5. HARRISON, H. E. and DARROW, D. C. The distribution of body water and electrolytes in adrenal insufficiency. *J. Clin. Investigation*, 17: 77, 1938.
  6. DARROW, D. C., HARRISON, H. E. and TAFFEL, M. Tissue electrolytes in adrenal insufficiency. *J. Biol. Chem.*, 130: 487, 1939.
  7. GAUDINO, M. and LEVITT, M. F. Influence of adrenal cortex on body water distribution and renal function. *J. Clin. Investigation*, 28: 1487, 1949.
  8. STERN, T. N., COLE, V. V., BASS, A. C. and OVERMAN, R. R. Dynamic aspects of sodium metabolism in experimental adrenal insufficiency using radioactive sodium. *Am. J. Physiol.*, 164: 437, 1951.
  9. PERERA, G., PINES, K. L., HAMILTONE, H. B. and VISLOCKY, K. Clinical and metabolic study of 11 dehydro-17 hydroxy corticosterone acetate in hypertension, Addison's disease and diabetes mellitus. *Am. J. Med.*, 7: 56, 1949.
  10. SPRAGUE, R., POWER, M. H., MASON, H. L., ALBERT, A., MATHIESON, D. R., HENCH, P. S., KENDALL, E. C., SLOCUMB, C. H. and POLLEY, H. J. Observations on the physiologic effects of cortisone and ACTH in man. *Arch. Int. Med.*, 85: 199, 1950.
  11. SOFFER, L. J., LEVITT, M. F. and BAEHR, G. The influence of cortisone and ACTH in acute disseminated lupus erythematosus. *Arch. Int. Med.*, 86: 558, 1950.
  12. KEMPNER, W. Treatment of heart and kidney disease and of hypertensive and arteriosclerotic vascular disease with the rice diet. *Ann. Int. Med.*, 31: 821, 1949.
  13. DOLE, V. P., DAHL, L. K., COTZIAS, G. C., EDER, H. A. and KREBS, M. E. Dietary treatment of hypertension. Clinical and metabolic studies of patients on the rice-fruit diet. *J. Clin. Investigation*, 29: 1189, 1950.
  14. GAUDINO, M., SCHWARTZ, I. L. and LEVITT, M. F. Inulin volume of distribution as a measure of extracellular fluid in dog and man. *Proc. Soc. Exper. Biol. & Med.*, 68: 507, 1948.
  15. GAUDINO, M. and LEVITT, M. F. Inulin space as a measure of extracellular volume. *Am. J. Physiol.*, 157: 387, 1949.
  16. SCHWARTZ, I. L., SCHACTER, D. and FREINKEL, N. The measurement of extracellular fluid in man by means of a constant infusion technique. *J. Clin. Investigation*, 28: 1117, 1949.
  17. SMITH, H. W., FINKELSTEIN, N., ALIMINOSA, L., CRAWFORD, B. and GRABER, M. The renal clearances of substituted hippuric acid derivatives and other aromatic acids in dog and man. *J. Clin. Investigation*, 24: 388, 1945.
  18. SCHREINER, G. Determination of inulin by means of resorcinol. *Proc. Soc. Exper. Biol. & Med.*, 74: 117, 1950.
  19. BRATTON, A. C. and MARSHALL, E. K., JR. A new coupling component for sulfanilamide determination. *J. Biol. Chem.*, 128: 537, 1939.
  20. GREGGSON, M. I. and STEWART, J. D. Simultaneous determination of the plasma volume with T-1824 and the "available" fluid volume with sodium thiocyanate. *Am. J. Physiol.*, 125: 142, 1939.
  21. SCHALES, O. and SCHALES, S. S. A simple and accurate method for the determination of chlorides in biological fluids. *J. Biol. Chem.*, 140: 897, 1947.
  22. LEVITT, M. F. and GAUDINO, M. Measurement of body water compartments. *Am. J. Med.*, 9: 208, 1950.
  23. WESTFALL, B. B. and LANDIS, E. M. The molecular weight of inulin. *J. Biol. Chem.*, 116: 727, 1936.
  24. BUNIM, J. J., SMITH, W. W. and SMITH, H. W. The diffusion coefficient of inulin and other substances of interest in renal physiology. *J. Biol. Chem.*, 118: 667, 1937.
  25. SMITH, H. W. *The Physiology of the Kidney*. New York, 1937. Oxford University Press.
  26. KRUHÖFFER, P. Inulin as an indicator for the extracellular space. *Acta physiol. Scandinav.*, 11: 16, 1946.
  27. BERGER, E. Y. and DUNNING, M. Personal communication.
  28. SMITH, W. W., FINKELSTEIN, N. and SMITH, H. W. Renal excretion of hexitols and their derivatives and of endogenous creatinine like chromagen in dog and man. *J. Biol. Chem.*, 135: 231, 1940.
  29. MANERY, J. F. and HASTINGS, A. B. The distribution of electrolytes in mammalian tissues. *J. Biol. Chem.*, 127: 657, 1939.
  30. CRANDALL, L. A. and ANDERSON, M. X. Estimation of the state of hydration of the body by the amount of water available for the solution of sodium thiocyanate. *Am. J. Digest. Dis.*, 1: 126, 1934-1935.
  31. DOMINGUEZ, R., CORCORAN, A. C. and PAGE, I. H. Mannitol: kinetics of distribution, excretion and utilization in human beings. *J. Lab. & Clin. Med.*, 32: 1192, 1937.
  32. SOBERMAN, R., BRODIE, B. B., LEVY, B. B., AXELROD, J., HOLLANDER, V. and STEEL, J. M. The use of antipyrine in the measurement of total body water. *J. Biol. Chem.*, 179: 31, 1949.
  33. BERGER, E. Y., DUNNING, M. F., STEELE, J. M., JACKENTHAL, R. and BRODIE, B. B. Estimation of intracellular water in man. *Am. J. Physiol.*, 162: 318, 1950.
  34. WILDE, W. S. The chloride equilibrium in muscle. *Am. J. Physiol.*, 143: 666, 1945.
  35. SCHWARTZ, I. L. Measurement of extracellular fluid by means of a constant infusion technique without collection of urine. *Am. J. Physiol.*, 160: 526, 1950.
  36. CRAWFORD, B. and GAUDINO, M. Personal communication.
  37. WINKLER, A. W., ELKINTON, J. R. and EISENMAN, A. Comparison of sulfocyanate with radioactive chloride and sodium in the measurement of extracellular fluid. *Am. J. Physiol.*, 139: 239, 1943.
  38. WALLACE, G. B. and BRODIE, B. B. The distribution of administered bromide in comparison with chloride and its relation to body fluids. *J. Pharmacol. & Exper. Therap.*, 65: 214, 1939.
  39. MOKOTOFF, R. and ROSS, G. Personal communication.
  40. SOFFER, L. J., GABRILOVE, J. L. and JAILER, J. W. ACTH in Cushing's syndrome and in virilism. *J. Clin. Endocrinol.*, 10: 594, 1950.
  41. WARREN, J. V., MERRILL, A. J. and STEAD, E. A. The role of the extracellular fluid in the maintenance of a normal plasma volume. *J. Clin. Investigation*, 22: 635, 1943.

42. HELLMAN, L., WESTON, R. E., ESCHER, D. J. and LEITER, L. The effect of adrenocorticotropin on renal hemodynamics and uric acid clearance. *Federation Proc.*, 7: 52, 1948.
43. INGBER, S. J., RELMAN, A. S., BURROWS, B. A., KASS, E. H., SISSON, J. H. and BURNETT, C. H. Changes in normal renal function resulting from cortisone and ACTH. *J. Clin. Investigation*, 29: 824, 1950.
44. EARLE, D., FARBER, S. J., ALEXANDER, J. D. and PELLAGRINO, E. J. Observations on the relation of renal function changes to the electrolyte and glycosuric effects of ACTH. Proc. Second ACTH Conference, 1950. (In press.)
45. WESTON, R. E., HELIMAN, L., ESCHER, D. J. and LEITER, L. The effect of low sodium and Kempner diets on renal hemodynamics and electrolyte excretion in hypertensives. *Federation Proc.*, 7: 132, 1948.
46. CHASIS, H., GOLDRING, W., BREED, E. S., SCHREINER, G. E. and BOLOMEY, A. A. Salt and protein restriction. Effects on blood pressure and renal hemodynamics in hypertensive patients. *J. A. M. A.*, 142: 711, 1950.
47. SMITH, H. W. *The Kidney: Structure and Function in Health and Disease*. New York, 1951. Oxford University Press.



# Experience with Methimazole (Tapazole) in the Treatment of Hyperthyroidism\*

## *A Report of Thirty-five Cases*

BERNARD L. HALLMAN, M.D. and PHILIP K. BONDY, M.D.

Atlanta, Georgia

THE introduction of antithyroid drugs into the treatment of hyperthyroidism has greatly improved the prognosis and course of patients with this disease. The earliest substances used as antithyroid agents proved to be moderately toxic, but recently, with the intro-

The high potency of methimazole suggested that it might be less toxic than any of the other antithyroid medications since the required dose would be very much less than for the other substances. Preliminary reports on the effects of the medication in hyperthyroidism looked promising.<sup>8,9</sup> A series of patients was therefore treated with methimazole and their response, course and toxic reactions evaluated.




FIG. 1. Structural formulas of thiouracil and methimazole. Dark portions represent the thiourea nucleus, which is probably responsible for antithyroid activity.

duction of 6-n-propyl-2-thiouracil and 4-methyl-2-thiouracil, the problem of toxicity has been greatly reduced. Both of the newer agents, however, have proved to be toxic in a small number of cases<sup>1-5</sup> and the search for less dangerous materials continues.

Stanley and Astwood<sup>6</sup> in 1947 screened thirty-two compounds for antithyroid activity and showed that the most potent material when tested in man was 2-mercaptoimidazole. Further studies of mercaptoimidazole derivatives showed that 1-methyl-2-mercaptoimidazole (methimazole, "tapazole") is even more active, being approximately 100 times as effective as thiouracil.<sup>7</sup>

The mercaptoimidazole group of goitrogens differs from the thiouracil group structurally in that the former possesses a five-member ring whereas the thiouracil ring has six atoms. The basic thiourea grouping, which is present in both molecules, is probably responsible for their activity. (Fig. 1.) The mode of action of the two types of goitrogens is therefore similar.

\* From the Department of Medicine, Emory University School of Medicine, and the Medical Service, Grady Memorial Hospital, Atlanta, Ga.

### MATERIAL AND METHODS

Thirty-five patients with proved hyperthyroidism were studied. (Table 1.) All of the patients at Grady Memorial Hospital in whom the diagnosis of hyperthyroidism was first made between July, 1950, and April, 1951, were included in the study. The diagnosis of hyperthyroidism was based upon the clinical picture, the serum protein-bound iodine concentration (PBI) and, in many cases, on the basal metabolic rate and I<sup>131</sup> uptake. The unreliability of the basal metabolic rate and the ready availability of the protein-bound iodine determination<sup>10</sup> led us to rely more on the latter; however, it was never accepted blindly without adequate clinical evidence of toxicity in the form of tachycardia at rest, heat intolerance, weight loss or changes of the hair and skin texture. In most cases all of these changes were observed and in almost all instances a definite goiter was also palpable.

In addition to this series a few patients were included from other hospitals. These patients were all seen personally by one of the authors, and the criteria for diagnosis were similar to those outlined for the Grady group.

None of the patients studied in this series had been receiving any other antithyroid medication before the administration of methimazole was begun. One patient had received propylthiou-

racil one year previously but had eloped from the clinic and had received no medication for a year when methimazole was started.

# RESULTS

*Effect on Hyperthyroidism.* Of the thirty-five patients treated with methimazole all but one

the patients gained weight (from 4 to 36 pounds, average 15.5 pounds) except for eight patients who had been in congestive failure when treatment was started. In this group evaluation of weight changes was complicated by the presence of edema, which invariably disappeared during therapy. The weight change in the patients with

TABLE I

CLINICAL DATA PERTAINING TO THIRTY-FIVE PATIENTS TREATED WITH METHIMAZOLE

1. Age.....	3 to 69 (avg. 39.6)		
2. Sex.....	12 males (34%)		
3. Race.....	21 colored (60%)		
4. Initial PBI (micrograms/100 ml.).....	9.3 to 28.7 (avg. 15.5)		
5. Initial basal metabolic rate.....	+28 to +96 (avg. +58); test available in only 25 patients		
6. Results of therapy.....	Euthyroid	33	
	Improved but not yet euthyroid	1	
	Dead	1	
7. Maximum dose of methimazole		Smooth goiters	Nodular goiters
	10 mg./day	1	
	20 mg./day	17	7
	30 mg./day	4	
	40 mg./day	3	2
	60 mg./day	1	
8. Duration of therapy to euthyroidism.....	Smooth goiters (26)	8.8 ± 1.1 wk.*	
	Nodular goiters (9)	8.2 ± 2.2 wk.	
9. Change during therapy (from initial observation to euthyroid state).....	Weight +4 to +36 (avg. +15.5)†		
	PBI -0.4 to -21.2 micrograms % (avg. 7.8 micrograms %)		
	Pulse -10 to -55 (avg. 30)†		
10. Final disposition of euthyroid group.....	Subtotal thyroidectomy	22	
	Radioactive iodine	9	
	Prolonged medical treatment	3	

\* Mean ± standard error of the mean.

† Eight patients with congestive failure were omitted from these figures.

showed dramatic improvement. (Table I.) The single exception was a fifty-eight year old man with severe congestive heart failure, who died of his cardiac disease after less than one week of therapy. During this time his serum protein-bound iodine had dropped from 16.6 to 11.7 micrograms/100 ml., indicating that he had begun to respond to the medication.

The remaining thirty-four patients were evaluated critically at one-week intervals for three or four weeks and thereafter at two-week intervals. An attempt was made to estimate the time required for control of hyperthyroidism by determining the approximate date at which the patient first became "euthyroid." Obviously such an estimate is subject to considerable error; but since the criteria for euthyroidism were standardized and the evaluations were performed by the same observers, it seems likely that they have a rough validity. It took from two to twenty-two weeks to control the toxic symptoms completely. During this time all of

heart disease varied from a loss of 15 pounds to a gain of 11 pounds. The pulse rate became less rapid in all patients. In those with normal cardiac function the heart rate was reduced from 10 to 55 beats per minute (average 30); when congestive failure was present, the decrease was similar but its significance is difficult to evaluate because many of these patients also had auricular fibrillation, and an accurate estimation of the mean pulse rate was difficult.

Methimazole caused a rapid decrease in the serum protein-bound iodine. There was a poor correlation between the fall of PBI and clinical improvement of the patient. The reduction of PBI often occurred before the patient improved clinically, an observation which has been noted previously.<sup>10</sup> The chief value of the PBI under these circumstances was to indicate that the circulating thyroid hormone concentration was kept within normal limits until the peripheral metabolism could fall to normal. By following the PBI it was possible to avoid the development

of myxedema, and clinical relapses could be predicted and prevented by observing a sudden rise of PBI before any change in the clinical status had occurred. During the course of therapy the PBI fell 0.4 to 21.2 micrograms/100 ml. (average fall was 7.8 micrograms/100 ml.).

*Dose.* The dosage of methimazole varied from 20 to 60 mg. per day, in divided doses of 5 mg. Table 1 shows how the doses were distributed. In general, a dose of 20 mg. per day was adequate; but if the PBI failed to show improvement within a week or two, it was usually increased to 30 or 40 mg. per day. In only one case were 60 mg. per day given—to a woman whose response to 40 mg. per day for eleven weeks had been inadequate. Since this patient was rather uncooperative, it is not certain that she had actually been taking the prescribed dose. On admission to the hospital the dose was raised to 60 mg. and at this level she showed rapid improvement and was ready for subtotal thyroidectomy within a week.

The patients with nodular goiter did not require any larger dose of methimazole or longer treatment than did the patients whose goiters were not nodular. (Table 1.) This is contrary to the experience with propylthiouracil, which has often proved to be relatively ineffective in patients with large or nodular goiters.

*Toxicity.* No toxic reactions were observed among the patients receiving methimazole. A specific search for leukopenia, agranulocytosis, fever and rash was made each time the patients were seen in the clinic. None of these complications were observed. In no case was it necessary to stop or reduce the dose because of toxic complications.

In addition to the main series of patients, methimazole was also administered to two patients in whom toxic manifestations had previously developed while propylthiouracil was being received. In these cases methimazole was well tolerated.

#### CASE REPORTS

**CASE I.\*** A sixty year old woman was admitted with classical symptoms and signs of thyrotoxicosis complicated by congestive heart failure and rapid auricular fibrillation. She had a large, smooth goiter. Her basal metabolic rate was +85 and her serum PBI was 14.9 micrograms/100 ml.

\* We are grateful to the staff of Piedmont Hospital, Atlanta, Ga., for permission to report this case.

Treatment was started with 300 mg. propylthiouracil daily in divided doses. After one week of therapy a fever of 103°F. developed which persisted for a week and disappeared within twenty-four hours after stopping the propylthiouracil. The propylthiouracil was resumed two days later and the dose increased to 400 mg. per day without return of the fever. After three weeks of therapy the PBI was 15.9 micrograms/100 ml. and the basal metabolic rate +75. The white count remained normal throughout. In the fourth week of treatment the patient was found to have a white count of 4,000 and propylthiouracil was promptly discontinued. The next day a severe sore throat developed, with ulceration of tonsils and palate. The temperature rose to 100.4°F. and the white count was found to be 3,650, with 2 per cent mature segmenters, 4 per cent myelocytes and 4 per cent premyelocytes. The white count returned to normal after three weeks of supportive and antibiotic therapy but thyrotoxicosis became worse. At this time treatment with 5 mg. methimazole four times daily was started. Within five days the PBI had dropped to 10.7 micrograms/100 ml. One month later the hyperthyroidism was greatly improved, no granulocytopenia had developed and the patient was being prepared for thyroidectomy.

*Comment.* In this patient large doses of propylthiouracil failed to control the hyperthyroidism but did produce serious agranulocytosis. Methimazole produced prompt improvement of the thyrotoxicosis and failed to cause toxic symptoms.

**CASE II.\*** This fifty-six year old farmer was admitted to the hospital because of a hemolytic streptococcus bacteremia which responded well to antibiotic therapy. On recovery it was apparent that he also had typical signs of hyperthyroidism. The serum PBI was 16.4 micrograms/100 ml. and the BMR +68. Treatment was begun with 375 mg. propylthiouracil per day, and the dose was increased to 500 mg. per day nineteen days later. On the twenty-fourth day of therapy he began to spike daily fevers of 103° to 104°F. Blood cultures were sterile and penicillin failed to control the fever. The propylthiouracil was therefore stopped and the fever subsided within a few days. A second trial of propylthiouracil therapy was begun after the temperature had been normal for three days. After ten days of treatment the spiking fever

\* The staff of Lawson Veterans Hospital kindly supplied the clinical data.



returned and continued for a week until propylthiouracil was discontinued. Five days later 20 mg. of methimazole per day were started and continued without difficulty for a month, at which time his PBI was 6.4 and he appeared

< .01).<sup>12</sup> The dose of propylthiouracil ranged from 300 to 700 mg. per day—more than ten times the amount of methimazole required. There were two instances of leukopenia in the fifty cases, an incidence of 4 per cent of poten-

TABLE II

CLINICAL DATA PERTAINING TO FIFTY PATIENTS TREATED WITH PROPYLTHIOURACIL IN 1948-1950

1. Age.....	15 to 77 (avg. 40.3)	
2. Sex.....	13 males (26%)	
3. Race.....	34 colored (68%)	
4. Maximum dosage.....	Less than 300 mg./day	1
	300 to 390 mg./day	16
	400 to 490 mg./day	20
	500 to 590 mg./day	5
	600 mg. or more/day	7
5. Duration of therapy until euthyroid state.....	6 to 104 wk. (avg. $30 \pm 3.6^*$ wk.)	
6. Results of therapy.....	Euthyroid	46
	Stopped therapy because of agranulocytosis	2
	Therapy incomplete	2

\* Standard error of the mean.

euthyroid. He was then given a therapeutic dose of 10 mc./I<sup>131</sup>.

*Comment.* Drug fever developed twice when the patient was given propylthiouracil. He was able to take methimazole without difficulty and had an uneventful recovery.

*Potency Compared with Propylthiouracil.* In view of the excellent effectiveness and apparent lack of toxicity of methimazole it seemed desirable to compare its potency with that of propylthiouracil. With this in mind we have studied a group of fifty patients previously treated with propylthiouracil during the two years prior to the present series. These patients were selected at random from the files of the Grady Memorial Hospital endocrine clinic and represent a fair cross section of the clinic population suffering from hyperthyroidism observed from January, 1948, to June, 1950. (Table II.) The propylthiouracil-treated group was examined to eliminate patients who received inadequate dosage of the drug (less than 300 mg. per day was considered inadequate)<sup>11</sup> or who had suffered complications which might have interfered with a smooth convalescence under the influence of the goitrogen. Four cases were dropped; one because of inadequate dosage, one because of pregnancy and two because of agranulocytosis.

Forty-four of the remaining forty-six patients became euthyroid after a course of treatment lasting six to 104 weeks (average thirty; standard error 3.6). When this is compared with the duration of treatment for the methimazole-treated group (average  $8.5 \pm 0.9$  weeks) the difference proves to be highly significant ( $p$

tially serious toxicity, as compared with the complete absence of toxicity in the methimazole group.

*Ultimate Disposition of Treated Patients.* Hyperthyroid patients who have been brought to the euthyroid state can be carried on maintenance doses of antithyroid medication for prolonged periods in the hope that a permanent remission will occur; or they can be treated in a definitive manner by either subtotal thyroidectomy or irradiation with I<sup>131</sup>. It is the practice of this clinic to give almost all hyperthyroid patients some form of definitive therapy because the nature of the clinic population makes it inadvisable to embark on a course of therapy which may continue over a period of many months. Of the patients in this series, therefore, twenty-two have been subjected to subtotal thyroidectomy and nine have received radioactive iodine. Final disposition has not been determined for the remaining four patients.

*Effect on Morphology of Thyroids Removed at Operation.\** Eighteen thyroid glands were available for pathologic study. They varied in weight from 18 to 160 gm., the average weight being 53 gm. Seven glands displayed a distinctly nodular architecture on gross examination.

A minimum of five tissue blocks were prepared from each gland. Tissues were fixed in Zenker's fluid with 5 per cent glacial acetic acid and were stained with phloxine and hematoxylin.

The glands of fourteen patients showed histologic changes of active hyperplasia. In most

\* This section was prepared by Dr. Abner Golden, Pathologist of Emory University Hospital, who reviewed all of the histologic slides available on these patients.

areas the follicular epithelium had a tall columnar appearance; it occasionally projected into the follicular spaces as small papillary infoldings. The nuclei were prominent and slightly pleomorphic, and mitotic figures were not infrequently encountered. Some nuclear fragmentation, cell degeneration and desquamation were apparent but these changes were unaccompanied by an inflammatory reaction. The follicles were generally decreased in size and contained pale staining, vacuolated or "scallop" colloid. A distinct increase in stromal vascularity frequently occurred.

The glands of four patients revealed marked involution, indicated by large colloid-filled follicles lined by flat or low cuboidal epithelial cells. Three of these patients had received pre-operative iodine therapy. None of the glands showed evidence of neoplastic change.

The histologic findings of active glandular hyperplasia are entirely comparable to those described following the administration of thiouracil to patients in the euthyroid or hyperthyroid states.<sup>13,14</sup>

*Effect of  $I^{131}$  Uptake.* The patients who received radioactive iodine received no methimazole for three days before the administration of  $I^{131}$ , in accordance with the recommendations of Williams et al.<sup>15</sup> Estimates of the percentage uptake of radioiodine were made in each case by direct counting over the neck, using a Geiger-Müller tube at 1 meter. The radioactivity over the neck was compared with the radioactivity of the therapeutic dose, counted before administration and extrapolated to twenty-four hours later on the  $I^{131}$  decay curve.\* Measurements were made twenty-four hours after the administration of the therapeutic dose, which ranged from 4 to 8 mc. The uptake by methimazole-treated patients was  $56 \pm 6.9$  per cent (range 40 to 85 per cent), which is approximately the same as that seen in hyperthyroid patients before treatment ( $>31$ ; average 54) and is definitely higher than the normal value (range 7 to 35; average 21). It is thus apparent that preparation of the patient with methimazole prior to irradiation of the thyroid gland with radioactive iodine does not interfere with the uptake of the isotope, nor with successful irradiation of the gland.

\* The data pertaining to  $I^{131}$  uptakes were generously supplied by Dr. Charles M. Huguley and Mr. William Miller of the Winship Clinic and Surgical Research Laboratory, Emory University, who also were responsible for the treatment of these patients with  $I^{131}$ .

#### COMMENTS

When a new drug is introduced for the treatment of hyperthyroidism, it must satisfy several criteria to justify acceptance. It should be less toxic than the medications already in use; it must be at least as effective; and it should not interfere with further therapeutic maneuvers such as subtotal thyroidectomy or irradiation with  $I^{131}$ .

In the group of patients treated with methimazole no instance of toxicity has been observed; in two cases methimazole proved to be non-toxic in patients who had had reactions to propylthiouracil. The number of patients studied, however, is too small to justify a categorical statement that methimazole is non-toxic; indeed, it seems almost certain that when a sufficiently large number of patients have been treated with the drug, toxic manifestations will appear. On the basis of our experience, however, it seems unlikely that the incidence of toxic manifestations will be as great as that found with propylthiouracil. One factor of importance in the low incidence of toxicity is probably the relatively small quantity of methimazole required to produce a therapeutic response. Not only is the daily dose about one tenth as large as the minimum effective daily dose of other antithyroid compounds but also the response is so rapid that the period of treatment required to attain the euthyroid state can be shortened substantially. As a result, the patient is exposed to a smaller amount of medication for a shorter period of time than when propylthiouracil is used.

The administration of methimazole is followed by a rapid improvement in the signs and symptoms of hyperthyroidism. We have attempted to compare the rate of response to methimazole with the rate of response to propylthiouracil. Since the methimazole-treated group was not studied simultaneously with the propylthiouracil-treated patients, the two groups cannot be considered completely comparable. Supervision of the methimazole group may have been somewhat closer than of the propylthiouracil patients, and other unrecognizable differences between the groups may have biased the results somewhat. In spite of these reservations as to the validity of the comparison it still seems probable that the response to methimazole was more rapid than to propylthiouracil. Since the drugs act in a similar fashion, it is difficult to explain this difference. The factor limiting the rate of response of a thyrotoxic patient to an



agent blocking the formation of thyroxine should be the rate of destruction of thyroxine by the tissues and the rate of dissipation of the effects of the hormone. There is no evidence to suggest that either methimazole or propylthiouracil affects the peripheral metabolism of thyroxine. The simplest explanation of the discrepancy, therefore, is that methimazole blocked the formation of thyroxine more nearly completely than did propylthiouracil. The difference may merely be one of dosage; but if this is the case, the effectiveness of methimazole must be even more than ten times as great as propylthiouracil.

It is a matter of considerable importance to be certain that an antithyroid drug does not complicate the task of the surgeon. For the most part, the patients operated upon in the present study received iodine for two or three weeks prior to operation. With this premedication the surgeons found the thyroids to be relatively easy to handle. Excessive vascularity and friability of the gland did not occur. In two cases operation was carried out without prior iodine preparation. In these cases the glands were somewhat friable and vascular, which discouraged the surgical staff from further attempts to operate without iodine medication.

The use of goitrogens in preparation for the administration of  $I^{131}$  may not be necessary in every case. However, it is known that irradiation causes a temporary increase in the circulating thyroxine<sup>16</sup> and that exacerbations of hyperthyroidism have occasionally occurred following irradiation. This may make it desirable to bring the patient to an euthyroid state before irradiating the gland, particularly when the patients in question have heart disease or are otherwise debilitated, as was the case in most of the patients who received  $I^{131}$  in this series. It is therefore important to find that preparation of patients for radioiodine therapy with methimazole does not interfere with the effectiveness of this type of therapy.

#### SUMMARY

1-methyl-2-mercaptoimidazole (methimazole, "tapazole")\* has been used in the treatment of thirty-five patients with hyperthyroidism. A good therapeutic response occurred in all cases. No toxic effects developed. Most patients responded to a dose of 5 mg. four times daily; occasionally as much as 40 mg. per day were given. The rate of response was significantly

\* Kindly supplied by Dr. D. C. Hines, Eli Lilly & Co., Indianapolis 6, Ind.

faster than for a group of fifty patients treated with propylthiouracil. Methimazole in combination with iodine provided satisfactory preparation for subtotal thyroidectomy. The drug did not interfere with the uptake of radioactive iodine in therapeutic amounts.

Methimazole ("tapazole") appears to be the antithyroid drug of choice at the present time.

#### REFERENCES

1. EISENMENGER, W. J. and STEELE, J. M. Leukopenia following the use of propylthiouracil. *J. A. M. A.* 135: 510, 1947.
2. MCGAVACK, T. H., GERL, A. J., VOGEL, M. and SCHUTZER, S. A clinical comparison of the effectiveness of 6-n-propylthiouracil and 2-thiouracil as antithyrotic agents. *Am. J. Med.*, 2: 144, 1947.
3. CRADDOCK, C. G., JR., SHOTTON, D., CROCKETT, C. L. and LEAVELL, B. S. Severe hypoprothrombinemia following propylthiouracil treatment of thyrotoxicosis. *New England J. Med.*, 244: 549, 1951.
4. MCCORMICK, R. V. Periarthritis occurring during propylthiouracil therapy. *J. A. M. A.*, 144: 1453, 1950.
5. BARFRED, A. Methylthiouracil in the treatment of thyrotoxicosis. *Am. J. M. Sc.*, 214: 349, 1947.
6. STANLEY, M. M. and ASTWOOD, E. B. Determination of the relative activities of antithyroid compounds in man using radioactive iodine. *Endocrinology*, 41: 66, 1947.
7. STANLEY, M. M. and ASTWOOD, E. B. 1-methyl-2-mercaptoimidazole: an antithyroid compound highly active in man. *Endocrinology*, 44: 588, 1949.
8. REVENO, W. S. and ROSENBAUM, H. Treatment of hyperthyroidism with 1-methyl-2-mercaptoimidazole. *J. A. M. A.*, 143: 1407, 1950.
9. BEIERWALTES, W. H. 1-methyl-2-mercaptoimidazole and 1-ethyl-2-mercaptoimidazole: a clinical evaluation of two new antithyroid drugs. *J. Lab. & Clin. Med.*, 36: 861, 1950.
10. HALLMAN, B. L., BONDY, P. K. and HAGEWOOD, M. A. Determination of serum protein-bound iodine as a routine clinical procedure. *Arch. Int. Med.*, 87: 817, 1951.
11. CODINGTON, A. B. and BONDY, P. K. Inadequacies of propylthiouracil in the treatment of thyrotoxicosis. *Am. J. Med.*, 5: 318, 1948.
12. SNEDECOR, G. W. *Statistical Methods*. 4th ed. Ames, Iowa, 1948. Iowa State College Press.
13. DONIACH, I. and SHARPEY-SCHAFER, E. P. Effect of 2-thiouracil on the histology of the normal human thyroid. *J. Endocrinol.*, 5: 131, 1947.
14. HALPERT, B., CAVANAUGH, J. W. and KELTZ, B. F. Structural changes in the thyroid glands of patients treated with thiouracil. *Arch. Path.*, 41: 155, 1946.
15. WILLIAMS, R. H., TOWERY, B. T., JAFFE, H., ROGERS, W. F., JR., and TAGNON, R. Radioiodotherapy. *Am. J. Med.*, 7: 702, 1949.
16. WERNER, S. C., QUIMBY, E. H. and SCHMIDT, C. Radioactive iodine,  $I^{131}$ , in the treatment of hyperthyroidism. *Am. J. Med.*, 7: 731, 1949.



# Seminars on Arteriosclerosis

## Arteriosclerosis—Some Clinical Implications\*

WILLIAM DOCK, M.D.

*Brooklyn, New York*

THE great upsurge of interest in and hope for the control of atherosclerosis should not blind the clinician to the fact that this is only one facet of the problem of arteriosclerosis, and that arteriosclerosis is only one factor in the loss of function encountered in an aging population. Since arteriosclerosis of various types has predilection for various sites, the emphasis in this discussion will be on local manifestations and on the interplay between arteriosclerosis and involutional changes of other sorts in leading to symptoms, signs and dysfunction.

### BRAIN

When we observe cerebrovascular accidents occurring in subacute endocarditis and in auricular fibrillation in people under thirty years of age, we are impressed by the lack of change in personality, in spite of a chronic illness, progressive crippling and repeated insults to the brain. On the other hand, in people over fifty and especially in those over seventy, we are struck by the loss of interest, energy and judgment which so often follows small cerebrovascular accidents. Alvarez,<sup>1</sup> in his contributions to the problem of "little strokes" with great deterioration in personality and with outstanding digestive or "neurotic" complaints, has cited many of these instances in which repeated strokes, with negligible neurologic disturbance, caused bizarre and misdiagnosed illnesses. At autopsy the elderly subjects have disseminated cortical atrophy and dilated cerebral ventricles as a result of involutional changes, as well as focal softening or hemorrhage, while the young ones have only focal lesions. We conclude that when the brain is young and its neurone content high, cerebral accidents produce much less change in personality than when the brain is already atrophic from involutional loss of neurones.

Avitaminosis, combined system disease and

hypothyroidism are other factors which accentuate the functional effects of cerebral anoxic injury. In combined system disease and in cortical atrophy the Schmidt-Kety method reveals practically normal blood flow, with low O<sub>2</sub> and glucose uptake.<sup>2</sup> Those factors which are reversible with therapy, or at least held in check while the patient adjusts himself to his disability, must always be in the clinician's mind, along with the environmental stresses peculiar to aging, namely, retirement, loss of relatives and friends, declining buying power of fixed incomes, conflicts with children and aggressive young associates. Evaluation of the role of cerebral arteriosclerosis and of "little strokes" in the development of senile dementia and psychoneuroses is difficult even with a good history and a complete autopsy. During life we must not ascribe to cerebral vascular disease all the manifestations of cerebral aging, which may in part be reversible with B<sub>12</sub> or other water-soluble vitamins, or greatly improved by correcting the social environment and the attitude of relatives and associates.

The cerebral arteries are subject to several types of sclerosis. The simplest is the intimal proliferation which occurs with syphilitic mesarteritis or the periarteritis of diffuse vascular disease or of tuberculous meningitis. Any of these may predispose vessels in the meninges to thrombosis or to occlusion hastened by atheroma developing secondarily. Similar internal thickening and secondary atherosclerosis occurs in aneurysmal weakening of basal arteries on a congenital basis. In the small arteries in the brain substance calcific or cystic medial disease and endarterial proliferation follow cortical atrophy and predispose the vessels to minute atheromas and to rupture. Thus the cerebral involution which accentuates functional loss from small strokes also plays a part in the pathogenesis of the vascular disease which causes such

\* From the Department of Medicine, State University of New York, State University Medical Center at New York, Brooklyn, N. Y.

accidents. Familial tendency to cerebral arteriosclerosis, rather than coronary disease, may be due in part to the pattern of cortical aging and in part to body build. Wilens has correlated low intracranial pressure, due to the upright posture and a relatively long spinal canal, with the relatively high ratio of cerebral to coronary accidents.<sup>3</sup>

#### RETINA

While the branches of the ophthalmic artery are not free from disease due to syphilis, tubercles or other inflammatory lesions, they suffer chiefly from "degenerative" lesions associated with hypertension, nephritis and diabetes. Cystic degeneration of the retina may predispose to vascular disease in the same way as cortical atrophy, but for the most part aging normotensives show only tortuosity of the arteries, without focal vascular lesions. It is a curious fact that in uremia and hypertension associated with chronic pyelonephritis or congenital cystic disease of the kidneys retinal sclerosis is somewhat less marked than in uremia due to glomerulonephritis. In some middle-aged uremics with these diseases the retina may be practically normal as it is in obstructive uropathies. When hypertension or diabetes precedes proteinuria by several years, retinal vascular disease is nearly always severe when urea retention is marked. The retina, like the kidney, has a high rate of blood flow per gram of functioning tissue, and in diabetes both organs show lesions far out in the arteriolar system, or even in the capillaries and juxtacapillary venules. In toxemia of pregnancy and malignant sclerosis the arterial component in the retina is complicated by papilledema and venous engorgement due to high intracranial pressure.

#### KIDNEYS

Three types of arteriosclerosis are manifested in the renal arterial bed, namely, reduplication of the medial elastica, atherosclerosis and regenerative intimal thickening due to atrophy or to adjacent foci of inflammation. In chronic pyelonephritis, glomerulonephritis, periarteritis and disseminated lupus all three processes are at work simultaneously in most of our patients, and we must go to the Orient or Central Africa to study these conditions uncomplicated by atherosclerosis. The speed with which renal failure develops after the acute inflammatory process is spent depends to a large degree on the

rate at which atherosclerosis develops. This in turn depends on the degree of hypertension and alteration in plasma lipids. In the kidney, as in the retina, occlusive lesions of the main arterial trunks are rare, and disseminated sclerotic lesions, combining two or more of the types mentioned, affect the arterioles all the way into the glomerulus and, in diabetics with Kimmelstiel-Wilson syndrome, even into the glomerular capillaries.

In pyelonephritis the severity of disease may vary greatly from one lobule to another, but in other forms of Bright's disease a single needle biopsy of the cortex gives an excellent sample of the whole cortex.<sup>4</sup> Obtaining such samples involves less discomfort or risk than biopsy of the liver, so that the kidney can now be studied histologically and functionally more accurately and safely than any other organ save the skin. Our knowledge of the evolution of renal vascular disease and of its control by therapy is therefore certain to be more precise than that of other organs. Since its atheromatous lesions closely parallel those of the retina, careful study of either organ will enrich our understanding of disease in the other. This will be particularly valuable in following the effects of lowering arterial hypertension, since the role of elevated pressure in initiating or aggravating atherosclerosis is still disputed.

Because parenchymal atrophy and failing efficiency of enzyme systems occur in the renal tubules with aging even when vascular disease is minimal, the fixation of specific gravity of urine and decline in maximal rate of para-amino hippurate excretion with old age cannot be taken as accurate measures of functional impairment due to arteriosclerosis. Nor does the most precise study of urinary sediment provide decisive information on the relative importance of arteriosclerosis and of continuing glomerulitis in cases of chronic glomerulonephritis with hypertension. While the clinical management of these cases does not depend on such evaluations, there can be no doubt that understanding of the course of the disease in each individual will be enlarged by careful study of serial cortical biopsies.

#### HEART

The coronary system presents problems the reverse of those encountered in the previous sections. Arteriolar sclerosis is practically nonexistent even in malignant phases of hyperten-

sion. Inflammatory disease is a rare cause of internal thickening or thrombosis. Syphilis rarely extends beyond the ostia. Coronary atherosclerosis of the large branches lying in the epicardium is almost the only type of "arteriosclerotic heart disease," a term which should be abandoned and replaced by coronary sclerosis, coronary insufficiency or myocardial infarction. Even in the epicardial arteries the smaller branches are much less severely diseased than the main trunks.

Although coronary atherosclerosis and insufficiency are all too frequent, they have been blamed for many conditions for which they are not responsible. As in the brain, functional failure of the heart is often ascribed to vascular disease or to the myocardium outgrowing the coronary supply, when the real fault is an involutional or metabolic defect. The coronary blood flow studies of Bing<sup>5,6</sup> and Goodale<sup>5</sup> have confirmed the postmortem perfusion studies of Dock<sup>7</sup> and of Dresdale.<sup>8</sup> In cardiac hypertrophy and even in heart failure, blood flow per 100 gm. of ventricle per minute remains within the normal range; and if chronic anemia is present, the coronary flow, even in large failing hearts, may be far greater than normal, as is the perfusibility postmortem.<sup>8</sup> Insufficiency of coronary flow can be due only to coronary atherosclerosis, arteritis or embolism, or to low perfusion pressure. The latter occurs in the congenital displacement of coronary ostia into the pulmonary artery and in diastolic hypotension. When this is chronic, as in aortic insufficiency, coronary hyperplasia may provide normal coronary flow even for a greatly hypertrophied heart.<sup>5</sup>

When myocardial infarction strikes elderly subjects, congestive heart failure not infrequently occurs acutely or within a few months. Experience with infarction in young soldiers proved that heart failure was extremely rare, acutely or after recovery from large infarcts. In the heart, as in the brain and kidney, involutional changes prepare the way for organic failure with vascular accidents which, in young people, would be the cause of transient and trivial dysfunctions. Since the heart, like the brain, is subject to metabolic injury from poor diet, and this happens more often in poor old people than in others, what is called "arteriosclerotic disease" not infrequently proves to be greatly benefited by a diet adequate in protein and water-soluble vitamin, especially thiamine. "Ar-

teriosclerotic," that is, coronary heart disease, can be evaluated only when anemia, Graves' disease and malnutrition have been corrected. Involutional change in myocardial function seems to be particularly susceptible to correction by digitalis which does not improve coronary flow, but an exact estimation of this element in heart failure can scarcely be made even after autopsy and the most meticulous clinical study.

Since atherosclerosis of the main coronary arteries may be advanced in normotensive young men who have no other visible atheromas, local factors, and notably the great thickness of the intima here as contrasted with the myocardial coronary twigs and other arteries of similar size, seem to be paramount. Without this, the metabolic disturbance shown by the plasma lipid pattern might only cause asymptomatic aortic lesions years or decades later. Family differences and even sex differences in the size, number and distribution of coronary intimal cushions at birth, and in the diffuse intimal thickening limited to the epicardial branches even in senescence, may explain some of the features of this form of atherosclerosis. Today this "middle-aged tax-payer's friend" has replaced pneumonia, the "old man's friend" in Sir William Osler's day, as Captain of the Men of Death.

The electrocardiograph, and more recently the ballistocardiograph, have been of great value in detecting and evaluating coronary disease after it has caused mild symptoms or even before symptoms have occurred. Figure 1 is based on data collected by Drs. H. and R. Mandelbaum on people free of symptoms and signs (including electrocardiographic changes) of cardiovascular disease. It shows how the incidence of ballistocardiographic abnormality in such subjects parallels the incidence of 3+ coronary sclerosis in a large series of miscellaneous autopsies. Starr<sup>9</sup> had already reported the high incidence of later coronary accidents among normal subjects with abnormal ballistic patterns. Such a method may help in selecting patients who might benefit from studies of plasma lipids and attempts at modifying abnormal patterns of lipid content of the blood.

#### LEGS

Vascular disease of the legs is far more striking in the veins than in the arteries but the latter show far more sclerosis than do the corresponding arteries in the arms and hands. The legs are colder, the arteries dilate less when metabolism



rises, and blushing stops above the waistline. While these facts may be related to crural susceptibility to both atherosclerosis and medial calcification, it seems probable that the greatest factor is a difference in arterial pressure. The pulse pressure and systolic pressure rise as the

shows that involution of joints and the nervous system has greatly increased the effort of walking, intermittent limp, like angina of effort, is due largely to vascular disease not involution. While angina is usually associated with pure atherosclerosis, intermittent limp is often a




FIG. 1. The crosses show the incidence of 3+ coronary disease disclosed at autopsy at the Mayo Clinic in a miscellaneous group of patients.<sup>11</sup> The black areas show percentage of maximal, the cross-hatched areas the percentage of marked, and the stippled areas the percentage of definite abnormality in ballistocardiograms of supposedly normal subjects (H. and R. Mandelbaum). The inflection on all curves in males at fifty-five years presumably is due to deaths from coronary disease exceeding rate of development of atheromas. The curves of severe and marked abnormality parallel the curves of incidence of coronary sclerosis although these are very different in men and women. The relatively great rise in minimal abnormality in older men and women probably is due to poor physical training and obesity.

pulse wave goes down the aorta. Even in recumbency, pressure is higher in the legs than in the arms. Erect posture adds many centimeters of hydrostatic pressure (although never as much as the elevation of the heart above the vessel). Except in diabetics and a small group of heavy smokers, arteriosclerosis of the legs rarely causes symptoms before the age of fifty, but calcification in the femoral and tibial arteries may be demonstrated by x-ray for decades without symptoms. Presumably this is due to Mönckeberg's type of medial calcification which does not encroach on the lumen. Atherosclerosis involves both large and minute arteries in the legs, but arteriolar disease is limited to inflamed regions. These occur in a small percentage of heavy smokers, who have thromboangiitis obliterans, and also secondarily to varices or to chronic edema in heart disease and cirrhosis.

Although the hobbling gait of many old people

result of mixed vascular lesions. An effective method of controlling atherosclerosis probably would greatly reduce the vascular disease of the legs of diabetics but would still leave a group of patients with limp or gangrene due to angiitis and other vascular disorders.

#### AORTIC, ILIAC AND SPLANCHNIC ARTERIES

In the aortic, splenic, carotid and temporal arteries we can observe the most frequent form of arteriosclerosis, a form closely related to the venous involution which causes varices. This is called fibrosis with ectasia. Elastic and muscular elements atrophy, collagen increases and the loss of distensibility is compensated by dilatation and increase in length. The resulting tortuosity and the dilatation are evident on the temples of many men before they are thirty, and in the chest films of some middle-aged or aged people the aortic tortuosity and dilatation mimics that

due to syphilis. A striking difference is that secondary atherosclerosis and calcification, which often occurs in the ascending part of the aorta with syphilis, is rarely seen proximal to the aortic knob, on the descending limb of the arch, in the other group. In women, and in men with coarctation of the aorta, tortuosity of the common carotid or innominate arteries may also cause pseudo-aneurysms. In women, too, extreme tortuosity of the abdominal aorta is more frequent than in men. There is little evidence that these involutional changes greatly alter the rate of atherosclerosis or cause other vascular lesions or organic disease.

As men age, at least in North America, there is an increasing calcium content of the media, especially in the abdominal part of the aorta, but true medial sclerosis rarely involves the aorta, carotids or large splanchnic arteries. Cystic degeneration and necrosis, the precursor of sudden medial separation (dissecting aneurysm), also is a rare disorder, having a significant occurrence only in severe hypothyroidism combined with severe hypertension.

Atherosclerosis affects the arterioles of the adrenal glands, pancreas and, to a lesser degree, those of the gastrointestinal tract and liver only in severe hypertensive and diabetic vascular disease. Splenic arterioles also are affected and show intimal hyaline in many elderly people. But in the splanchnic bed the large arteries rarely show atherosclerosis comparable to that of the coronaries. This is nearly always associated with severe atherosclerotic ulceration and calcification of the aorta. In elderly patients, usually women, chronic post-prandial distress (abdominal angina) may be due to severe sclerosis of branches of the celiac axis. Occasionally fatal retroperitoneal hemorrhage occurs as a result of ulceration of abdominal aortic or iliac atheromas. Similar hemorrhages may also occur from rupture of aneurysms of renal arteries, especially on the right side. The differential diagnosis is important since surgical treatment is more likely to succeed with the renal arterial lesion. The use of roentgen-opaque media, injected by a needle inserted into the aorta through the left flank, may serve to demonstrate such aneurysms of the renal artery. In retroperitoneal hemorrhage from either type of lesion, pain and paralytic ileus are prominent features, and discoloration of the scrotum or vulva and abdominal wall may develop within a day.

#### GENERAL CONSIDERATION ON PATHOGENESIS AND CONTROL

It is now generally recognized that arteriosclerosis rarely causes trouble in normotensives before the age of fifty unless the heart or the legs are involved, and in such cases a disturbed cholesterol-phospholipid balance is usually demonstrable. As patients age the ratio of lesions in the brain and in the aorta rises and the cholesterol-phospholipid levels approximate those of people who have minimal sclerosis. Diet and defects of metabolism appear to be of much less significance in atherosclerosis manifest after sixty-five years of age than in that evident before fifty.

Excessive smoking has been noted in a higher percentage of those suffering coronary accidents before fifty than in control populations, and the reduced life expectancy of smokers established by Pearl must in large part be related to the very striking actions of nicotine on the cardiovascular system.<sup>10</sup> Ballistocardiographic changes equivalent to those of coronary disease occur in a few normal young smokers for many minutes after a cigarette, and seem to reveal poor habituation to tobacco. Severe electrocardiographic changes occur much more rarely from smoking. It may be safely stated that smoking is a definite factor in some cases of arteriosclerosis of the legs, probably contributes to coronary disease, and is almost a negligible factor in retinal, renal and cerebral arteriosclerosis. There is no evidence whatever that alcohol or any other agent abolishes the effect of smoking on the circulation.<sup>10</sup>

#### ARTERIOSCLEROSIS AND SMOKING

While there is a growing body of evidence that exogenous (dietary) cholesterol forms the bulk of the atheromatous deposits in cholesterol-fed rabbits,<sup>10</sup> and that dietary cholesterol as well as fat is a matter of some importance to individuals who develop coronary disease, there is far less evidence that smoking is an important factor in the rate of evolution of atheroma. This evidence has recently been summarized by Roth.<sup>11</sup>

The effects of tobacco are apparent as vasoconstriction in the extremities of all subjects, minor changes in the ballistocardiogram in one-third and in the electrocardiogram in one-half of normal subjects. Marked changes in the ballistocardiogram occur in over 5 per cent, in the electrocardiogram in less than 1 per cent. The cumulative effects of such changes probably contribute to the reduced life expectancy of

smokers reported by Pearl<sup>12</sup> but a far greater contribution to that mortality may be due to effects on the respiratory system. Not only cancer of the lung but chronic sinusitis, chronic bronchitis, emphysema and reduced resistance to pneumonic infection could be cited. Hence, while the physician can advise the inquiring patient with vascular disease of the leg that smoking will surely curtail blood flow in the diseased tissue, he cannot be dogmatic about the hazard to the patient with angina or coronary disease. Modern technics make possible detection of those whose cardiac function shows no disturbance and others who show striking deterioration. Objective proof of no effect is reassuring, and evidence of actual impairment of function due to half a cigarette may help a patient to give up this habit.

## SUMMARY

Atherosclerotic accidents to the coronary circulation begin at an age when myocardial function is sound and in subjects of this age rarely cause heart failure. At older ages, however, the net effect of coronary disease and of myocardial involution is a steady rise in the incidence of heart failure with coronary atherosclerosis.

Arteriosclerotic accidents in the brain are infrequent prior to the age at which disseminated cortical atrophy, an involutional loss of neurones, usually begins to reduce mental acuity and memory. In nearly all cases in which personality changes follow cerebrovascular accidents involutional change is an important factor in reducing ability to resist injury. In some elderly subjects this is further complicated by inadequate supply

of water soluble vitamins, including B<sub>12</sub>, so that a *a priori* assumption that all deterioration in mentation and emotional adjustment of oldsters is due to arteriosclerosis is not merely erroneous but harmful, since it eliminates from proper therapy many whose condition can be improved.

## REFERENCES

1. ALVAREZ, W. C. Small commonly unrecognized strokes. *Postgrad. Med.*, 4, 96, 1948.
2. SCHEINBERG, P. Cerebral metabolism in pernicious anemia. *J. Clin. Investigation*, 29: 843, 1950.
3. WILENS, S. L. Orthostatic influences in the distribution of atheromatous lesions in the cerebral and other arteries. *Arch. Int. Med.*, 82: 431, 1948.
4. IVERSEN, P. and BRUN, C. Aspiration biopsy of the kidney. *Am. J. Med.*, 11: 324, 1951.
5. BING, R. J., HAMMOND, M. M., HANDELSMAN, J. C., POWERS, S. R., SPENCER, F. C., ECKENHOFF, J. E., GOODALE, W. T., HAFKENSHEL, J. H. and KETY, S. S. The measurement of coronary blood flow, oxygen consumption and efficiency of the left ventricle in man. *Am. Heart J.*, 38: 1, 1949.
6. BING, R. J. and DALEY, R. Behavior of the myocardium in health and disease as studied by coronary sinus catheterization. *Am. J. Med.*, 10: 711, 1951.
7. DOCK, W. The capacity of the coronary bed in cardiac hypertrophy. *J. Exper. Med.*, 74: 177, 1941.
8. DRESDALE, D. T. Personal communication.
9. STARR, I. The ballistocardiograph. *Harvey Lect.*, 42: 194, 1947.
10. BIGGS, M. W. and KRITCHEVSKY, D. Observations with radioactive hydrogen in experimental atherosclerosis. *Circulation*, 4: 34, 1951.
11. ROTH, G. Tobacco and the Cardiovascular System. Springfield, Ill., 1951. Charles C Thomas.
12. PEARL, R. Tobacco smoking and longevity. *Science*, 87: 216-217, 1938.
13. WHITE, N. K., EDWARDS, J. E. and DRY, T. J. The relationship of the degree of coronary atherosclerosis with age, in men. *Circulation*, 1: 645, 1950.



# Diet and Lipotropic Agents in Arteriosclerosis\*

JACK D. DAVIDSON, M.D.†

*New York, New York*

THE use of low cholesterol diets and the administration of various lipotropic compounds are two types of therapy that have been advocated in recent years for the treatment of arteriosclerosis. This paper attempts to review the evidence for their value.

Both of these treatments are predicated upon the hypothesis that arteriosclerosis is due to a disorder of lipid metabolism which results in the deposition of lipids in the arterial wall as a principal factor in the formation of the lesion. The evidence for this hypothesis has been presented in prior articles in this symposium.<sup>1-3</sup> The low cholesterol diet is intended to cut down the supply of the most characteristic lipid of the lesion, while the use of lipotropic agents is directed at mobilization and removal of the lipids from the lesions. The two therapeutic measures can be evaluated on the basis of: (1) their theoretic mode of action, (2) their effects in experimental arteriosclerosis in animals and (3) a limited amount of data reported on their clinical application.

## EFFECT OF LOW CHOLESTEROL DIETS

*Theoretic Basis.* The low cholesterol diet off-hand seems a reasonable approach to prevention of cholesterol deposition in the arterial walls. Recent studies in cholesterol metabolism indicate, however, that a large amount of this substance is synthesized by the body from small molecules available from fat, protein or carbohydrate. Gould<sup>2</sup> has shown that dietary cholesterol depresses endogenous cholesterol synthesis in animals, a finding which may imply the converse, that if dietary intake is insufficient cholesterol synthesis will be increased. Although all three basic types of food can supply the

small molecules for cholesterol synthesis, fats seem to be most efficiently converted and this provides a rationale for restriction of all fats as well as cholesterol in the dietary treatment of arteriosclerosis.

*Animal Experiments.* There is not much direct evidence from animal experiments for or against the low fat, low cholesterol diet. Horlick et al.<sup>4</sup> found that low fat diets did inhibit the spontaneous arteriosclerosis which occurs in chickens. Most experimental arteriosclerosis is produced in chickens, rabbits and dogs by feeding high cholesterol diets. This precludes studies of low fat, low cholesterol diet but can itself be taken as indirect support for the value of low cholesterol diets. It is also true that upon return to normal diets these animals all show regression of their arteriosclerotic lesions.<sup>2,5,6</sup> Equally indirect is the finding that inadequate caloric intake has an inhibiting effect upon the production of this type of experimental arteriosclerosis, possibly mediated by depletion of the pool of precursors for cholesterol synthesis.<sup>7</sup>

*Clinical Studies.* There is more support for low fat, low cholesterol diets from clinical studies. Thus statistical surveys of various segments of the population indicate that there is a higher incidence of arteriosclerosis in those groups whose diet is inherently richer in fats and cholesterol than in groups whose diets are poor in fats and cholesterol.<sup>8-11</sup> These studies, however, are open to criticism because the groups compared differ in other significant but uncontrolled factors such as heredity, nutritional status and differing age distribution. Nutritional status was studied by Wilens<sup>8,12</sup> in a series of autopsies. He found increased arteriosclerosis in obese individuals and less than the average incidence in

\* From the Research Service, First (Columbia) Division, Goldwater Memorial Hospital, Department of Hospitals, New York City and the Department of Medicine, College of Physicians and Surgeons, Columbia University, New York. Aided by grants from the National Heart Institute, The Public Health Institute of the City of New York and the Albert and Mary Lasker Foundation.

† Present address: Francis Delafield Hospital, New York, N. Y.

people who were below average weight. Age has a recognized correlation with arteriosclerosis so that in groups in which life expectancy is low due to infectious diseases fewer individuals survive to the age at which arteriosclerosis becomes common.

More reliable data are available from studies of people experimentally placed on low cholesterol, low fat diets. Morrison<sup>13</sup> reported a direct influence of diet on arteriosclerosis in terms of diminished mortality in a group on a diet containing 25 gm. of fat as compared with a control group on a normal diet. Although his figures are statistically significant, the number of deaths involved is rather small and the work should be extended. There is also some question from the reports<sup>14</sup> as to whether these patients were not also receiving choline.

The ultimate in low cholesterol, low fat diets was used by Kempner<sup>15</sup> in his rice diet study. His diet contained no cholesterol and less than 5 per cent fat. Studying hypertensive cardiovascular disease principally, he found that in 363 patients with initial serum cholesterol levels over 220 mg. per cent there was an average drop of 74 mg. per cent after treatment with the rice diet; 148 patients with initial serum cholesterol below 220 mg. per cent showed an average decline of only 15 mg. per cent on the rice diet. Starke<sup>16</sup> subsequently presented data on 154 of Kempner's patients in greater detail, coming to the same conclusion. Watkin et al.<sup>17</sup> at this institution independently studied Kempner's rice diet and found that thirty-five of forty-one patients showed a lowering of serum cholesterol while on the rice diet, the average decline being 40 mg. per cent. Their data included the frequent cholesterol determinations necessary to average out random fluctuations. They also followed the weight of each patient, indicating that the observed fall in serum cholesterol was attributable to the qualitative nature of the diet and not to starvation. Mellinkoff, Machella and Reinhold<sup>18</sup> noted a fall in serum cholesterol averaging 85 mg. per cent in fourteen patients with intestinal disease when treated with a protein hydrolysate and dextri-maltose preparation that was low in fat. This fall they attributed to the low fat therapy but this is somewhat questionable due to the patients' underlying diseases. Hildreth, Malinkoff, Blair and Hildreth<sup>19</sup> have recently reported a closely controlled study of fat restriction in three healthy subjects. Reduction of the dietary fat from their normal levels of 97, 114

and 138 gm. to 9, 10 and 62 gm., respectively, in these three subjects caused serum cholesterol declines of about 45 mg. per cent in each case. Restoration of pure vegetable oil to the diets produced significant rises in the serum cholesterol levels, proving the necessity for fat as well as cholesterol restriction. Other studies of diet in relation to serum cholesterol level were made by Wilkinson, Blecha and Reimer<sup>20</sup> and by Gertler, Garn and White<sup>21</sup> in which a higher minimum level of fat and cholesterol intake existed and little or no apparent effect upon the serum cholesterol level was noted.

It, therefore, seems likely, as Keyes et al.<sup>22</sup> have pointed out, that the blood cholesterol level is independent of intake over a wide range of cholesterol in the diet but that as the intake approaches "zero," which they believe to be probably less than 200 mg. per day, the blood level does decline. It also seems established that to achieve significant lowering of serum cholesterol the diet must not only be extremely low in cholesterol but must also be restricted markedly in all fats. The addition of vegetable fat to such diets low in cholesterol results in a rise in the serum cholesterol level toward the pre-diet level. With really low cholesterol, low fat diets the serum cholesterol can be reduced by 40 to 80 mg. per cent in most patients and particularly in patients who have hypercholesterolemia. The ultimate value of this degree of serum cholesterol reduction in the therapy or prevention of arteriosclerosis remains to be determined.

In addition to studies on the effects of diet on arteriosclerosis mortality and serum cholesterol level, there is the work of Gofman et al.<sup>23</sup> on the quite striking influence of a low cholesterol, low fat diet upon the level of the  $S_{12-20}$  lipoprotein fraction of serum. In a previous article in this series<sup>3</sup> Jones et al. described the correlation between the level of  $S_{12-20}$  lipoprotein of human serum and arteriosclerosis. They found that a diet of less than 50 gm. of fat and 200 mg. cholesterol evoked a fall in the  $S_{12-20}$  lipoprotein concentration in the serum of most patients. They reported that in fifty-six patients with  $S_{12-20}$  concentrations above 80 mg. per cent, who were allegedly on this diet for a year after a myocardial infarction, there were no recurrent infarctions among the fourteen patients who showed a fall in their  $S_{12-20}$  lipoproteins whereas sixteen recurrent infarctions occurred in the forty-two patients whose  $S_{12-20}$

level remained above 80 mg. per cent. Unfortunately this study does not indicate the effect of the diet upon recurrent myocardial infarction directly since the authors used the trend of the  $S_t$  12-20 level as their index of degree of adherence to the diet.

In contrast to Gofman's observations, a preliminary report of the  $S_t$  10-20 lipoproteins of patients on the rice diet by Hatch and Kendall<sup>24</sup> states that this drastic diet resulted in substantial increases in the  $S_t$  10-20 level in three patients, no change in three, and a decrease in only one. This is not discordant with Gofman's findings because the rice diet is much lower in fat than that studied by Gofman, and differs significantly in protein content as well. This rice diet effect is being studied further.

#### EFFECT OF LIPOTROPIC AGENTS

The case for or against the use of lipotropic agents in arteriosclerosis is much less clear and the available data are conflicting. Some of the conflicts are the result of inadequate knowledge at the time of early work and can now be resolved in the light of more recent findings. Some of the reports, however, seem to be based upon inadequate observations.

*Theoretic Basis.* The theoretic basis for the application of lipotropic agents to the management of arteriosclerosis is twofold, their lipotropic action and the fact that some are phospholipid components. Their use was suggested initially by the gross similarity of the fatty deposits in arteriosclerosis to those in fatty livers, which had just been reported responsive to substances now known as "lipotropic agents." This analogy is very superficial. The phrase "fatty liver" covers a number of different types of lipid infiltration of the liver arising from multiple causes. None of these lipid deposits are very similar chemically or histologically to the lesions of arteriosclerosis. This reasoning by analogy is even more faulty because the lipotropic agents are not uniformly effective against all fatty livers. Some types of fatty liver respond, others do not.

The lipotropic agents comprise five chemical compounds: choline, methionine, betaine, inositol and  $\beta$ -propiethetin.<sup>25</sup> The last can be dismissed in that it is thus far known to occur in seaweed, is unstudied in foods, and has not been applied to arteriosclerosis. The first three are interrelated in that methionine and betaine act by providing labile methyl groups for the syn-

thesis of choline.<sup>26</sup> Chemically, inositol stands in a class by itself, apparently unrelated to choline. A number of more complex substances have lipotropic activity by virtue of their choline content or their ready conversion to this compound. Thus lecithins, for example those derived from soy beans or egg yolk, yield choline upon hydrolysis. Proteins are lipotropic due to their methionine content, which in turn gives rise to choline. Lipocaic from pancreas is still too indeterminate for consideration.<sup>27</sup> This leaves just two compounds to be evaluated, choline and inositol.

Choline was the first compound shown to have lipotropic activity. This was accomplished by Best in 1932<sup>28</sup> when he recognized it as the active component in lecithin. It is effective in removing fat from livers made fatty by choline deficient diets or by high fat diets but not from fatty livers produced by carbon tetrachloride or phosphorus.<sup>29</sup> It is therefore not a universally effective lipotropic agent for fatty livers, to say nothing of arteries. Gavin and McHenry reported a curative action of inositol upon fatty livers arising from high fat diets.<sup>30</sup> There is still some controversy over its general effectiveness and whether it does anything that choline does not.<sup>31,32</sup> Initially, it was believed to have some specificity for the cholesterol type of fatty liver but Best<sup>25,32</sup> recently has evaluated it carefully and now believes that inositol does nothing that choline will not do except in very short (less than two weeks) experiments. It is not as strongly lipotropic in maximally effective doses, and is completely ineffective if fat is present to the extent of over 2 per cent in the diet. Most of the confusion in the past was apparently due to chance biologic variation when small numbers of animals were used, and to different effects dependent upon the duration of experiments and variation in the diets used by several investigators. In general, the specificity of action of lipotropic compounds for certain types of fatty livers and the conditions necessary for demonstration of their activity even in livers leave little theoretic basis for their use in the prevention or treatment of arteriosclerosis.

The second theoretic basis for the use of choline in arteriosclerosis came later when the probable importance of the phospholipids in the pathogenesis of the disease became apparent.<sup>2</sup> There is increasing evidence to indicate that the ratio of serum phospholipids to serum cholesterol may be more important than the



level of the serum cholesterol alone. If the serum phospholipids are involved in the prevention of arteriosclerosis, certainly choline and possibly inositol might act favorably by providing components of the phospholipids and thereby contributing to a desirable high level of these lipids. There is in this a reasonable rationale for testing these substances in the therapy or prophylaxis of arteriosclerosis, regardless of any consideration of lipotropic activity demonstrable in the liver.

*Animal Experiments.* Quite a large number of studies have been made of the effects of various lipotropic agents upon experimental arteriosclerosis in animals, since in such experiments it is possible to determine the degree of arteriosclerosis at autopsy as well as to study the effect on serum cholesterol levels. Rabbits, chickens and dogs have been used, the majority of published reports relating to rabbit experiments. With both rabbits and chickens there have been conflicting conclusions as to the effectiveness of the substances tested. The number of reports in favor of the lipotropic agents exceeds the unfavorable reports but this is a misleading index of results because of the reluctance and lack of interest of many investigators in publishing their negative experiments. Critical analysis of the published reports reveals that many of them must be rejected as inconclusive for various reasons.

Experimental arteriosclerosis, as produced in rabbits by cholesterol feeding, is subject to a large number of factors which influence the incidence and severity of the lesions. It is now recognized that even with all known variables controlled there is such marked biologic variation in experimental arteriosclerosis in rabbits that it is hazardous to use less than about twenty animals for each test regimen, with an equal number of control experiments limited to feeding the same quantity of cholesterol alone.<sup>33</sup> In setting up an experiment the control and test groups should be of identical strain, sex distribution, age and weight. They should be carried on experiment simultaneously for the same length of time. Frequent serum cholesterol determinations should be made and the rates of weight gain should be followed. Neglect of any one of these requirements in the evaluation of a drug in the prevention or cure of experimental arteriosclerosis in rabbits leaves the results of the experiment open to question. The requirements of strain or heredity control is

fundamental. Age has been shown<sup>34,35</sup> to be an important factor in determining the ease with which arteriosclerosis can be produced in rabbits. Sex has been noted to play a role in that some male rabbits seem to be rather refractory to the development of hypercholesterolemia and arteriosclerosis on this regimen.<sup>36</sup> The initial weight and the rate of weight gain during the experiment are recognized as having an appreciable influence upon the arteriosclerosis.<sup>7,37</sup> The thriving, weight-gaining rabbits seem to develop more arterial lesions than their lean brethren. The necessity for simultaneous controls cannot be emphasized too strongly, a requirement that is frequently neglected because it requires more in the way of facilities and personnel. There is also the temptation to use a single control group for comparison with several successive test groups. In our own laboratory we have seen strictly comparable control groups of rabbits in successive years show significant differences in the incidence and severity of arteriosclerosis. This is most pronounced if the seasons happen to differ, because rabbits do not do well in summer. Contemporary controls in the same animal quarters also serve as a check on such intangible variables as differences in commercial food lots and unrecognized intercurrent disease in the rabbit colony. These can inhibit the development of arteriosclerosis. Frequent serum cholesterol determinations serve to check the comparability of the control and test animals (when the test substance is to be used in a curative experiment), or to reflect possible effects of the test substance upon the serum cholesterol (in the preventative type experiment). This latter use of cholesterol levels is hazardous because mere acceleration of the dietary cholesterol through the gastrointestinal tract by an irritative effect of the test substance can probably result in lower values. Finally, if all of these conditions have been complied with to achieve the ideally controlled experiment, at the end of such an experiment it usually is impracticable to perform the minimum of twenty control and twenty test animal autopsies within an insignificant interval of time.

As a result of all these difficulties compromises have been made by all investigators with resultant invalidation of many of the experiments. Moreover, if a test group showed less arteriosclerosis than the control group, the test agent was likely to become the subject of a published report; if there was no difference between the

groups, the investigator was generally inclined to dismiss the experiment as inconclusive. Custom seems to require much better data for negative reports than for positive ones.

Most of what has been said of the design and interpretation of rabbit experiments applies in general to chicken and dog studies. Rodbard et al.<sup>38</sup> have shown that age and season of the year profoundly influence the development of arteriosclerosis in chickens. Birds on a uniform regimen show progressive development of arteriosclerosis from the eighth to the twentieth week of age, but after twenty weeks the arteriosclerosis slows down and may even regress. Studies on birds under eight weeks of age showed that arteriosclerosis could be produced in the winter season but not during the summer. In both chickens and dogs spontaneous arteriosclerosis occurs with old age. Thus 50 per cent of chickens have arteriosclerosis at the age of one and one-half years,<sup>4</sup> and Bevans has found arteriosclerosis in all of twenty-seven dogs between nine and nineteen years of age.<sup>39</sup> Katz and co-workers have favored the use of young chickens, large numbers of which can be conveniently used to cancel out biologic variations. Kendall and his group have used dogs because they believe that the experimental canine disease is more closely analogous to human arteriosclerosis and that there is less variation in the dog than in the rabbit.

Most of the experiments cited by pharmaceutical firms in support of the use of lipotropic agents in arteriosclerosis are open to criticism on one or more of the above counts. When the literature is reviewed with these requirements for adequately controlled experiments in mind, there are only a few that escape rejection. In general, the results of even these experiments are suggestive rather than conclusive.

Steiner<sup>36</sup> in 1938 reported that choline administered to nineteen six-months-old male rabbits on a high cholesterol diet over periods of 40 to 100 days resulted in only nine grossly arteriosclerotic animals, compared with fifteen diseased animals in the group of nineteen control rabbits on the same regimen without added choline. He stated that "on a statistical basis the results are not striking," and concluded that "choline delays, but does not prevent, arteriosclerosis in cholesterol-fed rabbits." That same year Steiner<sup>40</sup> reported a curative type of experiment in which twenty six-month-old male rabbits were fed cholesterol for 110 days. This

lot of arteriosclerotic animals was then divided into three groups. Five rabbits were sacrificed at once; the remainder were placed on plain food without cholesterol for sixty more days with ten of the latter receiving 0.5 gm. of choline daily. At the end of the experimental period all rabbits were sacrificed. The incidence of arteriosclerosis in the first five rabbits sacrificed at the end of the cholesterol feeding was the same as in the five which were on plain food alone for sixty more days, and both of these groups showed a higher incidence of arteriosclerosis than the remaining ten rabbits which had received choline. The number of animals used in these experiments was small and the serum cholesterol levels of the group which served as controls were somewhat higher than those of the choline group. In the same year Himsworth<sup>41</sup> studied the effects of choline in cholesterol-fed rabbits. He used littermate five to six month old animals divided into six groups with too few animals per group to permit any conclusions. He believed, however, that choline was without any effect upon the arteriosclerosis. A quite similar experiment was carried out by Baumann and Rusch<sup>42</sup> that year with the same conclusions, but again based on two experiments each involving only twelve rabbits which were divided into three groups. Cholesterol levels were controlled but the animals varied too much in weight and were too few per group to be conclusive. The Keston and Silbowitz study<sup>43</sup> in 1942 of soya lecithin and choline in rabbits fed cholesterol involved only seven or eight rabbits per group, and included no mention of any serum cholesterol levels. Their control group of eight rabbits showed arteriosclerosis in seven, while three test groups of seven, eight and eight on lecithin or choline showed two, two and three diseased animals, respectively. These earlier experiments were all of a pilot nature and, in the light of what has since become known about factors influencing experimental arteriosclerosis in rabbits, were inadequately controlled.

More recently several studies of choline in rabbits have reported it to be effective against experimental arteriosclerosis. Morrison<sup>44,45</sup> investigated the prevention of arteriosclerosis in cholesterol-fed rabbits. He reported an incidence of 75 per cent moderately severe (+++) to marked (++++) arteriosclerosis, with 5 per cent of animals free of lesions in his twenty control rabbits after ninety-two days on the diet. Twenty-nine similar animals which received



0.5 gm. choline hydrochloride per day in addition to the diet showed not even moderately severe (+++) arteriosclerosis, and 55 per cent were entirely free of lesions. On twice this choline dosage 78 per cent of the thirty-two rabbits were free of lesions. These figures are impressive but there is no mention of serum cholesterol levels and no statement as to whether the test and control animals were run simultaneously. If these two variables were not controlled, the observed differences could well have arisen through either of these important factors. Morrison<sup>45</sup> further reported a curative type of experiment in which arteriosclerosis was produced in forty-four rabbits by cholesterol feeding for 184 days. The rabbits were then divided and half got plain food alone as controls, while the other half received plain food plus choline for the next 185 days. In the twenty-one surviving control rabbits there were none free of arteriosclerosis. Of the twenty-two choline-treated animals 74 per cent were free of lesions. Again there were no serum cholesterol or weights mentioned to assure comparability of the two groups, and it is not stated that they were run simultaneously. Broun et al.,<sup>34</sup> using a total of 224 cholesterol-fed rabbits, fed groups of six to eighteen animals choline, betaine, methionine and inositol at various dosage levels. Some groups of rabbits were three months old, others six to eight months old. The eighteen rabbits on 40 to 160 mg. per day of betaine showed no effect of the medication. Similarly, 40 mg. of methionine and 40 mg. of inositol in groups of six rabbits each were inactive. In the case of choline, 40 mg. per day (the largest dose) did nothing in the six to eight month old rabbits, but in thirty month old rabbits doses from 2 to 500 mg. of choline were tested and the doses of 75 mg. per day and up seemed to inhibit the production of arteriosclerosis. This might have been due to depression of appetite by the choline but the authors reported that the various groups of animals showed "little difference in weight gain." These three month old rabbits showed serum cholesterol levels of only 400 to 700 mg. per cent as compared with several times this level in the older animals, so this might be a partial choline effect which can be overcome by greater hypercholesterolemia. The choline doses reported as effective are in line with those that have been generally used by other investigators. It is most interesting that when the regular diet was supplemented with 1 gm. of Crisco per day

the maximum 200 mg. of choline per day lost its effectiveness in protecting the animals against arteriosclerosis. This suggests that the basic diet in these experiments might have been unusually low in fat or deficient in choline. Steiner<sup>46</sup> has repeated and extended his earlier work on choline in rabbits. This time twenty-five controls received a high cholesterol diet alone and twenty-nine rabbits got this plus choline hydrochloride at two dosage levels, 500 mg. per day for eighteen animals and 1.0 gm. per day for eleven animals. Animals were sacrificed in pairs from each group (four animals) at intervals of five to ten days between the 40th and the 100th day of the experiment. None of the choline-treated animals sacrificed between the 40th and the 80th day showed arteriosclerosis, while ten of eleven control rabbits showed lesions at the end of the same period. The serum cholesterol levels of the two groups were quite similar but a little lower in the control group. No data were given as to the weights of the animals so there was no quantitative control on the relative rates of gain in the control and test groups. Recently this has been claimed to have a significant effect in rabbit arteriosclerosis<sup>7</sup> and could readily be a factor when the diet of one group is flavored by choline.

These conflicting past reports on the effect of lipotropic agents upon experimental arteriosclerosis in rabbits have stimulated several carefully controlled studies recently. Meyer et al., in an unpublished experiment,<sup>47</sup> studied choline for possible curative action. Arteriosclerosis was produced in forty-three chinchilla rabbits by cholesterol feeding for 107 days. Serum cholesterol determination were obtained every two weeks and the figures averaged to yield a mean serum cholesterol level for the entire 107-day period for each animal. On the basis of these cholesterol averages the rabbits were then divided into three groups, matched in their distribution of mean serum cholesterol levels and therefore as comparable as possible in the incidence and severity of their arteriosclerosis. Group A, containing thirteen rabbits, was sacrificed at once to determine the degree of arteriosclerosis in this representative sample of the colony. The average of the mean serum cholesterol levels in this group was 1005 mg. per cent. Autopsies were performed and the arteriosclerosis found was graded on a scale of 0 to 4+ in (1) the ascending aorta, (2) the thoracic aorta, (3) the abdominal aorta and (4)



the pulmonary artery. As an additional index of the severity of the lesions, chemical analyses were made of each individual aortic intima and media for its content of various lipids. The remaining two groups of rabbits were continued on experiment. Group B, containing fourteen

designed to study the effect of choline in prevention of arteriosclerosis. The animals were all males from a single colony and between ten and twelve months of age. Serum cholesterol levels were determined weekly and each rabbit was weighed twice a week. Three groups of ten

TABLE I

Average Serum Cholesterol mg. % during:			Gross Pathology Atherosclerotic Involvement (Scale of 0-4+)				Analysis of Aortic Intima and Media Lipids (gm./100 gm. Dry Wt.)						
Choles- terol Feeding Period		Regres- sion Period	Ascend- ing Arch of Aorta	Thoracic Aorta	Ab- dominal Aorta	Pulmo- nary Artery	Dry Wt. gm.	Cholesterol			Phospho Lipid (Lipid P × 25)	Total Lipid	Neutral Fat
								Total	Free	Ester			
A	1005	...	3.1	1.7	1.7	2.8	0.220	2.52	1.20	1.32	2.97	23.4	16.9
B	1068	600	3.7	2.4	2.0	2.6	0.176	10.10	4.08	6.02	4.55	25.6	6.4
C	1067	646	3.7	2.5	2.3	2.6	0.166	11.88	5.07	6.81	5.63	28.7	6.2

animals with an average mean serum cholesterol level of 1068 mg. per cent, was placed on a plain stock diet as controls against further changes in the lesions. Group C, containing twelve rabbits with an average mean serum cholesterol level of 1067 mg. per cent, was placed on the same plain stock diet with 1 gm. of choline hydrochloride in aqueous solution mixed with a small first portion of each rabbit's daily food. This technic assured consumption of the choline without having the rabbits' entire diet flavored by the choline. After 112 days on their respective regimens the rabbits in groups B and C were sacrificed and studied in the same manner as group A. The results in terms of averages for each group of rabbits are given in Table I. It should be noted that rabbits in groups B and C remained hypercholesterolemic during the regression period of the experiment despite no cholesterol intake. The arteriosclerosis in these groups was accordingly more severe than in group A. This rabbit experiment gave no evidence of any effect of choline upon the experimental arteriosclerosis. A similar failure to demonstrate effects of choline on regression of arteriosclerosis in rabbits (in unpublished experiments) was mentioned by Duff.<sup>1</sup>

Moses and Longabaugh<sup>37</sup> have reported quite a similar cholesterol-fed rabbit experiment

rabbits each were used and all were fed the same high cholesterol diet. One group received nothing else and served as controls. Group two had 1 gm. of choline salt mixed with the food of each rabbit daily. Group three similarly received choline but in doses of 4 gm. per day per rabbit. After six weeks on the experimental diets all animals were sacrificed and examined. Addition of the grades of arteriosclerosis in the animals in each group showed a score of 24 for the controls, 19 for the second group (on 1 gm. choline) and 26 for the third group (on 4 gm. choline). These figures fail to show an influence of choline upon the lesions. Despite this, there was one animal free of lesions in group one, as opposed to two in group two and three in group three. This might be interpreted as indicative of a protective effect of choline but is offset by a higher incidence of very severe arteriosclerosis in the choline-fed animals plus the fact that the average weight gain in the five test animals free of lesions was 12 gm. as opposed to an average gain of 416 gm. in the entire control group. This was a well controlled experiment but is marred by the marked biologic variation in the small numbers of animals in each group and the observed lower weight gain in the choline-fed animals, a complicating factor which could inhibit arteriosclerosis. Both factors

make it impossible to conclude that choline had any unequivocal influence upon the arteriosclerosis.

The effect of inositol has recently been studied in rabbits in two unpublished experiments. Ellenbogen and Kendall<sup>48</sup> used twenty-eight

gained less and had normal serum cholesterol levels, as would be expected. When the animals were examined after twelve weeks on their diets, the grade of arteriosclerosis (on a 0 to 4+ scale) in the cholesterol-without-inositol group averaged 2.6+; in the cholesterol-plus-inositol

TABLE II

Regimen	No. of Rabbits	Average Serum Levels			Wt. Gain (gm.)	Estimated Degree (Scale of 0-4+)	
		Cholesterol (mg. %)	Lipid Phosphorus (mg. %)	Moles Chol./Phos.		Arteriosclerosis	Fatty Liver
Stock diet. ....	7	56	4	1.1	625	0	0
Stock diet + 3 gm. inositol per week. ....	7	45	4	1.0	570	0	0
Cholesterol-containing diet. ....	7	667	12	4.6	550	1+, 1+, 1+, 1+, 1+, 2+, 2+	1+, 1+, 2+, 2+, 3+, 4+, 4+
Cholesterol-containing diet + 3 gm. inositol per week. ....	6	781	13	4.7	635	1+, 2+, 2+, 2+, 3+, 3+	2+, 3+, 3+, 3+, 3+, 4+

rabbits of the same stock. These were divided into four groups of matching body weights. Two groups were controls on stock diet and stock diet plus inositol, respectively. The third group was fed a high cholesterol diet alone, while the fourth group received the high cholesterol diet plus 0.5 gm. inositol six days a week. In all animals the serum cholesterol and phospholipids were followed and the gain in weight was measured. The animals were on experiment simultaneously in the same animal quarters. All animals were sacrificed and examined after eighty-four days. The results are given in Table II. This experiment was well controlled but suffered from the compromise made in using too few rabbits. There is, however, no indication that inositol had any effect upon the serum lipid levels, the amount of fat deposited in the liver, or the amount of arteriosclerosis produced. The second inositol study was carried out by Moses.<sup>49</sup> Though designed completely independently of Ellenbogen et al. it followed strikingly parallel lines. Moses used all male albino rabbits. He simultaneously had seven rabbits on cholesterol diet alone, nine on cholesterol diet plus 2 gm. inositol per rabbit daily, and nine on plain food plus 2 gm. inositol per rabbit daily. Weight gain and serum cholesterol levels were determined and were strictly comparable in the two cholesterol-fed groups. The third group on inositol without cholesterol

group 3.7+; and in the inositol-without-cholesterol group 0.0+. There was no evidence of any inositol effect but the numbers of animals were small so Moses did a second experiment. This time three groups of twenty-six comparable rabbits were used. Each group was fed a different level of cholesterol in its diet. Of each twenty-six, eight received no inositol and served as controls, while the remaining eighteen in cages of six were given 4, 8 and 12 gm. of inositol per cage in their drinking water daily. This division of animals permitted the study of a range of inositol dosage in rabbits under various dietary cholesterol loads. Serum cholesterol levels and the weights of the animals were followed and there was no evidence of any effect of inositol upon these measurements. Two animals from each of the twelve groups on different regimens were sacrificed after four, six and eight weeks on experiment. This added various time intervals to the already different regimens. The arteriosclerosis in the inositol-fed rabbits was in all instances comparable to that found in the corresponding control animals. In this experiment inositol was studied over a range of dosages and under different conditions of time and cholesterol feeding. This reduced the number of animals per subgroup to only two or four, numbers too small to be significant if a group differed from the bulk of the data. Since there was no such deviation of any of the groups, the

experiment as a whole seems highly significant and provides rather convincing evidence that inositol has no influence upon the development of arteriosclerosis under these conditions.

Another unpublished study of lipotropic agents in cholesterol-fed rabbits has been made by Kellner.<sup>50</sup> Soya lecithin preparations containing both choline and inositol were employed. He used approximately 100 rabbits divided into experimental groups of ten and twelve. Some were maintained on plain food, some on plain food plus lecithin, some on a cholesterol-containing diet alone, and some on this last diet plus lecithin. The doses of lecithin were calculated to be proportional to the doses recommended by the manufacturer for use in human subjects. All animals were followed as to weight and both serum cholesterol and phospholipid levels. Kellner found that lecithin caused no change in the serum phospholipid levels in the animals on plain food. In the cholesterol-fed rabbits there was no difference between the controls and the lecithin-treated groups in regard to serum cholesterol-to-phospholipid ratios or in the resultant arteriosclerosis.

Evaluation of the lipotropic agents in arteriosclerosis in chickens has been reported frequently by Herrmann<sup>51-54</sup> and by Stamler et al.<sup>55,56</sup> The former believes that the lipotropic agents are effective while the latter has been unable to demonstrate any favorable action.

Herrmann used old hens as his experimental bird, because of their high incidence of spontaneous arteriosclerosis. He followed the blood and tissue cholesterol concentrations as indices of the efficacy of the drugs. He fed choline,<sup>51</sup> methionine<sup>53</sup> and inositol<sup>52</sup> in efforts to reduce the blood cholesterol levels. All of these agents produced statistically significant blood cholesterol reductions in apparently adequate numbers of birds as compared with control bleedings in the same birds. Upon analysis of the heart, aorta and liver of the treated and simultaneous control hens there was significantly less cholesterol found in the organs of the treated hens. The experimental data appear to justify Herrmann's conclusions that choline, methionine and inositol did reduce the blood cholesterol level and diminish the tissue cholesterol content in his old hens. The significance of these findings with regard to arteriosclerosis is chiefly inferential. No report has been made of the direct effect of the lipotropic agents upon the arteriosclerotic lesions.

Stamler et al. studied the effect of choline and inositol upon the development of arteriosclerotic lesions in young chicks. The lesions studied were of three types: those induced by cholesterol feeding,<sup>56</sup> those induced by stilbestrol<sup>55</sup> and spontaneously occurring lesions.<sup>55</sup> In the cholesterol-induced arteriosclerosis experiment ninety-six birds were divided into six groups of twelve test birds each and one control group of twenty-four birds. All ninety-six birds were fed the same basic diet but one pair of test groups was given 0.25 per cent cholesterol in their diet; a second pair, 0.5 per cent cholesterol; and the third pair 2.0 per cent cholesterol. One of each pair of groups on a given diet was also given a combination of 1 per cent choline and 1 per cent inositol in its particular cholesterol-enriched diet. All birds were weighed weekly, food intake was recorded and bloods were drawn at intervals for complete serum lipid fractionation. Birds from test and control groups were sacrificed at intervals after fifteen to forty-seven weeks of experimental feeding. Under these conditions choline and inositol had no prophylactic effect against hypercholesterolemia, hyperlipemia, aortic lipidosis or atherosclerosis. On the contrary, the lipotropic factors tended to aggravate both the hyperlipemia and the atherosclerosis. In the stilbestrol-induced arteriosclerosis experiment twenty-four birds were implanted with stilbestrol pellets and twelve of these served as controls on regular diet while twelve were fed the same diet containing 1 per cent choline and 1 per cent inositol. The conduct of the experiment was the same as in the case of the study of cholesterol-fed birds. There was no evidence of any effect of the lipotropic agents upon blood or tissue lipids or the arteriosclerosis. The study of spontaneous arteriosclerosis was carried out in an analogous manner with forty-eight birds. Half were on regular diet and half on the lipotropic diet. The results were completely negative. These experiments differ in many ways from those of Herrmann but seem to bear more directly on the problem. They showed no evidence of any inhibitory action of choline and inositol upon arteriosclerosis in chicks.

The influence of choline and inositol upon the development and regression of experimental arteriosclerosis in the dog has been investigated by Kendall and co-workers.<sup>57</sup> Experimental arteriosclerosis can be produced in dogs by feeding cholesterol and thiouracil.<sup>58-60</sup> The experimental canine arterial lesions seem mor-



phologically and anatomically more similar to human arteriosclerosis than either the rabbit or chicken lesions. This as well as the fact that the dog is omnivorous like man has made our group favor this experimental animal. Choline was studied first and was tested for a possible inhibitory effect upon the production of arteriosclerosis.<sup>57</sup> In a pilot experiment three dogs were fed cholesterol, thiouracil and 5 gm. of choline hydrochloride per day for fourteen months. Their hypercholesterolemia was comparable to what had been observed previously in five dogs on the same regimen without choline. At autopsy all three animals had extensive gross arteriosclerosis. A better controlled experiment was then performed. Fourteen mongrel dogs were started on experiment at four months of age and fed a 5 per cent cholesterol diet ad libitum. The dogs were individually given 0.6 gm. thiouracil daily. Eight of the dogs were maintained on this regimen as controls while six were given choline hydrochloride mixed with their food in an amount equal to 2.5 per cent of the diet. Serum cholesterol determinations were made in all dogs every two weeks and four to seven of the samples of serum from each dog were also analyzed for phospholipid content. All of the choline-fed dogs were on experiment at the same time but only four of the control dogs were on experiment concurrently with them. After four months on the experiment all dogs were sacrificed and examined. It was not possible to demonstrate any effect of the choline upon the treated dogs as compared with the controls in respect to serum levels of cholesterol or phospholipids, in degree of fatty infiltration of the liver, or in the extent and severity of the arteriosclerosis which was present in all animals. In a subsequent unpublished experiment<sup>61</sup> choline was studied for possible curative effects. Arteriosclerosis was produced in six dogs by cholesterol and thiouracil feeding for six months. The dogs were then taken off this regimen and placed on plain food for four months. Under these conditions the arterial lesions are known to regress.<sup>5</sup> During this last four-month regression period 2½ per cent choline hydrochloride was added to the plain diet. At the end of the regression period the extent of the arterial lesions in these six choline-treated animals was compared with that in nine dogs observed after being placed on the same experimental regimen without added choline. The controls were not all run concurrently with the test dogs. There

was no difference between the two groups of dogs when animals of similar degree of effective hypercholesterolemia were compared.

Inositol was studied in ten dogs in the preventative type of experiment with no apparent effect of the drug on the serum lipids or the resultant arteriosclerosis, as compared with fifteen control dogs. The inositol dosage was 3 gm. per dog per day. The effect of inositol upon the regression of arteriosclerosis in dogs is currently being investigated.

In summary, the data on studies of the value of lipotropic agents in arteriosclerosis in experimental animals are conflicting. It does seem, however, that the more recent and better controlled studies of the action of inositol and choline in both rabbits and chickens have not demonstrated any effect of these agents either in preventing or curing arteriosclerosis, or in influencing the serum lipid pattern in what would seem to be a favorable manner. Only our own group has studied these agents in the dog but no beneficial effects have been demonstrated in this species.

*Clinical Studies.* Very few clinical studies of the effects of lipotropic agents upon human arteriosclerosis have been reported despite wide use of these compounds in medical practice. Controlled clinical studies are difficult because of the chronicity and marked variability of the disease. The diagnosis of arteriosclerosis is most secure in patients who have had a myocardial infarction so these are the patients in whom the drugs can best be evaluated. If evaluation is to be direct, the patients have to be followed while on the medication until sufficient numbers of deaths or complications from arteriosclerosis have arisen in comparable series of treated and control patients. This requires large numbers of patients and many years of observation. Throughout this period there is always the danger that the patients on the drug may get more careful medical attention than the controls and that thereby factors other than the drug under test may favorably influence the test patients' course.

One such "direct" study has been reported by Morrison and Gonzales in several articles.<sup>14, 62-64</sup> They used "alternate unselected patients" who survived proven coronary thromboses as their control and test groups. There were 115 patients in each group, with the sex, age and hypertension distributions reported comparable. The test group was given choline

in doses of 6 to 32 gm. per day, the average dose being 12 gm. These patients were then followed for periods up to three years and the mortality tabulated. There were thirty-five deaths in the control group and fourteen in the treated group. These figures are significantly different statistically and remain so upon breakdown into the deaths from recurrent coronary thrombosis (19 vs. 6), cardiac failure (10 vs. 5) and non-cardiac causes (6 vs. 3). This last category raises the question as to whether greater medical care for the patients given choline may not have influenced the results. The authors have not made this point entirely clear and there seems to be no specific accounting for individuals in this study of 230 patients who must have been lost to follow-up. A completely satisfactory experiment of this type should utilize placebo medication in the control group.

Less difficult indirect studies of lipotropic agents for human arteriosclerosis have been made by using the level of the serum cholesterol and serum phospholipids as indices of the possible action of the drugs. As with animals, this index is probably valid. In this type of experiment the chief pitfalls are the marked spontaneous fluctuation of the serum cholesterol level in patients with arteriosclerosis<sup>65,66</sup> and the fact that weight loss can cause reduction of the serum cholesterol level.<sup>67</sup> The latter is emphasized because both choline and inositol in doses that have been used can cause gastroenteritis and anorexia with possible resultant weight loss.

Soya lecithin was studied by Steiner and Domanski<sup>68</sup> in six patients with coronary arteriosclerosis and two patients with rheumatoid arthritis. A reduction of serum cholesterol was observed in all patients, with no difference between the two types of patients. The average maximum serum cholesterol decline was 68 mg. per cent. The decline, however, was maintained for only five weeks after which the serum cholesterol rose again despite continued soya lecithin feeding. Sufficiently frequent and numerous serum cholesterol determinations were made but there was no report of the weights of the patients. A fall in weight upon initiation of the lecithin feeding with later adaptation to the regimen and maintenance of weight could account for the findings in this experiment.

Herrmann has published several reports of studies of the effects of choline, methionine and inositol upon the serum lipid levels in humans.<sup>64,69,70</sup> He gave choline in doses of 1 gm. three times a day to forty patients for

periods of one and one-fourth to three and three-fourths months and observed an average reduction of 20 per cent in the total serum cholesterol. In the case of 0.5 gm. methionine given three times daily for two to four weeks to six patients, the reduction was reported as 10 to 15 per cent, with one patient showing a 33 per cent increase. In twenty patients given 0.5 gm. of inositol four times a day for twenty-five to thirty days there was a 19 per cent average reduction of serum cholesterol level. In all of these reports there is no mention of what happened to the weights of the patients or any record of the number of baseline serum cholesterol levels made in each patient to evaluate random fluctuation. The number of patients in these studies was generally too small to permit any definite conclusions. More recently Herrmann<sup>74</sup> has reported on larger numbers of patients on choline and on inositol regimens. Only average figures are given for the serum cholesterol levels before and after treatment. The serum cholesterol reductions produced amounted to 10 to 20 per cent of the initial levels. Similar studies on soya lecithin showed no significant reduction.

Morrison<sup>71</sup> attempted to study the effects of choline on the serum cholesterol levels of forty-eight patients who had had recent coronary thromboses. He reported that in these patients given 6 gm. of choline per day there were such marked fluctuations of the serum cholesterol level that it was impossible to determine whether a reduction or an increase predominated. In contrast to these arteriosclerotic patients a small group of normal people fed choline by Paul, Daum and Kemp<sup>72</sup> showed very little variation in their serum cholesterol levels, with as many showing a rise as a fall.

Inositol has been studied in diabetics by Felch.<sup>73</sup> A group of thirty diabetics selected for hypercholesterolemia and stability of their diabetes were fed 3 gm. of inositol daily for eight weeks. Serum cholesterol and lipid phosphorus determination were made two or more times before the start of inositol and after one, two, four and eight weeks on the drug. Both lipids were found to fall appreciably while the patients were on inositol. In this study variations in the degree of control of the diabetes alone could have accounted for the results. This possibility is further suggested by the fact that the lipids returned only part way to their baseline levels while the patients were followed for six weeks after inositol was discontinued. Unless



placebo medication is used in a control study in a group such as this greater medical attention during the trial of the drug may inadvertently result in more strict adherence to diet, with consequent better regulation of the diabetes and a favorable effect upon the blood lipid levels.

In all of these clinical studies, as in the animal studies, it is always difficult to be certain that reductions in serum cholesterol levels upon administration of lipotropic agents are not due to secondary factors such as weight loss. This is very likely to occur when choline or inositol are fed, due to production of anorexia and diarrhea by these compounds.

#### CONCLUSIONS

Under certain circumstances low cholesterol, low fat diets are capable of lowering the levels of serum cholesterol. The degree of dietary lipid restriction necessary to achieve this effect in some patients may preclude the use of this regimen. It is not established that the maximum depression of serum cholesterol level obtainable by dietary measures will affect arteriosclerosis in man. Evidence in the experimental animal, however, suggests that a restricted lipid intake produces regression of experimentally induced arteriosclerotic lesions.

Consideration of all data available to date indicates that despite individual positive reports there is no general agreement that choline or inositol have any specific influence upon arteriosclerosis or the serum cholesterol level in man or the experimental animal.

#### REFERENCES

1. DUFF, G. L. and McMILLAN, G. C. Pathology of atherosclerosis. *Am. J. Med.*, 11: 92, 1951.
2. GOULD, R. G. Lipid metabolism and atherosclerosis. *Am. J. Med.*, 11: 209, 1951.
3. JONES, H. B., GOFMAN, J. W., LINDGREN, F. T., LYON, T. P., GRAHAM, D. M., STRISOWER, B. and NICHOLS, A. V. Lipoproteins in atherosclerosis. *Am. J. Med.*, 11: 358, 1951.
4. HORLICK, L., KATZ, L. N., STAMLER, J. The effect of a low fat diet on the spontaneously occurring arteriosclerosis of the chicken. *Am. Heart J.*, 37: 689, 1949.
5. BEVANS, M., DAVIDSON, J. D. and KENDALL, F. E. Regression of lesions in canine arteriosclerosis. *Arch. Path.*, 51: 288, 1951.
6. HORLICK, L. and KATZ, L. N. Retrogression of atherosclerotic lesions on cessation of cholesterol feeding in the chick. *J. Lab. & Clin. Med.*, 34: 1427, 1949.
7. FIRSTBROOK, J. B. Effects of changes in body weight on atherosclerosis in rabbits. *Science*, 111: 31, 1950.
8. WILENS, S. L. Bearing of general nutritional state on atherosclerosis. *Arch. Int. Med.*, 79: 129, 1947.
9. OPPENHEIM, F. Review of 100 autopsies of Shanghai Chinese. *Chinese M. J.*, 39: 1067, 1925.
10. SNAPPER, I. Chinese Lessons to Western Civilization, p. 160. New York, 1941. Interscience Publishers.
11. STEINER, P. E. Necropsies on Okinawans. *Arch. Path.*, 42: 359, 1946.
12. WILENS, S. L. The resorption of arterial atheromatous deposits in wasting disease. *Am. J. Path.*, 23: 793, 1947.
13. MORRISON, L. M., ZWERLEIN, M. and WOLFSON, E. The effects of low fat-low cholesterol diets on the serum lipids. *Circulation*, 2: 475, 1950.
14. MORRISON, L. M. Arteriosclerosis. *J. A. M. A.*, 145: 1232, 1951.
15. KEMPNER, W. Treatment of heart and kidney disease and of hypertensive and arteriosclerotic vascular disease with the rice diet. *Ann. Int. Med.*, 31: 821, 1949.
16. STARKE, H. Effect of the rice diet on the serum cholesterol fractions of 154 patients with hypertensive vascular disease. *Am. J. Med.*, 9: 494, 1950.
17. WATKIN, D. M., FROEB, H. F., HATCH, F. T. and GUTMAN, A. B. Results with unmodified Kempner rice diet in fifty hospitalized patients. *Am. J. Med.*, 9: 441, 1950.
18. MELLINKOFF, S. M. and MACHELLA, T. E. and REINHOLD, J. G. The effect of a fat-free diet in causing low serum cholesterol. *Am. J. M. Sc.*, 220: 203, 1950.
19. HILDRETH, E. A., MALINKOFF, S. M., BLAIR, G. W. and HILDRETH, D. M. Effect of vegetable fat ingestion on human serum cholesterol concentration. *Circulation*, 3: 641, 1951.
20. WILKINSON, C. F., BLECHA, E. and REIMER, A. Is there a relation between diet and blood cholesterol? *Arch. Int. Med.*, 85: 389, 1950.
21. GERTLER, M. M., GARN, S. M. and WHITE, P. D. Diet, serum cholesterol and coronary artery disease. *Circulation*, 2: 696, 1950.
22. KEYS, A., MICKELSON, O., MILLER, E. V. O. and CHAPMAN, C. B. The relation in man between cholesterol levels in the diet and in the blood. *Science*, 112: 79, 1950.
23. GOFMAN, J. W., JONES, H. B., LINDGREN, F. T., LYON, T. P., ELLIOT, H. A. and STRISOWER, B. Blood lipids and human atherosclerosis. *Circulation*, 2: 161, 1950.
24. HATCH, F. T. and KENDALL, F. E. Effect of a cholesterol-free, low-fat diet upon the serum lipids in patients with essential hypertension. *Circulation*, 4: 472, 1951.
25. BEST, C. H., LUCAS, C. C., PATTERSON, J. M. and RIDOUT, J. M. The rates of lipotropic action of choline and inositol under special dietary conditions. *Biochem. J.*, 48: 452, 1951.
26. MCHENRY, E. W. and PATTERSON, J. M. Lipotropic factors. *Physiol. Rev.*, 24: 128, 1944.
27. DOLAN, R. A. Choline and other lipotropic factors. *Minnesota Med.*, 31: 1198, 1948.
28. BEST, C. H. and HUNTSMAN, M. E. Effects of components of lecithine upon deposition of fat in liver. *J. Physiol.*, 75: 405, 1932.
29. BEST, C. H. American Lecture Series No. 21. Diabetes and Insulin and the Lipotropic Factors, p. 25. Springfield, Ill., 1948. Charles C Thomas.
30. GAVIN, G. and MCHENRY, E. W. Inositol: a lipotropic factor. *J. Biol. Chem.*, 139: 485, 1941.



31. MACFARLAND, M. L. and MCHENRY, E. W. Further observations on the lipotropic need for inositol. *J. Biol. Chem.*, 176: 429, 1948.
32. BEST, C. H., RIDOUT, J. H., PATTERSON, J. M. and LUCAS, C. C. A statistical evaluation of the lipotropic action of inositol. *Biochem. J.*, 48: 448, 1951.
33. SEEGAL, D. et al. Combined staff clinic on cholesterol metabolism and arteriosclerosis. *Am. J. Med.*, 6: 103, 1949.
34. BROUN, G. O., ANDREWS, K. R., CORCORAN, P. J. V. and VAN BRUGGEN, J. Studies on the effect of lipotropic agents in experimental cholesterol arteriosclerosis. *Geriatrics*, 4: 178, 1949.
35. POLLAK, O. J. Age and weight as factors in the development of experimental cholesterol atherosclerosis in rabbits. *Arch. Path.*, 43: 387, 1947.
36. STEINER, A. Effect of choline on production of experimental arteriosclerosis in rabbits. *Proc. Soc. Exper. Biol. & Med.*, 38: 231, 1938.
37. MOSES, C. and LONGABAUGH, G. M. Evaluation of choline in the prevention of experimental atherosclerosis. *Arch. Path.*, 50: 179, 1950.
38. ROBBARD, S., KATZ, L. M., BOLENE, B. A., PICK, R., LOWENTHAL, M. and GROS, G. The age factor in hypercholesteremia and atheromatosis in the chick. *Circulation*, 3: 867, 1951.
39. BEVANS, M., TAYLOR, B. and ABELL, L. L. Spontaneous canine arteriosclerosis. *Circulation*, 2: 477, 1950.
40. STEINER, A. Action of choline on experimental aortic atherosclerosis. *Proc. Soc. Exper. Biol. & Med.*, 39: 411, 1938.
41. HIMSWORTH, H. P. Experimental atherosclerosis. *Acta med. Scandinav., Suppl.*, 90: 158, 1938.
42. BAUMANN, C. S. and RUSCH, H. P. Effect of choline on atherosclerosis in the rabbit aorta. *Proc. Soc. Exper. Biol. & Med.*, 38: 647, 1938.
43. KESTON, H. D. and SILBOWITZ, R. Experimental atherosclerosis and soya lecithin. *Proc. Soc. Exper. Biol. & Med.*, 49: 71, 1942.
44. MORRISON, L. M. The prevention of experimental arteriosclerosis by choline feeding. *Geriatrics*, 4: 236, 1949.
45. MORRISON, L. M. and ROSSI, A. Absorption of aortic atherosclerosis by choline feeding. *Proc. Soc. Exper. Biol. & Med.*, 69: 283, 1948.
46. STEINER, A. Effect of choline in the prevention of experimental aortic atherosclerosis. *Arch. Path.*, 45: 327, 1948.
47. MEYER, W., ABELL, L. L. and KENDALL, F. E. Personal communication.
48. ELLENBOGEN, L. and KENDALL, F. E. Personal communication.
49. MOSES, C. Personal communication.
50. KELLNER, A. Personal communication.
51. HERRMANN, G. R. Effect of choline on blood and tissues with especial reference to cholesterol in old hens. *Proc. Soc. Exper. Biol. & Med.*, 61: 302, 1946.
52. HERRMANN, G. R. Inositol decholesterization in old hens. *Proc. Soc. Exper. Biol. & Med.*, 63: 436, 1946.
53. HERRMANN, G. R. Methionine decholesterization. *Proc. Soc. Exper. Biol. & Med.*, 64: 284, 1947.
54. HERRMANN, G. R. Some experimental studies in hypercholesterolemic states. *Exper. Med. & Surg.*, 5: 149, 1947.
55. STAMLER, J., BOLENE, C., HARRIS, R. and KATZ, L. N. Effect of choline and inositol on plasma and tissue lipids and on spontaneous and stilbestrol-induced atherosclerosis in the chick. *Circulation*, 2: 722, 1950.
56. STAMLER, J., BOLENE, C., HARRIS, R. and KATZ, L. N. Effect of choline and inositol on plasma and tissue lipids and atherosclerosis in the cholesterol-fed chick. *Circulation*, 2: 714, 1950.
57. DAVIDSON, J. D., MEYER, W. and KENDALL, F. E. Effect of choline upon experimental canine arteriosclerosis. *Circulation*, 3: 332, 1951.
58. STEINER, A. and KENDALL, F. E. Atherosclerosis and arteriosclerosis in dogs following ingestion of cholesterol and thiouracil. *Arch. Path.*, 42: 433, 1946.
59. STEINER, A., KENDALL, F. E. and BEVANS, M. Production of arteriosclerosis in dogs by cholesterol and thiouracil feeding. *Am. Heart J.*, 38: 34, 1949.
60. BEVANS, M., DAVIDSON, J. D. and ABELL, L. L. The early lesions of canine arteriosclerosis. *Arch. Path.*, 51: 278, 1951.
61. KENDALL, F. E., MEYER, W., ABELL, L. L., BEVANS, M., ELLENBOGEN, L. and DAVIDSON, J. D. Unpublished data.
62. MORRISON, L. M. and GONZALES, W. F. Results of treatment of coronary arteriosclerosis with choline. *Am. Heart J.*, 38: 471, 1949.
63. MORRISON, L. M. and GONZALES, W. P. Effect of choline as a lipotropic agent in the treatment of human coronary atherosclerosis. *Proc. Soc. Exper. Biol. & Med.*, 73: 37, 1950.
64. MORRISON, L. M. and GONZALES, W. F. Results of treatment of coronary arteriosclerosis with choline. *Am. Heart J.*, 39: 729, 1950.
65. STEINER, A. Cholesterol in arteriosclerosis with special reference to coronary arteriosclerosis. *M. Clin. North America*, 34: 673, 1950.
66. STEINER, A. Significance of cholesterol in coronary arteriosclerosis. *N. Y. State J. Med.*, 48: 1814, 1948.
67. WALKER, W. J. and WIER, J. A. Plasma cholesterol levels during rapid weight reduction. *Circulation*, 3: 864, 1951.
68. STEINER, A. and DOMANSKI, B. Effect of feeding "soya lecithin" on serum cholesterol level of man. *Proc. Soc. Exper. Biol. & Med.*, 55: 236, 1944.
69. HERRMANN, G. R. Cholesterol levels in various diseases and the effects of decholesterizing agents. *Texas State J. Med.*, 42: 260, 1946.
70. HERRMANN, G. R. Hypercholesterolemia and the effects of lipotropic agents. *Tr. A. Am. Physicians*, 60: 160, 1947.
71. MORRISON, L. M., CHANEY, A. L., HALL, L. and GONZALES, W. The role of lipid metabolism in production of coronary arteriosclerosis and atherosclerosis. *Am. J. Med.*, 6: 388, 1949.
72. PAUL, W. D., DAUM, K. and KEMP, C. R. Action of choline on blood lipid fractions in cirrhosis of the liver, diabetes mellitus and related conditions. *J. Iowa M. Soc.*, 37: 146, 1947.
73. FELCH, W. C. and DOTTE, L. B. Depressing effect of inositol on serum cholesterol and lipid P in diabetes. *Proc. Soc. Exper. Biol. & Med.*, 72: 376, 1949.
74. HERRMANN, G. R. Some newer approaches to the problems of arteriosclerosis. *Arq. brasil. de cardiol.*, 1: 257, 1948.

# Conference on Therapy

## Use of Curare and Curare-like Agents

THESE are stenographic reports of conferences by the members of the Department of Pharmacology and of Medicine of Cornell University Medical College and New York Hospital, with collaboration of other departments and institutions. The questions and discussions involve participation by members of the staff of the college and hospital, students and visitors. A selected group of these conferences is published in an annual volume, Cornell Conferences on Therapy, by the Macmillan Company.

DR. GEORGE READER: This therapy conference will be on the use of curare and curare-like agents. Dr. Wescoe and Dr. Riker, who have studied the pharmacology of these drugs, are here. Dr. Wescoe will open the discussion.

DR. W. CLARKE WESCOE: Curare, as you all know, is a very old drug with a fascinating history. Its medical usage has largely been confined to the last sixteen years. Prior to that time only understandardized crude preparations of curare were available and the results obtainable from its use were equally crude and unpredictable. In 1935 the active principle, *d*-tubocurarine chloride, was isolated by King. Later, in 1943, Wintersteiner obtained the same alkaloid from another species of plant.

This active principle, *d*-tubocurarine chloride, is the drug most commonly used today to produce muscular relaxation or paralysis. It is important to note that it still must be biologically assayed to determine its potency. The official assay method of the United States Pharmacopeia is the rabbit head drop test. Essentially this is a test for failure of neuromuscular transmission. A series of injections of curare is given fifteen seconds apart. The end point of the assay is the inability of the rabbit to raise its head in response to a physiologic stimulus. By comparison with the activity of a standardized preparation of crystalline *d*-tubocurarine the potency of any curare sample may be measured.

The mechanism of action of this compound merits some discussion. When an impulse travels down a motor nerve, it causes depolarization of the neuromuscular junction. This wave of negativity gives rise to the so-called action potential recorded by means of various electrical measuring devices. A rapidly falling potential results within the end plate which, when it reaches a certain critical level, excites the muscle to respond with a contraction. The membrane

of the end plate is rapidly repolarized in preparation for the next stimulus. Now, how does curare interfere with this process? It has been shown experimentally, chiefly by Eccles and his collaborators, that curare prevents the depolarization of the end plate, thus inhibiting development of the change in potential. If sufficient curare is given, the potential difference does not reach the critical level necessary to produce a muscle contraction. In essence, the motor impulse arrives normally at the end plate but its transmission to the muscle is blocked.

Since *d*-tubocurarine was first introduced, some compounds with similar actions have come into use. By adding two methyl groups to the very complicated structure of *d*-tubocurarine, it was possible to produce the dimethyl-*d*-tubocurarine which is sold commercially as metubine.<sup>®</sup> This compound has the same mechanism of action as curare but it is approximately ten times more potent. Whether or not it has any advantages I will leave to the clinicians to decide.

There is another agent which has been used in other countries and is now being introduced into America. This compound, which has the trade name flaxedil,<sup>®</sup> is quite different chemically from *d*-tubocurarine. It has a benzene structure, substituted with choline at three adjacent carbon atoms. In the choline nuclei ethyl groups are substituted for the usual methyl radicals. There is fairly good evidence that flaxedil, despite its chemical dissimilarity, has the same mechanism of action as *d*-tubocurarine.

There is still another group of drugs which produce the end results of curarization and which are sometimes called curarizing agents. Of course, a patient can be "curarized" only with curare; when paralysis is produced with another drug, it is incorrect to say the patient is "curarized." The terms "curare-like" cura-



riform" may be more properly applied to those agents which lead to an effect similar to that of curare.

The other agents to which I refer are the quaternary ammonium compounds which produce the same effects as curare but by a different means. The first of these compounds was developed in England by Ing, who combined two quaternary ammonium molecules separated by a straight chain of ten carbon atoms. It was named decamethonium bromide by Ing and is marketed under the trade name of syncurine.<sup>®</sup> The end result following administration of syncurine is the same as that produced by curare but the mechanisms of action of the two agents seem to be directly opposed. Syncurine prolongs depolarization at the end plate whereas curare prevents it. When the depolarization is prolonged, the end plate cannot repolarize to the normal resting state and the next nerve impulse cannot be transmitted to the muscle. Its action resembles that of acetylcholine which, if allowed to pile up at the myoneural junction, blocks the muscle response. Since syncurine acts in this manner it should theoretically be counteracted by *d*-tubocurarine. In the laboratory such is the case. The one paralyzing agent can be antagonized by the other if the experimental conditions are exactly adjusted. Actually many trials may be necessary to demonstrate the antagonism.

I do not know whether the prolonged depolarization produced by syncurine is desirable. On the whole, the mechanism of action of curare appears preferable although proof of harm from syncurine is lacking. It is known, however, that prolonged depolarization produced by other agents extended over a period of several days does lead to permanent damage.

There are a few other pharmacologic actions of these paralytic drugs which deserve mention. Curare has a histamine-like action if injected intra-arterially and, according to reports, when given intravenously. This may be manifested by a sudden drop in blood pressure, bronchiolar constriction and excessive secretion in the respiratory tract. Fortunately, such reactions have been encountered very rarely in the clinical usage of the drug. Curare and also syncurine produce a moderate degree of ganglionic blockade. I suppose there are some surgical circumstances in which this ganglionic blockade would not be desirable and use of such drugs might add another hazard to the procedure. Flaxedil does not alter ganglionic function. All

these agents at least partially inhibit vagal nerve function. Flaxedil in particular is almost as effective as atropine in blocking the cardiac effects of vagal nerve stimulation so that a moderate tachycardia often follows its administration.

Finally, I would like to speak about the problems involved in the use of curare and similar drugs. The greatest danger inherent in all these compounds is, of course, respiratory embarrassment. This may occur following administration of any of them and will continue to represent a hazard until a drug is developed that will spare the diaphragm while paralyzing all other skeletal muscles. Unfortunately, there seems to be nothing specific about the end plates of the diaphragm and I doubt that any such drug will be produced. There is no real difference among these compounds in their margin of safety since the occurrence of respiratory difficulties is related to production of the desired therapeutic paralysis. Therefore, I would say without reservation that these paralytic drugs should never be used unless means are available for instituting positive pressure artificial respiration at a moment's notice.

Artificial respiration must be considered the best physiologic antidote to curare and similar drugs. I think it important to know, however, that there are also available some chemical antagonists. The one most commonly used is neostigmine, also known commercially as prostigmine.<sup>®</sup> Recently a new series of compounds related to neostigmine has been developed at the New York Hospital-Cornell Medical Center. These have not yet been marketed and have no trade names. The one most practical for clinical use is referred to by its code designation, Ro 2-2561. These chemicals depolarize the end plate of the nerve, an action opposite to that of *d*-tubocurarine, metubine and flaxedil. Since the actions are opposite, they neutralize each other. The mechanism may be most simply explained by stating that Ro 2-2561 competes with curare for the neuroeffector cell. This does not occur in the case of syncurine, which produces paralysis by depolarization. Neostigmine, Ro 2-2561, and the others of the series will not antagonize the effects of syncurine but on the contrary will increase the paralysis. I believe that the only effective source of relief for cases of respiratory difficulty following use of syncurine is artificial respiration.

DR. READER: Dr. Artusio, as an anesthetist, would you care to comment?



DR. JOSEPH F. ARTUSIO, JR.: As you all realize, the primary aim in anesthetization of individuals is the production of unconsciousness but in many surgical procedures muscular relaxation is also mandatory. Until the introduction of curare, adequate relaxation could be produced only by very deep anesthesia which was associated with disturbances of many of the physiologic mechanisms which compensated for trauma, blood loss and the like. Naturally, individuals in such a state easily went into shock when exposed even to minor stress. In present day anesthesia we can maintain very light planes of anesthesia and by the use of curare or a similar drug retain the peripheral compensating mechanisms, with adequate relaxation. This has been a tremendous boon, especially for patients considered as poor risks. Today there is hardly an abdominal operation or an endotracheal intubation performed in this institution without the aid of one of the curare preparations.

We most commonly use *d*-tubocurarine. This is an extremely effective muscular relaxant and can be employed in combination with any of the anesthetic agents. When used in association with pentothal sodium or cyclopropane, the dose of curare necessary to paralyze the individual without producing severe respiratory depression is usually between 9 and 12 mg. When used in conjunction with ether anesthesia the dose necessary to produce the same effect is in the neighborhood of 3 to 6 mg. Thus with ether only about one-third as much curare is needed as with the other anesthetics.

I must emphasize that the doses mentioned are only approximations. The paralytic effects of *d*-tubocurarine are not exactly predictable even on a dose-for-weight basis. I agree emphatically with Dr. Wescoe's statement that no one should administer curare unless he is prepared to furnish artificial respiration to the patient immediately. The antagonists to curare and flaxedil provide another means of combating the effects of overdosage, but artificial ventilation is still the treatment of choice.

Our experience with flaxedil has been limited. From Dr. Marbury's work it would appear that the usual dose in man lies between 1 and 1½ mg. per kg. This produces good muscular relaxation comparable to that produced by *d*-tubocurarine, without interfering with respiration. Flaxedil has an advantage over curare in that its effects are much more predictable. We have seen the vagal blocking action of

flaxedil in patients lead to a moderate tachycardia but we have not yet decided whether this is important. Both flaxedil and curare produce their maximum effects within three minutes after intravenous injection and the effect lasts about twenty to forty-minutes.

Syncurine, the decamethonium compound, is even less predictable than *d*-tubocurarine. The dose is approximately 2 to 4 mg. in anesthetized humans and it makes little difference what anesthetic agent is used in association with it. There is not the distinction between ether and cyclopropane which must be considered when using curare.

All of these agents show cumulation if doses are repeated in less than sixty minutes. To produce the same effect as the initial dose, subsequent doses must be approximately halved.

We might add that flaxedil may eventually become the drug of choice since it does not have the ganglionic blocking action of *d*-tubocurarine. This is an advantage in that it preserves more of the peripheral compensating mechanisms to acute blood loss and trauma. Early observations, subject to confirmation, indicate that flaxedil has a second advantage over curare in that its effects are more predictable.

DR. READER: Are there any questions at this point?

DR. WALTER F. RIKER: I would like to ask Dr. Artusio, how complete is the ganglionic blockade with *d*-tubocurarine during the usual surgical procedure?

DR. ARTUSIO: That is difficult to say. I do not know any way to determine clinically the amount of ganglionic blockade. In accord with the work in the experimental laboratory, I would certainly choose a drug that did not produce a ganglionic blockade for use in those patients who may suffer trauma or acute blood loss.

DR. JOHN MCLEAN: Dr. Artusio has commented on the use of these agents in conjunction with intravenous and inhalation anesthesia. I wonder if he would say something about their use in conjunction with local anesthetics in the head region.

DR. ARTUSIO: In conjunction with local anesthesia, 3 to 6 mg. of *d*-tubocurarine will usually provide fair relaxation of the small muscles of the face with but little interference with the muscles of respiration. But again I must stress that the degree of paralysis produced by curare is not always what one would expect. Even

small doses of the drug if given to sensitive individuals may produce respiratory depression of a severe degree. I would never use it unless I was ready to offer the patient artificial respiration.

DR. McKEEN CATTELL: In relation to the fact that the compounds of the decamethonium type cause depolarization of the end plate, I wonder whether there is a discharge from the end plate resulting in muscular fibrillation, as is the case with neostigmine.

DR. WESCOE: I do not think it is seen clinically. Does Dr. Artusio see it?

DR. ARTUSIO: It is not seen clinically.

DR. WESCOE: Experience in the laboratory indicates that with any agents that block by depolarization it is possible to find some dose at which stimulation occurs.

DR. READER: In a patient in whom the effect of curare, or one of these curare-like agents is excessive, what is the first thing to do? Do you immediately try one of the chemical antagonists or do you first start artificial respiration?

DR. ARTUSIO: There are two emergency situations that can arise after curare has been given. First, the patient may have received too much curare either because of overdosage or of unusual sensitivity to the drug. Second, the patient may have been fully curarized when the operation was for some reason suddenly terminated, thus necessitating supportive measures until the paralytic effects disappear. If there is overdosage, the prime requisite is artificial respiration. The period during which this will be needed can be shortened markedly by the chemical antagonists, but I do not think a patient should be allowed to remain in apnea while someone is running to get an anti-curare compound or is drawing the compound into a syringe. If oxygen is available, it should be given under positive pressure. Once artificial respiration under optimal conditions is established the antagonist may be given at leisure. In the second circumstance when a patient has been curarized and his respiration reduced to barely adequate levels just as the operation must be terminated, the antagonists are effective. A single injection may overcome the paralysis and render breathing adequate.

DR. JANET TRAVELL: Do you ever have to give more than one dose?

DR. ARTUSIO: Yes, if the first is inadequate. The dose of an antagonist depends upon the degree of curarization, which is best measured by the respiration. We worked out the doses

required in man for one antagonist, Ro 2-2561, using the respiratory minute volume as a guide. For a completely paralyzing dose of *d*-tubocurarine 15 mg. of Ro 2-2561 were needed. When respiration was reduced to two-thirds, only 10 mg. of the antidote were needed. Five milligrams counteracted a respiratory depression of 30 to 50 per cent. For these trials the intravenous route of administration was employed.

The maximum effect of the antagonist is obtained within five minutes. If the dose is insufficient to antagonize all the curare, some paralysis will again become apparent. Then it may be necessary to give another dose of the antagonist, the dose to be calculated again on the basis of the degree of respiratory depression which remains.

DR. WESCOE: I do not think that Dr. Artusio is including neostigmine in his present discussion. The dose for neostigmine is much smaller than the other antagonists.

DR. ARTUSIO: I did not mean to include neostigmine in that dose schedule.

DR. READER: How do you tell when a patient has had too much curare? Are you immediately aware of it?

DR. ARTUSIO: Just observe the respiration, preferably by actual measurement of minute volume. The maximum effect of curare will appear within three minutes after its intravenous injection.

DR. READER: Do curare and the other drugs stimulate the central nervous system?

DR. WESCOE: Not within the dose ranges that are used clinically. All the effects are peripheral.

DR. READER: You stated that these agents affect the motor end plate of the nerve. Is there any change in the muscle cells themselves?

DR. WESCOE: The muscle remains perfectly normal. Even though paralyzed completely, so far as response to nerve stimuli is concerned, the muscle will react to direct electrical stimulation.

DR. DAVID B. THOMPSON: How are curare and other compounds handled by the body? Are there any disease states in which their use is contraindicated?

DR. READER: Dr. Riker, would you like to answer the first part of the question?

DR. RIKER: There is no information on the metabolism of *d*-tubocurarine by the animal organism. We do not know what happens to it after it leaves the myoneural junction. There is some not entirely conclusive evidence that it



may be eliminated in the urine. Further study of these aspects of the subject is badly needed.

DR. READER: Are there any chemical tests for curare?

DR. RIKER: No, their unavailability is at present a real handicap.

DR. THOMPSON: What are the contraindications to the curarizing drugs?

DR. READER: Dr. Artusio, do you have an opinion about that?

DR. ARTUSIO: The only disease which would make me use curare with caution is myasthenia gravis. There certainly is a possibility that administration of the drug to an unrecognized myasthenic might result in paralysis more profound than was anticipated. I cannot think of any other disease that might cause difficulty.

INTERN: How would you handle an obvious case of myasthenia being maintained on prostigmine, which is an antagonist to curare?

DR. RIKER: Just discontinue the prostigmine. These patients may come to you practically paralyzed.

DR. WESCOE: I would not even say that myasthenia is a contraindication to the use of curare. In my opinion that belief, while traditional, has little to support it. Curare can be given to a myasthenic but must be administered judiciously. Even if difficulty arises, nothing serious is going to happen so long as artificial respiration is maintained.

DR. READER: Would the antagonist of choice for a myasthenic be neostigmine?

DR. WESCOE: No. The antagonist of choice for the curare agents is one of the shorter acting compounds. Neostigmine in too large doses is very likely to cause trouble because it possesses a curariform action like that of syncurine. It is difficult to predict whether the stimulating or depressant effects will predominate. It is quite possible to augment the paralysis of curare by adding to it the paralytic effects of too much prostigmine. This cannot happen with the briefer acting newer antagonists.

DR. READER: Dr. Dunning, how do you feel about the question of contraindications? Are there any neurologic conditions which might increase the dangers of these paralytic drugs?

DR. HENRY S. DUNNING: I should think a patient who has had poliomyelitis with some residual weakness of his respiratory muscles would run greater risks of respiratory difficulties than would a normal individual.

DR. READER: How about amyotrophic lateral sclerosis?

DR. DUNNING: I would include that in the same category. We often see cases of amyotrophic lateral sclerosis, poliomyelitis and even neuritis in which the intercostal muscles are not working. This is often missed in an ordinary physical examination. I should think such a condition might increase the dangers. Surely the patient's respiratory apparatus should be very carefully examined before curare is given.

DR. READER: I was interested in your remark about poliomyelitis because there was once much agitation for the use of curare in acute poliomyelitis. Dr. Plum, could you give us a brief description of the use of curare in poliomyelitis? What was the rationale of this treatment and why was it abandoned?

DR. FRED PLUM: I never used it so I can only report the experience of others. My understanding is that curare was used in an attempt to break up joint limitation which otherwise occurred in individuals with painful and stiff muscles. In other words, the normal motion about a joint was limited because of pain and stiffness in adjacent muscles. By curarizing those muscles it was hoped to achieve earlier mobility and therefore earlier physical re-education. I believe the reason this therapy was given up was simply that long term studies showed that the patients who were curarized did not get well any faster. Indeed, histologic studies on muscles from those patients showed a good deal of actual rupture of the muscle fibers about the joints. Some observers believed that groups of paralyzed muscles were being overstretched and damaged. I have some difficulty visualizing how this could happen if they maintained muscles within the usual states of stress. Nevertheless, even though the curare was not detrimental, it accomplished no demonstrable benefit.

DR. READER: Curare was also popular at one time for patients with arthritis, based on the same reasoning Dr. Plum has described. Dr. Wescoe, was there any value in this therapy?

DR. WESCOE: I do not know what the status of therapy is there. Dr. Travell might have something to add.

DR. TRAVELL: There are very few people who use it. To do any good in a chronic disease like arthritis curare would have to be given on a long term, ambulatory basis. It is not possible to do



this safely. There are treatments more effective and more easily used.

DR. CATTELL: Are we trying to keep the chemical characteristics of these antagonists under a wrapper? It might be interesting to hear about their relationship to acetylcholine and the other members of that group.

DR. WESCOE: These compounds were originally developed because it was noted that neostigmine had several actions, one of which was a potent anti-cholinesterase action. Previously it had been shown by Stedman that much of the anti-cholinesterase action of neostigmine is due to the dimethylcarbamate grouping on its basic trimethylphenylammonium structure. We first requested the chemist to replace the carbamic acid end of the molecule with an acetyl radical, to give a compound closely related structurally to acetylcholine. Other compounds studied included one with a hydroxyl substitution of the carbamate grouping and also the basic trimethylphenylammonium salt. The anti-curare action of the three modifications of the neostigmine molecule are essentially similar.

INTERN: Originally you stated that the anti-curare compounds cause depolarization of the end plate of the nerve. That is also the way in which you said that syncurine paralyzes. Why then don't these antagonists block neuromuscular transmission?

DR. WESCOE: In the laboratory they do. Small doses of the antagonists cause only stimulation with marked muscle fasciculations. If larger doses are given, this brief period of stimulation is followed with a curare-like paralysis. But this paralysis is of brief duration, measured in minutes, and most importantly it has been impossible in the laboratory to show any summation between the paralytic actions of curare and the antagonists. If a patient manifests effects of curare, the antagonists do not add to the paralysis but instead counteract it. Within six minutes the actions of the antagonists are over and, if curare effects have returned, an additional dose can be given without any fear of adding to the paralysis.

DR. CATTELL: Is the antagonist entirely eliminated in a matter of minutes?

DR. WESCOE: That is a difficult question. The effects certainly disappear rapidly but we do not know the fate of the compound itself. We do not know how it is eliminated.

DR. READER: Some time ago I heard much discussion of a long-lasting curare-like com-

pound. Wasn't some new drug developed to provide a prolonged effect?

DR. WESCOE: That was probably mephenesin which can be given over relatively long periods to promote muscular relaxation. However, the paralysis is mediated by an action on the reflex center of the brain and spinal cord and thus does not resemble curare in its mechanism of action. No drug with a true curariform action and a prolonged period of action has been developed. The Romansky formula developed for repository penicillin has been used to provide slowly absorbed deposits of *d*-tubo-curarine. I do not know of anyone still using it.

DR. READER: Wasn't that quite dangerous?

DR. WESCOE: The unpredictability of absorption of curare was added to the unpredictable paralytic effects of the drug.

DR. TRAVELL: The total amount given in slowly absorbed preparations would have to be much greater than the usual single therapeutic dose. That might be a real hazard!

DR. WESCOE: I think that is true.

DR. LEON J. WARSHAW: I wonder whether anyone has used curariform drugs by intra-arterial injection when relaxation of the muscles of a limb is desired for the reduction of fracture.

DR. READER: Does anyone know about that? Dr. Wescoe or Dr. Artusio, have you ever heard of its use in such circumstances?

DR. ARTUSIO: No.

DR. WESCOE: The only intra-arterial studies are experimental.

DR. WARSHAW: Is it possible that by giving a small dose into an artery and blocking the venous return from the area, a local effect might be obtained without any systemic or respiratory changes?

DR. READER: That certainly sounds like a practical idea. Are there any other questions?

DR. RIKER: I would like to ask Dr. Artusio if he has noted any histamine-like or ganglionic blocking actions of *d*-tubocurarine during surgical procedures.

DR. ARTUSIO: We usually give curare intravenously and by this route its histaminic actions rarely occur. It is difficult to determine by clinical means the presence of ganglionic blockade. We have not been impressed by its occurrence during operations. Since, however, we know that it may happen, I think it is a factor to be considered before administering curare to patients who may suffer much trauma or blood loss during operation.

DR. CATTELL: I would like to ask whether Dr. Artusio always intubates the patient before using these drugs.

DR. ARTUSIO: We never give curare without prior intubation so that we have a ready pathway for artificial ventilation. That is the safest method to employ and the safest rule to make in a large service such as we have here.

DR. CATTELL: Would you say curare should never be used without intubating first?

DR. ARTUSIO: It should not be used unless the proper apparatus is at hand and the individual who is administering the curare can produce unobstructed artificial ventilation.

DR. READER: That brings up another point, however. Do you ever use curare in order to intubate someone?

DR. ARTUSIO: Oh yes, we do that very often but of course the intubation is accomplished immediately and that is the purpose in giving the curare, to produce an airway. I have not seen any individual who had had curare whom it was not possible to intubate.

DR. READER: In a patient with laryngeal spasm, or some other condition for which tracheotomy has heretofore been indicated, is curare plus intubation better than tracheotomy?

DR. ARTUSIO: I believe that an individual in severe laryngospasm should first be treated with positive pressure oxygen by means of a mask and bag in an attempt to break the spasm. Anesthetists usually have atropine available in their equipment. A large dose of atropine, 0.8 to 1 mg. intravenously, may be of great help in abolishing the laryngospasm. If curare is immediately available, and I mean immediately because laryngospasm cannot go on long without producing central nervous system damage, it may be used. However, if I could not break the laryngospasm with positive pressure oxygen, I would have the surgeon begin tracheotomy while intubation was being attempted.

DR. READER: Do you use curare with spinal anesthesia?

DR. ARTUSIO: It may have some value in spinal anesthesia if the muscular paralysis has worn off while the sensory paralysis is still intact. A small dose may give satisfactory relaxation for a while longer. We have not used curare in that fashion. If in spinal anesthesia the relaxation is poor, we convert to a general inhalation anesthesia.

DR. READER: Dr. Victor Marshall told me that he finds curare essential in doing prostatectomies

by the transurethral route because it combats priapism. Were those patients receiving spinal anesthesia?

DR. ARTUSIO: Some of them were under spinal anesthesia. We used small doses of curare, 3 to 6 mg., in the conscious patient, which diminished the priapism. We were, of course, always ready to intubate the patient.

DR. READER: Dr. Wescoe, do you have a comment?

DR. WESCOE: I question giving curare with a local anesthetic because it is my impression that the surgeon does not always take the patient seriously when he says the sensory anesthesia is diminishing. It would be an unpleasant experience not to be able to respond to a painful stimulus.

DR. ARTUSIO: We have given curare to a few conscious individuals with severe muscle spasm. They had low back muscle spasms which elicited terrific reflex pains whenever they moved. We gave them 3 to 6 mg. of *d*-tubocurarine intravenously, got them out of bed and walked them around. Much to my surprise the spasm was completely abolished and did not return.

DR. READER: Were they able to walk unassisted?

DR. ARTUSIO: Yes. You must remember that the condition of the patient partially determines the response to curare. Earlier I mentioned the fact that ether sensitizes the myoneural junction to curare so that the full curarizing dose in an etherized patient would be about one-third the dose for a patient during cyclopropane anesthesia. Without any anesthesia the dose would be slightly higher. In a conscious, unanesthetized human, 3 to 6 mg. of curare will produce no respiratory difficulties and only slight muscular weakness.

#### SUMMARY

DR. READER: Dr. Wescoe has discussed the pharmacology of *d*-tubocurarine and flaxedil which are believed to produce muscle relaxation by preventing depolarization of the nerve end plate. Syncurine, on the other hand, leads to the same result by prolonging depolarization. This distinction is important in the use of the anti-curare agents since the latter do not counteract the effects of syncurine. There is general agreement that artificial respiration is the best treatment for respiratory difficulties arising after administration of any of these neuromuscular blocking agents. Indeed, these drugs should not

ever be given unless facilities for proper ventilation are immediately available. However, with the exception of syncurine, the chemical antagonists may be useful in that they can immediately, completely and safely antagonize the effects of curare. It seems to be the opinion that flaxedil may prove superior to *d*-tubocurarine in that it does not alter ganglionic

transmission and its effects are more uniform than those of curare. The point has been made that curare and the other drugs are quite unpredictable in their actions. However, if this characteristic is taken into account, and if the means are available to produce artificial respiration, there are few contraindications to the use of these drugs.



# Clinico-pathologic Conference

## Miliary Pulmonary Infiltration with Hydrothorax and Ascites

STENOGRAPHIC reports, edited by Robert J. Glaser, M.D. and David E. Smith, M.D., of weekly clinico-pathologic conferences held in the Barnes Hospital, are published in each issue of the Journal. These conferences are participated in jointly by members of the Department of Internal Medicine and Pathology of the Washington University School of Medicine and by Junior and Senior medical students.

THE patient, E. H. (No. 188544), a sixty-five year old white married housewife, entered the Barnes Hospital on August 31, 1950, complaining of shortness of breath, swelling of the abdomen and fatigue. Because of the patient's condition the history was obtained with difficulty and was not considered adequate.

The family history was of interest in that one brother died of "lung disease" of unknown etiology. As far as could be determined the patient herself had enjoyed generally good health. There was no known previous serious illness; and although her appetite had been poor for years and her diet had been lacking in protein, an occasional non-productive cough had been her only known symptom. She had kept a dog in her home for many years.

One year before admission the patient developed uterine prolapse and a complicating urinary infection; she was admitted to a local hospital where a hysterectomy was performed. The general physical examination was said to have been negative at that time. Following the operation, however, the patient failed to regain full strength and she noted the onset of dyspnea on exertion. Six months prior to entry she developed a "severe cold" associated with a productive cough. These symptoms persisted and after three weeks the patient sought medical attention. Her physician told her that she had a high fever and made a diagnosis of virus pneumonia. She was treated with penicillin and a sulfonamide with some transient improvement, but she then developed increasing fatigue, dyspnea, anorexia and cough, and she lost weight.

Three months before admission to the Barnes Hospital, she was readmitted to the local hospital where she remained for ten days. At that

time her temperature was 97.4°F. and physical examination revealed pallor, malnutrition, signs of pleural fluid in the left lower chest, ascites, an enlarged, firm, non-tender liver and ankle edema. The laboratory findings included normal white cell and differential counts. The non-protein nitrogen was 37 mg. per cent and the cephalin-cholesterol flocculation test 4 plus. An electrocardiogram was said to have been "abnormal" and a chest film revealed cardiac enlargement and a left pleural effusion. A gastrointestinal x-ray series was negative, but cholecystograms revealed non-function of the gallbladder.

A thoracentesis was performed; smears of the fluid showed many intracellular organisms which were thought to be histoplasma capsulatum. In addition, there were cigar-shaped forms, thread-like structures resembling mycelia, and some acid-fast organisms which were "not characteristic" of tubercle bacilli. Cultures of the pleural fluid on Sabaraud's medium and on blood agar were negative. During her hospital stay the patient's temperature ranged between 99° and 100°F.

After discharge she experienced progressive swelling of the abdomen and legs and was once again admitted to the local hospital. Physical examination revealed that she was moderately dyspneic. Signs of fluid were present over both lung bases and the heart was moderately enlarged. The abdomen was distended with fluid and there was pitting edema of the legs. A chest film showed cardiac enlargement and bilateral pleural effusions, with the most marked effusion on the left. Thoracentesis and abdominal paracentesis were performed, but examination of the fluids failed to show the organisms previously described.

During her first hospital week the patient's

temperature ranged between 101° and 101.5°F. but subsequently it became almost normal. She was given small doses of digitalis and a transfusion and was discharged. After leaving the hospital she once again had increasing abdominal swelling and edema of the legs, and she was referred to the Barnes Hospital.

At the time of entry physical examination revealed the temperature to be 36.8°C., pulse 90, respirations 20 and blood pressure 155/90. The patient was thin, weak and appeared chronically ill. Her complexion was sallow. Examination of the skin revealed it to be of poor turgor, and there were several decubitus ulcers over her back. The retinæ showed moderate arteriosclerotic changes but no hemorrhages or exudates. Examination of the upper respiratory tract was negative except that the tongue was extremely red. Examination of the chest revealed flatness, absent fremitus, and diminished breath sounds over the entire right chest. The heart presented no abnormalities. The abdomen was moderately distended with fluid, but the liver could be felt 4 cm. below the right costal margin. The feet and legs were markedly edematous.

The laboratory data were as follows: Blood count: red cells, 4,910,000; hemoglobin, 16.3 gm; white cells, 9,100; differential count: stab forms, 13 per cent, segmented forms, 78 per cent; lymphocytes, 3 per cent; monocytes, 6 per cent. Urinalysis: specific gravity, 1.019; albumin, trace; sugar, negative; bilirubin, 1.2 mg. per cent; urobilinogen, 0.49 mg. per cent; centrifuged sediment: many clumped white blood cells and many bacteria. Urine culture: *B. aerogenes*. Stool examination: guaiac negative. Blood Kahn test: negative. Blood chemistry: Non-protein nitrogen, 18 mg. per cent; sugar, 79 mg. per cent; chlorides, 96 mEq./L.; total proteins, 6.9 mg. per cent; albumin, 2.6 gm. per cent; globulin, 4.3 gm. per cent; alkaline phosphatase, 4 Bodansky units; cephalin-cholesterol flocculation test, 3 plus; thymol turbidity, 9.1 units; total cholesterol, 91 mg. per cent; cholesterol esters, 50 mg. per cent; BSP retention: 60 per cent in thirty minutes; sodium bilirubinate, 0.44 mg. per cent; bilirubin, 1.64 mg. per cent. Venous pressure: 85 mm of water. Circulation time (decholin): fifteen seconds. Roentgenogram of the chest: there was distinct cardiac enlargement, chiefly of the left ventricle; a pleural effusion was present at the right base and there

was pulmonary congestion. In addition a fine mottling was seen throughout both lung fields. Electrocardiogram: slightly depressed S-T segments in leads II, III and AVF; inverted T waves in V<sub>4</sub>, V<sub>5</sub> and V<sub>6</sub>; flat T waves in leads I, II, III and AVF.

On the day following admission a right thoracentesis was performed and 1,600 cc. of pleural fluid were removed. The fluid had a specific gravity of 1.015. Sections made from a cell block revealed no malignant cells and gave no clue to etiology. An abdominal paracentesis was done and 700 cc. of cloudy yellow fluid was removed. This fluid had a specific gravity of 1.011 with a total protein of 0.88 gm. per cent. The cell count was 500 of which 50 per cent were segmented and 50 per cent mononuclear forms. No organisms were seen. Cell block section of the ascitic fluid revealed mycelia and spores, but these were considered to be contaminants. Sternal bone marrow aspiration was carried out. Smears of the marrow were negative for organisms, but on culture *Histoplasma capsulatum* was recovered two days before the patient's death. Many examinations of pleural fluid obtained by repeated thoracenteses were made but neither organisms nor tumor cells were seen. Histoplasmin and first strength PPD skin tests were negative, and the histoplasma complement fixation test was negative. Microscopic sections of cell blocks made from pleural fluid, in which organisms were originally described, were reviewed by a mycologist who suggested the possibility of sporotrichium infection. Because of the possibility of sporotrichosis potassium iodide was given in increasing doses without favorable effect. The patient became weaker and continued to be anorectic and lethargic. Edema of the legs became more marked, and the patient was given intravenous salt-free albumin. Because of poor food intake she was maintained by intravenous alimentation and vitamin supplements. Despite the latter substances, however, she developed acute glossitis and stomatitis. At the end of her second hospital week the following laboratory data were obtained: prothrombin time, 70 per cent of normal; total proteins, 7.0 gm. per cent; albumin, 2.4 gm. per cent; globulin, 4.6 gm. per cent; non-protein nitrogen, 12 mg. per cent; chlorides, 88 mEq./L. Red blood cell count, normal; hemoglobin, normal; white blood cell count, 13,350; differential, normal. Subsequent roentgenograms of

the chest again revealed bilateral miliary mottling in addition to the pleural effusions.

Three weeks after admission another paracentesis was performed. The fluid had the characteristics of a transudate and no organisms were seen on smear. The non-protein nitrogen rose to 32 mg. per cent and then to 61 mg. per cent. The patient had increasing dyspnea, she vomited frequently and adequate nutrition could not be maintained. Weakness, lethargy, anorexia, hypotension and dehydration were prominent. The cephalin-cholesterol flocculation test was 4 plus, and the thymol turbidity 5.3 units. Repeated electrocardiograms showed no evidence of hyperkalemia although the T waves were upright. During her hospital stay the patient exhibited a swinging type of low grade fever, but in the last few days it rose to between 38° and 39°C. She died on September 28, 1950, shortly after developing signs of bronchopneumonia.

#### CLINICAL DISCUSSION

DR. HARRY L. ALEXANDER: This case presents a most interesting problem in differential diagnosis. During the period that this patient was in another hospital in this community a diagnosis of histoplasmosis was made. Subsequently when she was admitted here, bone marrow examination was performed; and although examination of smears of the marrow failed to reveal *Histoplasma capsulatum*, culture of the marrow was reported positive for these organisms two days before the patient died. Dr. Elliott, would you begin the discussion by commenting on the chest films, particularly the presence of miliary mottling in the lung fields? What is the significance of this particular finding?

DR. GLADDEN V. ELLIOTT: The miliary infiltrate which was noted in this patient's chest roentgenogram was compatible with several etiologic factors. Such lesions are sometimes due to pulmonary congestion occurring in acute heart failure, and they may occur on the basis of hemosiderosis following long-standing heart failure. Excluding these two explanations one must certainly consider either histoplasmosis or miliary tuberculosis.

DR. ALEXANDER: In an attempt to substantiate the diagnosis of histoplasmosis, skin tests were performed and serum was collected for a complement fixation test. As was noted in the protocol the skin test was negative, and the complement fixation test, which was done by

Dr. Furcolow of the United States Public Health Service, was also negative. Dr. Wells has been particularly interested in this disease, and I have asked him to review the significance of these particular data.

DR. JOHN G. WELLS: The significance of the histoplasmin skin test and of the complement fixation test in substantiating the diagnosis of histoplasmosis has been discussed rather widely in the recent literature. It can be said at the outset that no definitive conclusions have as yet been reached. Several important observations have been made by Furcolow.<sup>1</sup> He noted that apparently healthy patients who had positive histoplasmin skin tests frequently lost their skin reactivity to histoplasmin during the course of various acute illnesses other than histoplasmosis. After recovery the patients again exhibited positive histoplasmin tests. The incidence of positive histoplasmin skin tests also decreased with increasing age. The experience at the Children's Hospital here is consistent with the foregoing results. Thus Dr. Klingberg<sup>2</sup> has recently summarized ten cases of histoplasmosis in which the diagnosis was proven in all by culture and in nine by subsequent pathologic examination. None of the nine patients who died, all of whom had repeated positive cultures, had a positive skin test or a positive complement fixation test. In the one child who recovered the skin and complement fixation tests were negative during the acute stage of the disease, but both became positive after the child recovered. In one of the recent Barnes Hospital cases in which the diagnosis of histoplasmosis was proven at autopsy, the skin test was negative. It seems clear, therefore, that negative skin tests and/or negative complement fixation tests in acutely ill patients do not rule out the diagnosis of histoplasmosis.

DR. ALEXANDER: Thank you for your summary, Dr. Wells. The information you have presented is pertinent in the case under discussion today. The fact that both the skin and complement fixation tests were negative in this gravely ill woman cannot be considered to militate against the diagnosis of histoplasmosis. Also of interest in this particular case was the presence of both pleural effusions and ascites. In the collected literature on the subject up to 1945, and

<sup>1</sup> FURCOLOW, M. L., EMGE, M. E. and BUNNELL, I. L. Depression of tuberculin and histoplasmin sensitivity associated with critical illness. *Pub. Health Rep.*, 63: 1290, 1948.

<sup>2</sup> KLINGBERG, W. G. Generalized histoplasmosis in infants and children. *J. Pediat.*, 36: 728, 1950.



indeed even later than that, not a single case of histoplasmosis with which ascites was associated was reported. Pleural effusion is also quite uncommon. The latter fact seems to me to be particularly unusual because the organism frequently attacks the lungs and the subpleural tissues; one would expect, therefore, that pleural effusion would be encountered in a significant per cent of cases. Dr. Wells, what have you found out about this particular point?

DR. WELLS: Pleural effusion was not observed in any of the ten cases at the Children's Hospital. One of the recent patients with histoplasmosis in this hospital had both a pleural effusion and ascites; but the concomitant association of severe hypoproteinemia and of congestive heart failure made it difficult to determine the role of the infection *per se* in the development of fluid in the two body cavities. In two other recent cases of proven histoplasmosis in children, one patient had a definite pleural effusion demonstrated by x-ray, and in the other an interlobar effusion was suspected.<sup>3</sup> Nonetheless as you have pointed out, Dr. Alexander, the accumulated clinical evidence indicates that ascites and pleural effusion are rare in histoplasmosis.

DR. ALEXANDER: You have had much experience with this disease, Dr. Goldman. Have you found pleural effusion in any of the patients you have followed?

DR. ALFRED GOLDMAN: No, I have not.

DR. ALEXANDER: Dr. Flance, would you comment on this point?

DR. I. JEROME FLANCE: I agree entirely with what has been said.

DR. ALEXANDER: It can be assumed then that if this patient indeed had histoplasmosis, she was one of the very few in whom ascites and pleural effusion have been encountered. Because of this fact the possibility that this patient's major disease was something other than histoplasmosis must be considered. Does anyone wish to suggest an alternative diagnosis?

DR. W. BARRY WOOD, JR.: Since Dr. Elliott believed that the miliary mottling in the chest film was compatible with tuberculosis, that diagnosis must certainly be considered. Further, at one point during her stay at the other hospital acid-fast organisms were present in the pleural fluid.

<sup>3</sup> WHEELER, W. E., FRIEDMAN, J. and SASLAW, S. Simultaneous non-fatal systemic histoplasmosis in two cousins. *Am. J. Dis. Child.*, 79: 806, 1950.

DR. ALEXANDER: I would agree that tuberculosis certainly is a distinct possibility although the fact that the pleural and ascitic fluids both had characteristics typical of transudates is disturbing. Dr. Massie, this patient had an enlarged heart and an abnormal electrocardiogram. What importance do you ascribe to cardiac failure *per se*?

DR. EDWARD MASSIE: When she was admitted to the Barnes Hospital the patient's venous pressure and circulation time were both normal. Although she may have had mild failure, I doubt that it played a major role in the general process.

DR. ALEXANDER: Are there other suggestions?

DR. ROBERT J. GLASER: One other diagnosis which should be mentioned is that of collagen disease. The bizarre clinical picture which she presented can be explained on that basis.

DR. GOLDMAN: It is possible that this patient had a fungus infection plus another disease process such as carcinomatosis or tuberculosis. I believe the chest findings were compatible with miliary carcinomatosis as well as with tuberculosis or histoplasmosis. I also agree with Dr. Glaser that collagen disease must be considered.

DR. ALEXANDER: If this patient had had disseminated carcinomatosis, wouldn't you have expected one of numerous cell blocks which were made to have shown malignant cells?

DR. GOLDMAN: I doubt that examinations of cell blocks made from pleural and ascitic fluids of patients with carcinomatosis are positive in more than 50 per cent of cases.

DR. ALEXANDER: Isn't the diagnosis more difficult in the case of ascitic fluid than pleural fluid?

DR. GOLDMAN: There are more false positives with ascitic fluid than with pleural fluid.

DR. ALEXANDER: Dr. Wood, you have suggested that this patient had miliary tuberculosis. In my examination of the record I discovered that fifteen tuberculin skin tests were done and all were negative. Does that finding disturb you?

DR. WOOD: It does to some extent but not completely. The situation in the terminal phase of tuberculosis may be similar to that to which Dr. Wells has alluded in histoplasmosis; that is, PPD skin tests may become negative. Such a sequence of events does not always occur in tuberculosis.

DR. ALEXANDER: Dr. Goldman, should disseminated sporotrichosis be considered in the absence of skin lesions?

DR. GOLDMAN: Usually sporotrichosis, of course, does involve the skin, but it may also produce pulmonary lesions.

DR. ALEXANDER: Does disseminated sporotrichosis occur in this country without skin lesions, Dr. Moore?

DR. MORRIS MOORE: I know of only one such case.

DR. GLASER: Dr. Alexander, when you prepared this protocol you must have been impressed with the fact that the patient kept a dog in her house because you specifically mentioned it. Would you tell us the significance of that information?

DR. ALEXANDER: When I first noted the statement about the dog in the history, I was not impressed with it. Subsequently, Dr. Rosecan pointed out to me that dogs may be infected with histoplasma, and he suggested that the statement should be included.

DR. WELLS: It is of interest in that regard, Dr. Alexander, that very exhaustive studies have demonstrated *Histoplasma capsulatum* in dogs, cats, Norwegian rats, house mice, skunks and chipmunks.<sup>4</sup> It is thus widely distributed among animal life.

DR. ALEXANDER: Dr. Kenamore, when this patient was first referred to you, what was your opinion about the etiology of her disease?

DR. BRUCE D. KENAMORE: We were quite puzzled by the clinical picture and laboratory data which did not seem to us consistent with a diagnosis of tuberculosis; we of course considered histoplasmosis seriously, and we asked Dr. Moore's help in delineating the problem.

DR. ALEXANDER: Would you comment, Dr. Moore, on your interpretation of the various cultures and smears?

DR. M. MOORE: Although acid-fast organisms were said to have been found in smears of the fluid obtained at the time of the first thoracentesis, I was never able to demonstrate any. I did find some intracellular organisms which suggested *Histoplasma capsulatum* but they were very few in number and, in the absence of a positive culture at that time, I expressed doubt as to that diagnosis. In regard to the cigar-shaped cells which were described, the possibility of sporotrichosis came to mind because in tissue this fungus may produce cigar-shaped cells. Against the diagnosis of sporotrichosis, however, was

the fact that these cigar-shaped bodies stained very well. Further, the presence of very large cells is quite uncommon in sporotrichosis. I was thus unable to make a definitive diagnosis, but on the basis that sporotrichosis was a possibility, albeit an unlikely one, I suggested that iodides be given. Iodide therapy was without effect.

DR. ALEXANDER: Iodide is specific for sporotrichosis, is it not?

DR. M. MOORE: Yes, it is the one fungus infection for which iodide is specific.

DR. WELLS: We asked Dr. Henry Pinkerton of the Department of Pathology of St. Louis University Medical School to examine the slides which Dr. Moore has described to see if he thought toxoplasmosis merited consideration. Dr. Pinkerton did not believe that the cellular forms were compatible with that diagnosis.

DR. ALEXANDER: Dr. Moore, you saw the culture of the bone marrow which was reported to be positive for histoplasma two days before the patient died. Did you confirm the report?

DR. M. MOORE: Yes, the culture was unequivocally positive for *histoplasma capsulatum*.

DR. ALEXANDER: On the basis of that finding, one certainly would assume that the patient had histoplasmosis.

DR. M. MOORE: There can be no question that the bone marrow culture was positive for the organisms; whether or not the patient had histoplasmosis, however, may be another problem.

DR. ALEXANDER: Do patients carry these organisms?

DR. M. MOORE: I do not know.

DR. CARL V. MOORE: In every instance of which I am aware, when histoplasma was cultured from the bone marrow, the patient was shown to have disseminated histoplasmosis.

DR. WOOD: I believe I saw this patient in consultation with Dr. Kenamore. If I remember correctly, I thought the evidence favored histoplasmosis as the most likely diagnosis. I must admit that at that time I was unaware of the infrequency with which pleural effusion and ascites associated with histoplasmosis. Had I had that information, I might not have been as willing to make the diagnosis. After the positive culture was recovered, we felt certain that the patient had histoplasmosis.

DR. ALEXANDER: I believe that the evidence distinctly favors the diagnosis of histoplasmosis. Other possibilities include tuberculosis and sporotrichosis, but these are less likely. I doubt

<sup>4</sup> EMMONS, C. W. Animal reservoirs and other sources of pathogenic fungus, *Histoplasma capsulatum*. *Am. J. Pub. Health*, 40: 436, 1950.

that any of us would be surprised if this patient were found to have more than a single disease entity responsible for the entire clinical picture.

*Clinical Diagnosis:* Histoplasmosis.

#### PATHOLOGIC DISCUSSION

DR. MENARD C. IHNNEN: At autopsy the patient was well nourished in spite of her long illness. The pertinent external findings were scattered petechiae on the flexor surfaces of the arms and anterior surfaces of the legs and slight edema of the legs. In the thorax there were 500 cc. of bloody pleural fluid on the right and 100 cc. of clear fluid on the left. There were rather dense fibrous adhesions over the middle and lower lobes of the right lung. On the surfaces of both lungs, but more clearly seen on the left, there were increased lymphatic markings accompanied by tiny, grey, firm nodules 1 or 2 mm. in diameter. Several slightly enlarged lymph nodes at the hilum of the right lung contained calcium and white granular material. Throughout the parenchyma of the lungs, particularly in the lower lobe of the right lung, there were fairly well circumscribed 2 mm. grey nodules. In the lower lobe of the left lung there was a lesion 3 cm. in diameter that was grey-red, indistinctly outlined and slightly granular. The pericardium was severely thickened, particularly over the right lateral surface. The tissue had a granular, firm, grey-yellow appearance; it was moist and did not dissect like fibrous tissue. The heart was not enlarged and it showed no other pertinent lesion.

In the abdominal cavity there were 600 cc. of clear fluid. The mesenteric lymph nodes were not unusual nor was the peritoneum. The liver weighed 1,000 gm. and the surface was finely nodular. These nodules were grey-yellow and slightly raised; the average was no larger than 4 mm. The organ did not cut with increased resistance, and nodules similar to those on the capsular surface were present throughout all parts. The spleen was of normal size and consistency; on the cut surface, in addition to many lymphoid follicles, scattered grey soft areas that were of a slightly irregular outline and a little larger than lymphoid follicles were apparent. The surfaces of the kidneys were very finely granular. Tiny grey nodules similar to but smaller than those in the liver were scattered over the surfaces and throughout the medulla and cortex. There were linear grey foci in the pyramids. The gastrointestinal tract was of

grossly normal appearance, as were the pancreas, adrenals, bone marrow and gallbladder.

DR. GUSTAVE J. DAMMIN: The usual gross features in histoplasmosis are a striking involvement of the lymph nodes with only occasional lesions of macroscopic proportions in the viscera. The clinical diagnosis of histoplasmosis seemed acceptable in this case, however, as parenchymal lesions are known to occur with a resulting picture of a disseminated granulomatous process similar to some other fungal diseases and tuberculosis. A particularly illustrative case has been described by Dublin, Culbertson, and Friedman.<sup>5</sup> With the clinical diagnosis of histoplasmosis confirmed by culture and knowledge of the possibility that gross lesions of that type might occur, very little else was considered in this case until the microscopic sections were studied.

In Figure 1 one of the small nodules described in the lungs is illustrated. The lesion is a typical tuberculoid focus with some necrosis in the center and a granulomatous reaction with two giant cells at the periphery. The large lesion in the lower lobe of the left lung had the microscopic appearance illustrated in Figure 2. There is total filling of the alveoli with cells and an exudate that has undergone caseous necrosis with consequent obliteration of the outlines of the alveolar walls. A section of the greatly thickened pericardium (Fig. 3) shows giant cells and the same type of granulomatous reaction with some fibrosis. An identical granulomatous thickening involved the visceral and parietal layers of both the pericardium and right pleura. The lymph nodes contained a great deal of necrosis toward their centers but a distinct granulomatous reaction at the periphery. Similar lesions were present in the spleen, and Figures 4 and 5 show the miliary granulomas in the liver and bone marrow. The cellular elements in the uninvolved areas of bone marrow are not remarkable. The particular granuloma illustrated happens to have more necrosis than some; others had more fibrous reaction. The nodules presented a continuous series representing various stages between necrosis and fibrosis. With the usual stains they were compatible with and were first accepted as lesions of histoplasmosis. In sections of the adrenal (Fig. 6), although no lesions were observed grossly, there are a few foci which have the appearance of those noted in the organs already described. The lesions

<sup>5</sup> DUBLIN, W. B., CULBERTSON, C. G. and FRIEDMAN, H. P. Histoplasmosis. *Am. Rev. Tuberc.*, 58: 562, 1948.






FIG. 1. A nodule of miliary tuberculosis in the lung; in this and all other lesions acid-fast bacilli were demonstrated; no fungi were seen despite the positive culture of *Histoplasma capsulatum* during the patient's life.

FIG. 2. A focus of tuberculous pneumonia in the lower lobe of the left lung.

FIG. 3. Tuberculous pericarditis; the pleura of the right side had a similar microscopic appearance.

FIG. 4. Miliary tubercle in the liver.

FIG. 5. Miliary tubercle with central necrosis in the bone marrow; the remaining marrow around this and similar lesions was not remarkable.

FIG. 6. Tuberculosis lesion in the adrenal.

recognized grossly in the kidney as pyelonephritis proved to be of a granulomatous nature similar to the miliary nodules.

No organisms resembling *Histoplasma capsulatum* were found in any of the routine hematoxylin and eosin sections. This was contrary to our experience in other cases of disease due to

this cause, but to make the recognition of any such fungi easier most of these sections were also stained by the periodic acid-Schiff reaction of McManus. This stain is particularly useful because of a substance in the capsule of the fungi that is presumably of a polysaccharide nature and stains specifically. We might also

mention that although there is an acid-fast component in histoplasma, it is not of such amounts that the acid-fast stain regularly demonstrates the fungi; about a third of the organisms will stain with the usual acid-fast stain. We have not been successful in demonstrating any fungi in these sections by any method, but with the Ziehl-Nielsen stain acid-fast bacilli are present in all the lesions described. The report of acid-fast bacilli in pleural fluid removed four months before the patient's death is, therefore, more consistent with the lesions found at autopsy than the various preparations interpreted as histoplasma.

The original slide on which a diagnosis of histoplasmosis was made contained organisms with the morphologic characteristics of that fungus, but a negative culture at that time was very disturbing as cultures are more often positive than is the histologic recognition of the organism. The same problem is well reflected in a startling report by Dr. Alan Raftery<sup>6</sup> that in 436 appendices which were examined microscopically over 10 per cent contained organisms morphologically resembling *Histoplasma capsulatum*. That paper was discussed by Drs. R. J. Parsons, H. Reimann and others and the consensus appeared to be that positive identification of the specific organism is difficult even with cultures and treacherous when based on morphologic criteria alone. In the present case on two occasions, once with pleural fluid and once with ascitic fluid, organisms were described as having the morphologic character-

istics of histoplasma, yet cultures made on the same material were negative. Cultures of blood, brain and lung at autopsy did not demonstrate any histoplasma. The successful culture of the fungus from bone marrow in which no organisms were recognized microscopically is the best evidence we have that this patient did have histoplasmosis. The lesions in the tissue are all so intimately associated with acid-fast bacilli there is no evidence any other causal agent was at work. It does not seem possible to decide whether this case represents a carrier state or if it is a matter of the histoplasma being present in the body without producing lesions. We know that the lesions of histoplasmosis can heal, but there is no evidence of more than one kind of process in the tissues of this case. We have seen at least one case of associated tuberculosis and histoplasmosis, and histoplasmosis has occurred with unusual frequency in association with other chronically debilitating diseases such as the malignant lymphomas; so it might be considered that the major part of the disease in this patient was tuberculosis with a coincidental infection with histoplasma which was not demonstrable terminally.

*Final Anatomic Diagnoses:* Miliary tuberculosis, caseous and fibrocaseous, involving the lungs, with tuberculous pneumonia in the lower lobe of the left lung; mediastinal lymph nodes, liver, spleen, adrenals, kidneys, and bone marrow; tuberculous pleurisy, right, and pericarditis; tuberculous pyelonephritis.

*Acknowledgment:* Illustrations were made by the Department of Illustrations, Washington University School of Medicine.

<sup>6</sup> RAFTERY, A. Subclinical histoplasmosis. *J. A. M. A.*, 145: 216-220, 1951.

# Case Reports

## Mobilization of Gouty Tophi by Protracted Use of Uricosuric Agents\*

T. F. YÜ, M.D. and ALEXANDER B. GUTMAN, M.D.

*New York, New York*

IT has been known for some time that salicylates effect a marked increase in urinary urate excretion when administered to gouty subjects in substantial dosage, about 5 gm. daily, particularly if given in conjunction with sodium bicarbonate or other alkalinizing

indicates that this may be accomplished. Once the uricosuric regimen was fully established, no new tophi have appeared over months or years of observation in any of our patients with chronic tophaceous gout, including those patients in whom crops of new tophi had de-




FIG. 1. Schematic representation of clinical course of gouty patient, A. G., from onset of symptoms in 1932 to June, 1950. Peaks represent attacks of acute gouty arthritis, height indicating the severity and width the duration of each episode. The solid baseline indicates increasing severity of persistent manifestations of chronic tophaceous gout. The course is divided into three phases of development.

agents.<sup>1-5</sup> More recently it has been found that carinamide (4'-carboxyphenylmethanesulfon-anilide)<sup>4-6</sup> and benemid® (p-(di-n-propylsulfamyl)-benzoic acid)<sup>4,5,7,8</sup> also are potent uricosuric agents and are better tolerated, particularly benemid.<sup>8</sup> The increased urinary urate excretion produced by these drugs is generally attributed to suppression of tubular reabsorption of urate.

We have been exploring the usefulness of these uricosuric agents in the management of chronic gout with the hope that by maintaining augmented excretion of urate it might be possible to minimize deposition of urate in the tissues and thus obviate further development of the deformities and disabilities of chronic tophaceous gout. Our experience thus far

developed at relatively short intervals in the period immediately preceding initiation of therapy. Moreover, interruption of treatment was followed in some instances by resumption of overt new tophus formation within a few months.

Recently we have been obtaining evidence in some cases of reduction in size of long established tophi. The following instance is a case in point:

### CASE REPORT

A. G., a sixty-two year old male, was first seen in June, 1950, for gout of eighteen years' duration. His first attack of acute gouty arthritis in 1932 involved the right big toe and incapacitated him for only a few days. The subsequent development of the disease can be divided into

\* From the First Medical Service, The Mount Sinai Hospital, and the Department of Medicine, Columbia University College of Physicians and Surgeons, New York, N. Y. Supported in part by a grant from the John A. Hartford Foundation and a research grant from the National Institute of Arthritis and Metabolic Diseases, National Institutes of Health, U. S. Public Health Service.



three stages, each of approximately six years' duration. (Fig. 1.) From 1932 to 1938 he was subject to frequent attacks of podagra, which recurred at about monthly intervals, with no residual complaints between attacks. In 1933 he noted one small tophus on his right ear. In 1938 the episodes of acute gouty arthritis began to run a longer course, involving more joints and responding less readily and completely to colchicine. During this period the ankles, knees and elbows were frequently involved, leaving the toes relatively free. On many occasions several joints were swollen simultaneously. He also suffered recurrent attacks of acute olecranon bursitis, with serous effusions requiring repeated taps. In 1939 tophaceous deformities of the feet developed and slowly increased in size, initially at the first metatarsophalangeal joints, then at the fifth metatarsophalangeal joints and finally on the dorsum of the feet and about the ankle joints. In consequence he had more difficulty in walking and standing and was obliged to use oversize shoes. In 1945 a large tophaceous swelling appeared on the fourth finger of the left hand, together with a smaller tophus at the second metacarpophalangeal joint. About this time the serous effusions of the left olecranon bursa subsided, leaving a large permanent swelling. Since 1948 crops of small tophi had appeared in the lobes of both ears.

As a result of increasing disability due to chronic tophaceous gout, the patient had been unable since 1943 to carry on any regular work as a salesman. Treatment during this latter period consisted of colchicine, 1 mg., and aspirin, 0.65 gm., every night.

On examination in June, 1950, he presented the picture of advanced tophaceous gout. There were tophaceous deposits in both feet, especially around the metatarsophalangeal joints. These were firmly attached to the underlying bones. There were no discharging sinuses. One large tophus was palpable along the flexor tendons of the left ankle. The left elbow was the site of a solitary tophus about the size of a hen's egg. The proximal phalangeal joint of the left fourth finger was immobilized by a cherry-sized tophus; another was present at the second metacarpophalangeal joint of the left hand and a third large mass was palpable on the ulnar side of the left wrist. These tophi were firmly attached to the underlying joints. About six or seven small tophi were present in both ear lobes. Roentgenologic examination showed punched-

out areas in the medial condyle of the humerus, the proximal and middle phalanges of the left fourth finger and the first metatarsophalangeal joints of both feet, with large soft tissue swellings around these areas. The knees, ankles, wrists, right hand and elbow showed no significant joint changes.

The serum urate consistently exceeded 11 mg. per cent. The mean urinary urate excretion on a low purine diet was 512 mg./24 hours. Red cell count was 4.12 million; hemoglobin 12.8 gm. per cent; white cell count and differential normal. ESR was 59 mm. in the first hour (Westergren). The urine showed 1+ albumin, with occasional granular and hyaline casts, and a few red blood cells and white blood cells. Serum NPN was 59 mg. per cent and PSP excretion 25 per cent in the first hour and 13 per cent in the second hour. Blood pressure was 140/80 mm. Hg.

On June 26, 1950, surgical excision of the tophus at the left elbow was attempted under local anesthesia. The major part of the tophus, which was quite vascular, was found to extend to and was attached to the articular capsule and the ulnar collateral ligament. About 10 gm. of the tophaceous mass was removed, leaving the greater portion behind. The wound healed well. Olecranon bursitis with serous effusion subsequently developed, requiring repeated aspirations. The urate content of the aspirated fluid was usually somewhat higher than that of the serum, sampled simultaneously.

Starting on June 29th the patient was given carinamide, 12 gm. a day for eighteen days, and then benemid, 2 gm. a day for eighty-four days. He was maintained on a low purine, low protein diet throughout this period. Prophylactic colchicine was continued. The mean daily urinary urate excretion on this regimen was 644 mg., a daily excess excretion of approximately 132 mg. Thus in a period of 102 days (from June 29th to October 8th) the total excess excretion was about 13.5 gm. The patient's serum urate level fell to 6.5–8.0 mg. per cent. Since he was still having a great deal of diffuse pain and stiffness in various joints, medication was changed on October 26th to sodium salicylate combined with sodium bicarbonate. It was found that with 3 gm. salicylate a day he was more comfortable but there was virtually no uricosuric effect with this dosage. Mild salicylism soon developed, with tinnitus, diminution in hearing and gastric irritation. Because of




FIG. 2. Case A. G., tophaceous deformities of feet. A, December, 1950; B, September, 1951, after prolonged treatment with uricosuric drugs.

FIG. 3. Case A. G., tophaceous deformities of hands. A, December, 1950; B, September, 1951.

salicylism enteric-coated sodium salicylate was tried. The patient tolerated 4 gm. daily of the enteric-coated sodium salicylate very well and his diffuse joint pains disappeared almost entirely. On this dosage of salicylate his mean daily urinary urate excretion increased to 800 mg./24 hours, with an associated fall in serum urate to  $6.0 \pm 0.5$  mg. per cent. The dosage of enteric-coated sodium salicylate was increased to 5.2 gm. a day on December 7th, a dosage which he tolerated well, and the diet was liberalized somewhat in respect to protein intake. His serum urate level then was maintained at 5.0–5.5 mg. per cent and there was a daily urinary excretion of 800 to 1,000 mg. for the next three months. Since March, 1951, the serum urate levels have remained within  $4.0 \pm 1.0$  mg. per cent. The urinary excretion of urate has continued at levels of 700–800 mg. per twenty-four hours.

DECEMBER, 1951

Figures 2A, 3A, 4A and 5A indicate the status of his visible tophi in December, 1950. Since institution of uricosuric therapy in June, 1950, no new tophi had appeared, nor was there any change in the established tophi. At the end of February, 1951, the patient began to note that his shoes were becoming too large and the number of tophi on his ear lobes was decreasing. By May the decrease in size of tophi became quite apparent and by September, 1951, the changes were quite striking. (Figs. 2B, 3B, 4B and 5B.) We estimate from the urinary urate determinations that an excess of 100 gm. of urate was mobilized and excreted during the period of uricosuric therapy.

At the time the photographs were taken in September, 1951, the serum urate was 3.3 mg. per cent and the urinary urate excretion was 700 mg. per day. Red blood count was 4.25 million; hemoglobin 13.5 gm. per cent; white cell count,




FIG. 4. Case A. G., tophaceous deformity of left olecranon bursa. A, December, 1950; B, September, 1951.




FIG. 5. Case A. G., tophi of left ear. A, December, 1950; B, September, 1951.

differential and platelets were normal. ESR was 23 mm. in the first hour. The urine still showed 1 + albumin; the sediment contained a few red cells and leukocytes but no casts. Serum NPN was 45 mg. per cent. The PSP excretion was 28 per cent in the first hour and 15 per cent in the second hour. Blood pressure was 160/90 mm. Hg. Prothrombin time was 14 seconds (normal, 12 seconds). Serum salicylate level was 20 mg. per cent.

The patient is much improved and now able to carry out part-time activities. Of interest is the fact that except for a minor acute attack of gout early in July, 1950, shortly after starting benemid therapy, and another in October, 1950, he has been free of acute gouty arthritis while receiving 1 mg. of colchicine daily for prophylaxis.

#### COMMENTS

Disappearance of established tophi as a result of treatment with uricosuric drugs, such as occurred in the case described, was not anticipated. It has always seemed likely that old urate deposits in the tissues are beyond the reach of humoral agents, and the minimal mixing of intravenously injected  $N^{15}$ -labeled uric acid with tophus content observed by Benedict et al.<sup>9</sup> appeared to confirm this belief. However, the observations in the case reported herein seem to be unequivocal (Figs. 2 to 5) and are supported by data indicating a markedly negative urate balance for about one year, with mobilization of approximately 100 gm. urate from the tissues. Presumably, mobilization of urate from the tissues is facilitated by the sharp reduction in serum urate level produced by these agents.

Sometimes tophi spontaneously decrease in size and disappear, particularly from the ear or elsewhere after breaking through the skin and discharging their contents. There was no spontaneous external evacuation of urate in the present instance nor have we encountered so general a spontaneous reduction of tophi anywhere in the literature, or in 270 unpublished case histories of gout we have reviewed, or in our personally supervised series of sixty gouty cases prior to treatment with uricosuric agents. There are, moreover, other instances of disappearance of tophi in our patients following prolonged administration of salicylates in 5 gm. daily dosage. Case V. P., who had very extensive and crippling urate deposits particularly in the hands, feet and olecranon bursae, has also noted unequivocal reduction in size of many tophi, with partial restoration of function of the



extremities. In this instance some 90 gm. of excess urate were mobilized and excreted in the urine over a period of six months. Case L. R. has observed virtual disappearance of a hen's egg size tophus of the olecranon bursa. Several patients on protracted benemid medication similarly have noted disappearance or decrease in size of tophi, for the most part small or recently established deposits, together with lessened swelling of chronically enlarged joints. This has been most striking in the feet, enabling these patients to walk better and to wear smaller sized shoes than they had been accustomed to for years. The results with benemid are particularly encouraging because very few of our patients have been able to tolerate for long the required dosage of salicylates, which in amounts less than 5 gm. per day usually has little or no uricosuric effect and in doses of the order of 2 gm. per day is apt to cause retention of urate.<sup>1-5</sup> Benemid ordinarily produces sustained uricosuria when given to patients with chronic gout in doses of 0.5 or 1.0 gm. per day, a dosage level of low toxicity.<sup>8</sup>

A matter of some concern in protracted treatment of chronic gout with uricosuric agents is the possibility of aggravating renal damage by increasing the excretory urate load on the kidneys. So far we have not observed any indication of this, provided dosage is carefully regulated according to the urinary urate level, and ample fluids and alkalinizing salts are given; in Case A. G., in fact, the serum NPN declined from 59 to 45 mg. per cent under treatment, perhaps reflecting mobilization of urate deposits in the kidneys. In any event, so far as can be determined these drugs do not cause any significant increase in the amount of urate formed in the body or any increase in glomerular filtration rate, but they apparently do suppress urate reabsorption in the tubules. The chief hazard in their use would therefore seem to be precipitation of urate in the collecting tubules and this apparently can be minimized by the precautions indicated. The evidence that uric acid has a direct "toxic" action in concentrations of the order of magnitude under consideration is unsubstantial.<sup>10</sup>

As indicated in the case report, it is our practice to institute a low purine, restricted protein diet in conjunction with uricosuric

therapy in order to minimize urate production while augmenting urate excretion, so that maximal negative urate balance may be achieved. Even under these conditions mobilization of visible urate deposits is a slow process demanding the utmost patience and perseverance. Proper regulation throughout this period requires frequent determinations of serum and urinary urate levels, constant attention to diet and dosage levels of medication, and vigilant search for evidences of untoward side reactions.

#### SUMMARY

A case of advanced tophaceous gout exhibiting unequivocal reduction in size of many tophi in the course of protracted uricosuric therapy is described. It may be possible by proper use of uricosuric agents not only to minimize new tophus formation but also to cause the disappearance of old established tophi.

#### REFERENCES

1. JENNINGS, G. H. The value of sodium salicylate in the treatment of gout. *Reports on chronic rheumatic diseases*. No. 3: 106, 1937.
2. BAUER, W. and KLEMPERER, F. Medical progress: treatment of gout. *New England J. Med.*, 231: 681, 1944.
3. ROBINSON, W. D. and BLOCK, W. D. The effect of salicylates on the blood concentration and urinary excretion of urates and non-urate chromogens. *Proc. Central Soc. Clin. Research*, 19: 69, 1946.
4. GUTMAN, A. B. In Combined Staff Clinic on uric acid metabolism and gout. *Am. J. Med.*, 9: 799, 1950.
5. GUTMAN, A. B. Some recent advances in the study of uric acid metabolism and gout. *Bull. New York Acad. Med.*, 27: 144, 1951.
6. WOLFSON, W. Q., COHEN, C., LEVINE, R. and HUDDESTON, B. Transport and excretion of uric acid in man; physiologic significance of the uricosuric effect of carinamide. (Abstr.) *Am. J. Med.*, 4: 774, 1948.
7. GUTMAN, A. B. and YÜ, T. F. Benemid, (p-(di-n-propylsulfamyl)-benzoic acid) as uricosuric agent in chronic gouty arthritis. *Tr. A. Am. Physicians*, 64: 1951 (in press).
8. BISHOP, C., RAND, R. and TALBOTT, J. H. The effect of benemid (p-(di-n-propylsulfamyl)-benzoic acid) on uric acid metabolism in one normal and one gouty subject. *J. Clin. Investigation*, 30: 889, 1951.
9. BENEDICT, J. D., FORSHAM, P. H., ROCHE, M., SOLOWAY, S. and STETTEN, DEW., JR. The effect of salicylates and adrenocorticotrophic hormone upon the miscible pool of uric acid in gout. *J. Clin. Investigation*, 29: 1104, 1950.
10. GUTMAN, A. B. and YÜ, T. F. Gout, a derangement of purine metabolism. *Advances Int. Med.*, 5: 1951 (in press).

# Localized Amyloid Deposition in the Lower Respiratory Tract\*

AARON SCHOTTENFELD, M.D., LEON M. ARNOLD, M.D., JOHN G. GRUHN, M.D.  
and A. DAVID ETESS, M.D.

*New York, New York*

WE have recently had the opportunity to study a patient with amyloid infiltration of the trachea and main bronchi who was admitted to the hospital with the provisional diagnosis of bronchogenic carcinoma. The correct diagnosis was not suspected until the pathologist examined the bronchoscopic biopsy. We were therefore stimulated to review the literature for more information about the nature and frequency of this unusual lesion.

Local amyloid tumefactions confined to a single organ have been reported in the larynx,<sup>1-7</sup> urinary bladder,<sup>8-11</sup> conjunctiva,<sup>12-14</sup> mouth and nasopharynx<sup>15-18</sup> and a variety of other organs.<sup>19-23</sup> They have frequently been mistaken clinically for neoplasms. Local amyloidosis of the upper respiratory tract, especially the larynx, is more common than involvement of the lower respiratory tract including trachea, bronchi and pulmonary parenchyma. We were able to find twelve well documented cases and a questionable thirteenth case of local amyloid deposition in the lower respiratory tract in addition to our own case.

## CASE REPORT

M. L. (Montefiore Hospital No. 50886), a forty-three year old white male house painter, entered Montefiore Hospital on May 5, 1950. He first noted wheezing and shortness of breath on effort five years before admission but attributed his complaints to exposure to paints. Shortly thereafter a cough developed which was productive of up to one-half cupful of thick white sputum daily. Despite progressively increasing symptoms including dyspnea at rest he continued to work until the winter of 1945 when he was admitted to a hospital for pneumonia of the left lower lobe from which he rapidly recovered. He returned to work and was comfortable during

the following summer. Symptoms of dyspnea, wheezing and productive cough recurred with the onset of cold weather and persisted except for relatively asymptomatic intervals during the summer months. In the winter of 1948 he was again hospitalized for pneumonia, this time of the right lower lobe. Hemoptysis was noted for the first time on this occasion. Because of persistent right lower lobe consolidation bronchoscopy was advised but the patient refused consent. For the following fourteen months penicillin in oil was administered three times weekly and during this period he was well except for slight cough productive of a teaspoonful of sputum daily. During the winter of 1949 to 1950 there were two further episodes of pneumonia characterized by acute onset of fever, chills, productive cough, dyspnea and hemoptysis. Thereafter the sputum became more copious, viscid and greenish yellow. The symptoms did not subside during a vacation in Florida. After the patient's return home x-ray examination on March 29, 1950, revealed a shadow extending from the left hilum into the left lower lung field. Hospitalization was advised.

On admission to Montefiore Hospital the patient was found to be a well developed, well nourished, sun-tanned white male of forty-three years who was dyspneic and orthopneic. Temperature was 99°F., pulse rate 72, respiration 38 and blood pressure 110/70. The pertinent physical findings were as follows: The trachea was in the midline. Wheezes were palpable over the entire left chest. On auscultation inspiratory and expiratory wheezes were audible over both lung fields and there were rales over the lower left chest posteriorly. The leafs of the diaphragm were normally mobile with respiration. The heart was not enlarged or otherwise remarkable. The abdomen was soft and presented no masses

\* From the Medical and Laboratory Divisions and the Peroral Endoscopy Service, Montefiore Hospital, New York, N. Y.




FIG. 1. A, Chest film May 8, 1950; B, chest film January 25, 1951

or palpable organs. Rectal examination revealed a slightly enlarged, smooth prostate. The extremities were normal and clubbing was absent. No bone tenderness was elicited.

The admission chest films of May 8th (Fig. 1A) revealed a large area of irregular density extending from the left hilar region out into the left middle and lower lung fields and involving the base of the left lower lobe as well as the lower portion of the left upper lobe from hilum to lingula.

Initial bronchoscopy was performed on May 10th and disclosed an extremely extensive lesion consisting of projecting masses extending in irregular fashion from the subglottic area down to the carina, which was considerably broadened, and further into both main bronchi. The masses varied from pea-sized, globular projections into the lumen preventing distant visualization to broad, flat elevations. The main bronchi, particularly the left, were quite narrowed by the flat elevations. The masses were covered by intact mucous membrane and were not particularly friable. They were firm and hard, and were bitten away by the biopsy forceps with considerable difficulty. Very little bleeding was encountered on removing the projections either on the first examination or on subsequent ones. The bronchoscopist's initial impression was "externally infiltrating and metastasizing carcinoma." The pathologist found no tumor but reported amyloid deposition in the submucosa of the trachea.

Intensive investigation disclosed no evidence of generalized amyloid disease. The blood protein was 6.8 gm. per cent, albumin 5.0 gm. per

cent and globulin 1.8 gm. per cent. The Congo red test was negative (17 per cent removed after one hour). Bence-Jones proteinuria was absent. Marrow studies were negative for myeloma. A complete roentgenologic skeletal survey revealed no osseous abnormalities. Electrocardiograph was not remarkable.

The following additional laboratory studies were performed: Hemoglobin was 13.5 gm. per cent, red blood cells 4,170,000 per mm.,<sup>3</sup> and white blood cells 7,400 per mm.,<sup>3</sup> with a normal differential count. Urinalysis: specific gravity ranged from 1.012 to 1.022. All specimens were negative for sugar and albumin; sediments contained only occasional epithelial cells. Sputum smears and cultures were negative for acid-fast bacilli on three occasions. The Mazzini test was negative. Erythrocyte sedimentation rate was 9 mm. in one hour; fasting blood sugar 80 mg. per cent; blood urea nitrogen 8.0 mg. per cent; cephalin flocculation negative; thymol turbidity 1 unit; serum bilirubin 0.46 mg. per cent; cholesterol 254 mg. per cent.

Up to January 10, 1951, twelve additional bronchoscopies were performed to confirm the diagnosis and to remove the obstructing masses. Eleven biopsy specimens taken from the trachea and bronchi between May 10, 1950, and January 10, 1951, were available for review. Gross specimens varied in size from small, ruddy tissue masses less than 3 mm. in diameter to a mass of small fragments having an aggregate volume of 4 cc., the largest individual particle measuring 8 mm. in greatest dimension. Multiple sections of trachea and bronchi included only the mucosal and submucosal portions of the wall. (Fig. 2.)






FIG. 2. A, Bronchial biopsy; amyloid deposition through entire tunica propria; mucosa on left; bronchial gland lower left corner (original magnification 105  $\times$ ). B, bronchus; mucosa of pseudostratified ciliated columnar cells; amyloid and fat in tunica propria (original magnification 255  $\times$ ). C, amorphous acellular hyaline amyloid in tunica propria; mucosa at upper left (original magnification 550  $\times$ ). D, metaplastic bone and cartilage in tunica propria (original magnification 85  $\times$ ). E, foci of adipose tissue adjacent to amyloid deposits in tunica propria (original magnification 105  $\times$ ).

The mucosa consisted of a pseudostratified columnar ciliated epithelium with occasional goblet cells resting on a thick basement membrane. Beneath the mucosa were irregular masses and clumps of amorphous hyaline material which stained pink-red with hematoxylin-eosin, bright orange with various lots of Congo red, and purplish red with metachromatic dyes such as crystal and methyl violet. All stains for amyloid were confirmed on fresh frozen sections. Unequivocally positive reactions for amyloid were obtained on each of the eleven submitted specimens. Interspersed amidst the amyloid masses were occasional normal tracheal and bronchial glands, small irregular islands of metaplastic cartilage, a solitary focus of metaplastic bone with hematopoietic tissue within its medullary space and areas of mature adipose tissue. No giant cells were found.

The patient was treated by piecemeal removal of the obstructing nodular masses using an ordinary Yankauer type of bronchoscopic cutting biopsy forceps. The largest masses were in the upper and lower trachea. At first bronchoscopy was performed weekly, later every two weeks and, since hospital discharge on July 28, 1950,

approximately every month. Following each bronchoscopic removal of tissue there was subjective and objective improvement. After ten bronchoscopies the patient could comfortably perform a day's work. He still has a morning cough productive of yellowish sputum which is blood tinged on rare occasions, usually during the week following a bronchoscopy. He is no longer dyspneic on rest or ordinary activities. Wheeze is less prominent both subjectively and objectively. It is noted by the patient only after walking rapidly up two flights of stairs and during sexual intercourse. Repeated chest films from May 3, 1950, to November 25, 1950, demonstrated no remarkable change in the infiltration of the left lung. However, in the interval from November 25, 1950, to January 25, 1951, very appreciable resolution of the pulmonary density occurred. (Fig. 1B.)

Removal of the amyloid by forceps is still in progress but the therapeutic plan may include an attempt at further enlargement of the airway, particularly the lower bronchi, by electrocoagulation. It has been interesting to note that there has been little or no increase in size of the flat lesions. It has therefore been possible to keep

ahead of any growth of the lesions and to improve the airway even with therapy at monthly intervals.

#### COMMENTS

In Table I we have tabulated certain of the recorded data available from the thirteen<sup>24-36</sup> previously recorded cases and from our case.

dependent upon the exact location and extent of the amyloid deposition. When the deposition is small in amount or located in silent areas, no symptoms develop. Thus in Case 2 presenting only a bean-sized submucous nodule in the posterior tracheal wall and in Cases 8 and 9 in which amyloid masses were confined to the pulmonary parenchyma the diagnoses were made

TABLE I

Case	Author	Age	Sex	Duration of Symptoms Prior to Diagnosis	How Diagnosed	Associated Pathologic Condition	Location of Amyloid
1	Balser <sup>24</sup>	66	M	19 yr.; "bronchitis" with evidence of progressive broncho-stenosis	Autopsy	Chronic bronchiectasis and pneumonitis of R.U.L.; several ossified areas in right lung	Trachea and main bronchi
2	Kraus <sup>25</sup>	..	M	None recorded	Autopsy	Pulmonary emphysema and malnutrition; acute pneumonia	Trachea
3	Manasse <sup>26</sup>	63	M	None recorded	Autopsy	Perforated duodenal ulcer with generalized peritonitis	Larynx and trachea
4	Glockner <sup>27</sup>	76	M	None recorded	Autopsy	B.P.H.; cystitis, hydronephrosis and pyelonephritis	Larynx, trachea and main bronchi
5	Johanni <sup>28</sup>	66	F	15 mo.; hoarseness, cough; later progressive dyspnea	Autopsy	Incomplete description of associated pathologic condition	Larynx and trachea
6	Herxheimer <sup>29</sup>	65	M	40 yr.; hoarseness	Autopsy	Abscess of R.L.L.; right empyema	Larynx, trachea and lung
7	von Werdt <sup>30</sup>	71	M	6 yr.; intermittent hoarseness; in last year dyspnea with stridor	Surgical removal of tumor; autopsy 11 hr. postoperatively	Bronchiectasis R.U.L. and R.M.L.	Larynx, trachea and main bronchi
8	Meyer <sup>31</sup>	57	M	None recorded	Autopsy	Chronic interstitial nephritis, arteriosclerosis, early cirrhosis of the liver, fresh cerebral hemorrhage	Lung
9	Hallermann <sup>32</sup>	52	F	None recorded	Autopsy	Postoperative pneumonia of R.L.L., R.M.L., L.L.L.; right empyema; thoracotomy	Lung
10	Bauer <sup>33</sup>	30	M	1 yr.; dyspnea and hoarseness	Laryngoscopy with biopsy	Goiter	Larynx and trachea
11	Bloch and Soulas <sup>34</sup>	46	F	4 mo.; cough, increasing dyspnea and hoarseness	Bronchoscopy with biopsy	None described	Larynx and trachea
12	Falconer <sup>35</sup>	71	M	2 mo.; increasing dyspnea	Autopsy	Gastric ulcers; hepatic cirrhosis with esophageal varices	Trachea and main bronchi
13	Weismann <i>et al.</i> <sup>36</sup>	35	M	9 yr.; cough and intermittent hemoptysis	Right pneumonectomy	Pneumonitis R.L.L. and R.M.L.	Right main bronchus and branches
14	Authors' case	43	M	5 yr.; cough, dyspnea, wheezing	Bronchoscopy with biopsy	Pneumonic infiltration L.L.L. and L.U.L.	Trachea and main bronchi

**Age Incidence.** Amyloid deposition in the lower respiratory tract is apparently a disease discovered in later adult life, nine of the fourteen patients being over fifty, with one case in which the age was not stated. The youngest patient was thirty years of age, the oldest seventy-six.

**Sex Incidence.** Males predominated in this group in a proportion of eleven to three. This differs from primary systemic amyloidosis, Eisen<sup>37</sup> having reported twenty-six males and twenty-two females in the cases collected by him.

**Clinical Features.** The duration of recorded symptoms was quite variable in the cases reviewed. In Case 12 there was two months of increasing dyspnea until death. In Case 6 hoarseness was present for forty years. The clinical picture is extremely variable and is

as incidental findings at autopsy after death due to unrelated major disease.

When the larynx is involved in addition to the lower respiratory tract, as occurred in five instances (Cases 5, 6, 7, 10 and 11), hoarseness is a prominent symptom. Involvement of the larynx alone is not within the scope of this review and has been extensively reviewed elsewhere. Patients in this older age group with persistent hoarseness may be suspected of having laryngeal carcinoma.

Symptoms and signs resulting from involvement of the trachea and bronchi are expressions of the amount of obstruction produced. One may therefore encounter various degrees of dyspnea, cough and wheezing with the superimposition at times of the features of acute recurrent pneu-

monia or chronic bronchiectasis. It is in this type of involvement that the diagnostic possibilities will be diverse and will run the gamut of asthma, foreign body, unresolved pneumonia, tuberculosis, bronchiectasis, adenoma and carcinoma.

tive carcinoma from which this condition of amyloid deposition would require differentiation on the basis of bronchoscopic appearance. The appearance is almost unique and does not resemble that of Boeck's sarcoïd, tuberculosis or any of the lymphomatous diseases.

**Pathologic Features. Distribution of lesions:** The distribution of amyloid in the reported cases is summarized below from Table I.




FIG. 3. Schematic representation of the location and extent of the amyloid deposits in fourteen cases of amyloidosis of the lower respiratory tract.

Diagnosis of this condition is almost completely dependent upon pathologic study of biopsy material. The lesion is so uncommon that on gross inspection alone no observer would venture the diagnosis. As has been mentioned, in our case the initial gross impression on bronchoscopy was infiltrating carcinoma of the submucosally infiltrative type. When viewed in retrospect certain features speak against such a diagnosis. These include (1) the unusual extent of the lesion throughout the entire trachea, carina and both major bronchi, (2) the intact mucous membrane, (3) the firm, tough consistency of the masses and their unusual resistance to the cutting forceps and (4) the paucity of bleeding on removal of the masses. It is difficult to imagine any disease other than infiltra-

No. of Cases	Location of Amyloid
4	Larynx and trachea
2	Larynx, trachea and main bronchi
1	Larynx, trachea and lung
1	Trachea alone
3	Trachea and main bronchi
1	Right main bronchus
2	Lung (1 questionable)

The extent of involvement is shown schematically in Figure 3.

The absence of amyloid elsewhere in the body was confirmed by autopsy in eleven cases. The remaining three patients (Cases 11, 13 and 14) were alive at the time of publication of the case reports. In these cases there was no clinical or laboratory evidence of more generalized amyloidosis.

In all but one of the cases of tracheobronchial deposition amyloid was present within the submucosa and would have been available for bronchoscopic biopsy. In Case 13 if a biopsy had been obtained it would not have revealed the amyloid which was deposited in the external connective tissue coats around the right main bronchus and its branches, around the right pulmonary artery and inferior pulmonary vein.

**Staining properties:** As one of his five criteria for "atypical amyloidosis" Lubarsch<sup>38</sup> cited the variability of reaction with amyloid stains. Reference to Table II which summarizes the published data on staining reactions in the reported cases indicates that they conform in large measure to this generalization. The reports of Cases 10 and 11 did not include specific descriptions of staining properties. In both instances the pathologic description was "local amyloid," presumably on the basis of positive specific stains. Case 9 is questionable because of consistently negative staining reactions. It should be emphasized, however, that in the remaining eleven cases there were specific descriptions of at least one acceptable staining reaction identifying the material as amyloid. Perhaps too much emphasis has been placed on atypical staining



as a criterion of so-called atypical, primary or local tumor-forming amyloidosis.

*Secondary changes within the amyloid tumefactions:* Among the most conspicuous of the incidental changes associated with amyloid tumors of the lower respiratory tract are calcification and metaplastic bone formation. Calcification, ossi-

3. The clinical and pathologic features of thirteen previously reported cases and our fourteenth case are reviewed and tabulated.

## REFERENCES

1. SOTTI, G. Amyloidosis of larynx. *Pathologica*, 20: 106, 1928.

TABLE II

Case	Iodine	Iodine and H <sub>2</sub> SO <sub>4</sub>	Methyl Violet	Gentian Violet	Congo Red
1	Positive	Variable		Positive (variable)	
2	Inconclusive	Variable	Positive		
3	Positive			Positive	
4	Positive (not uniform)	Variable	Positive		
5	Positive	Variable	Positive		
6*	Positive	Positive	Positive		
7*	Positive	Inconclusive	Negative	Negative	
8* Small masses	Positive			Positive	
Large masses	Positive			Negative	
9	Negative		Negative	Negative	Negative
10	Reported as local amyloid; staining reactions not given				
11	Reported as local amyloid; staining reactions not given				
12			Positive		
13	Negative (gross)		Positive (not uniform)		Positive (not uniform)
14			Positive		Positive

\* Also positive van Gieson's stain

fication or both was described in ten of the twelve cases in which the pathology was given in detail. Similar changes are not uncommon in bronchial adenomas but are quite rare in carcinoma of the lung.

The occurrence of amyloid within the tracheal adipose tissue was noted in our case as well as in Case 3. In Case 7 amyloid was deposited in the peritracheal and hilar adipose tissue.

Giant cells were not encountered in our material but were noted as a conspicuous element in Cases 5, 6, 7, 8 and 10. Only in Case 5 did the author report the presence of amyloid substance within giant cells.

## SUMMARY AND CONCLUSIONS

1. A case of local amyloid tumefaction of the trachea and major bronchi is described. The condition was diagnosed by bronchoscopic biopsy and treated palliatively by periodic removal of amyloid masses.

2. Attention is called to this clinical entity in the differential diagnosis of stenotic lesions of the lower respiratory tract.

DECEMBER, 1951

2. GIOACCHINI, R. Amyloid tumors of the larynx. *Valsalva*, 3: 35, 1932.
3. KRIEGSMANN, G. Amyloid tumors of larynx. *Arch. f. Ohren-, Nasen-u. Kehlkopf.*, 130: 198, 1932.
4. JORGENSEN, A. Amyloid tumor of larynx. *Ztschr. f. Laryng., Rhin.*, 25: 209, 1934.
5. DELLA VEDOVA, A. JR. Tumoral amyloidosis of larynx in boy eleven years old. *Valsalva*, 10: 808, 1934.
6. KRAMER, R. and SOM, M. L. Local tumor-like deposits of amyloid in larynx; report of case with review of literature. *Arch. Otolaryng.*, 21: 324, March, 1935.
7. CLERF, L. H. Amyloid tumor of larynx. *Tr. Am. Laryng. A.*, 64: 146, 1942.
8. LUCKSCH, F. Local amyloid formation in the urinary bladder. *Verhandl. d. deutsch. path. Gesellsch.*, 7: 34, 1904.
9. CHWALLA, R. A case of amyloidosis at the bladder neck resembling a tumor. *Urol. & Cutan. Rev.*, 36: 381, 1932.
10. RUSCHE, C. F. and BACON, S. K. Solitary tumor-like amyloidosis of the urinary bladder. *Tr. Am. A. Genito-Urin. Surgeons*, 34: 163, 1941.
11. CORBITT, R. W., BRODERS, A. C. and POOL, T. L. Amyloidosis of urinary bladder. *J. Urol.*, 52: 153, 1944.
12. LEWKOJEW, E. F. Primary amyloid degeneration of cornea. *Klin. Monatsbl. f. Augenh.*, 85: 117, 1930.

13. KREIBIG, W. Amyloid degeneration of conjunctiva. *Ztschr. f. Augenh.*, 90: 129, 1936.
14. SMALTINO, M. Hyaline-amyloid degeneration of the cornea in trachoma. *Boll. d'ocul.*, 16: 126, 1937.
15. SCHNEIDER, H. Amyloid tumor of vestibule of mouth. *Ztschr. f. Stomatol.*, 36: 288, 1938.
16. SCHMIDT, M. Tumor-like amyloid degeneration in upper air passages with special regard to occurrence in nasopharynx. *Arch. f. Ohren-, Nasen- u. Kehlkopfch.*, 117: 300, 1928.
17. MUTSCHLER, H. H. Occurrence of solitary tumor-like amyloid growth in the palatine tonsil; case and study of pathogenesis and symptoms of amyloid tumors. *Ztschr. f. Laryng., Rhin.*, 24: 228, 1933.
18. CHIARI, H. Isolated local amyloid deposit in tonsil. *Monatschr. f. Ohrenh.*, 71: 666, 1937.
19. TILP, A. Local tumor-like amyloid of the urethra. *Centralbl. f. all. Path. u. path. Anat.*, 20: 913, 1909.
20. VON ALBERTINI, A. Local amyloid of the urethra. *Frankfurt. Ztschr. f. Path.*, 33: 248, 1925.
21. LEHMAN, G. Local amyloid deposition in the wall of the ureter. *Zentralbl. f. allg. Pathologie u. path. Anat.*, 68: 209, 1937.
22. WINER, L. H. Local amyloidosis of the skin. *Arch. Dermat. & Syph.*, 23: 866, 1931.
23. DELARUE, J. and THOREL, M. Amyloid tumor of fallopian tube. *Ann. d'anat. path.*, 15: 1050, 1938.
24. BALSER, W. Tracheo- and bronchostenosis with amyloid in the wall of the airway. *Virchows Arch. f. path. Anat.*, 91: 67, 1883.
25. KRAUS, F. New observations on amyloid masses. *Ztschr. f. Heilk.*, 7: 245, 1886.
26. MANASSE, P. Multiple amyloid tumefactions of the upper airway. *Virchows Arch. f. path. Anat.*, 159: 117, 1900.
27. GLOCKNER, A. Local tumor-like amyloid of the larynx, trachea and main bronchi with laryngo- and tracheostenosis produced thereby. *Virchows Arch. f. path. Anat.*, 160: 583, 1900.
28. JOHANNI, J. U. An amyloid tumor of the larynx and trachea. *Arch. f. Laryng. u. Rhin.*, 14: 321, 1903.
29. HERXHEIMER, G. Multiple amyloid tumors of the larynx and lungs. *Virchows Arch. f. path. Anat.*, 174: 130, 1903.
30. VON WERDT, F. Local amyloid in the entire respiratory tract. *Beitr. z. path. Anat. u. z. allg. Path.*, 43: 239, 1908.
31. MEYER, O. Local tumor-like amyloid of the lungs. *Frankfurt. Ztschr. f. Path.*, 8: 304, 1911.
32. HALLERMANN, W. Peculiar masses in the lungs resembling local tumor-like amyloid. *Frankfurt. Ztschr. f. Path.*, 36: 471, 1928.
33. BAUER, A. A case of local amyloidosis of the trachea. *München. med. Wchnschr.*, 79: 1596, 1932.
34. BLOCH, A. and SOULAS, A. Amyloid tumor of the trachea. *Bronchoscop., œsophagoscop. et gastroscop.*, pp. 211-213, 1937.
35. FALCONER, B. A case of amyloid tumors of the trachea and main bronchi with dyspnea. *Acta oto-laryng.*, 26: 353, 1938.
36. WEISMANN, R. E., CLAGETT, O. T. and McDONALD, J. R. Amyloid disease of the lung treated by pneumonectomy. *J. Thoracic Surg.*, 16: 269, 1947.
37. EISEN, H. N. Primary systemic amyloidosis. *Am. J. Med.*, 1: 144, 1946.
38. LUBARSCH, O. Recognition of atypical amyloid depositions. *Virchows Arch. f. path. Anat.*, 271: 867, 1929.

## AUTHOR INDEX VOLUME XI

- Abbott, L. D., Jr., 521, 621  
 Abildskov, J. A., 237, 618  
 Adair, C., 188, 523  
 Adams, J. E., 624  
 Adlersberg, D., 600  
 Adriani, J., 239  
 Aikawa, J. K., 237  
 Alexander, L. C., 517  
 Almy, T. P., 67  
 Alpert, L. K., 516  
 Antzis, E., 531  
 Arnold, L. M., 770  
 Atlas, L. T., 522  
 Austrian, R., 667
- Bader, M. E., 715  
 Bailey, C. C., 590  
 Bailey, O. T., 523  
 Barr, D. P., 468, 480  
 Barrow, J. G., 616  
 Bartels, C. C., 448  
 Bartels, E. C., 448  
 Bayrd, E. D., 202  
 Beck, G. J., 630  
 Beeman, E. A., 522  
 Belding, H. W., 243  
 Bell, C. E., Jr., 238  
 Bell, J. A., 522  
 Bellet, S., 431  
 Berenson, G. S., 617  
 Berger, E. H., 398  
 Berliner, R. W., 274  
 Berry, K., 431  
 Best, M. M., 240  
 Beyers, M. R., 521  
 Bickerman, H. A., 630  
 Bierman, H. R., 625  
 Bine, R., Jr., 626  
 Bloom, W. L., 238, 252, 618  
 Blumgart, H. L., 271, 331  
 Boger, W. P., 517  
 Boldrey, E. B., 624  
 Bond, E. D., 588  
 Bondy, P. K., 619, 724  
 Bortin, M. M., 40  
 Botwin, A., 250  
 Boyle, A. J., 517  
 Brainerd, H. D., 627  
 Brauer, R. W., 237  
 Brawner, D. L., 621  
 Brill, I. C., 410  
 Brown, E., 620  
 Brown, T. McP., 518  
 Brown, W. E., 244  
 Brun, C., 324
- Brunkow, C. W., 398  
 Bruyn, H. B., 627  
 Bunn, P. A., 188, 523  
 Burch, G. E., 237, 618  
 Burnell, J. M., 627  
 Burnett, C. H., 238  
 Burnett, S., 244  
 Burns, T. W., 239  
 Burrows, B. A., 238  
 Burson, N., Jr., 618, 620  
 Burtner, O. W., 621  
 Bush, O. F., 239  
 Byers, S. O., 626  
 Byron, R. L., Jr., 625
- Call, L. S., 123  
 Cannon, J. L., 622  
 Caplovitz, C. D., 617  
 Cargill, W. H., 239  
 Carpenter, F. M., 519  
 Casten, G., 247  
 Cayer, D., 240  
 Chamovitz, D. L., 255  
 Chapman, D. W., 26, 616, 617  
 Cheever, F. S., 523  
 Chesky, K., 520  
 Coe, W. S., 240  
 Coggins, W. J., 239  
 Cohen, E. B., 645  
 Cole, R. M., 522  
 Coleman, D. H., 179  
 Conan, N. J., 283  
 Cooper, G. R., 249  
 Corday, E., 410  
 Cornatzer, W. E., 240  
 Cronvich, J. A., 237, 618  
 Cosby, R. S., 31  
 Cournand, A., 667  
 Cummings, M. M., 252
- Da Costa, J. M., 559  
 Daines, M. C., 625  
 Daniels, J. B., 523  
 Davidson, J. D., 736  
 Davies, D. F., 240  
 Davis, W. D., Jr., 251  
 Davis, W. M., 639  
 Davison, K. B., 624  
 Devlin, B., 706  
 Dick, H. L. H., 625  
 Dimitroff, S. P., 31  
 Dinning, J. S., 241  
 Dock, W., 730  
 Donald, K. W., 667  
 Doolan, P. D., 244, 619, 622
- Doyle, J. T., 241, 253  
 Dresdale, D. T., 686  
 Duff, G. L., 92  
 Duke, T. W., 249, 622  
 Duncan, G. G., 535  
 Dunn, J., 531  
 Dunsmore, R. A., 516
- Earle, D. P., 283  
 Eder, H. A., 468, 480  
 Edgren, D. C., 252  
 Eichna, L. W., 283  
 Ellenberg, M., 170  
 Elliott, H. W., 624  
 Ellison, R. R., 521  
 Emlet, J. R., 241  
 Engel, F. L., 239  
 Entwisle, G., 461  
 Estes, E. H., Jr., 241, 253  
 Etess, A. D., 770  
 Evans, J. A., 448  
 Evans, R. S., 624
- Feltman, J. A., 706  
 Fenn, G. K., 403  
 Ferguson, J. H., 245  
 Fieber, M. H., 516  
 Fields, J., 410  
 Finch, C. A., 629  
 Findley, T., 251  
 Finnerty, F. A., Jr., 242  
 Finston, E., 410  
 Fischer, E., 615  
 Fitzhugh, F. W., 615  
 Fleming, J. W., 618  
 Flieg, W., 410  
 Foraker, A. G., 621  
 Frayser, R., 243  
 Freedberg, A. S., 44, 255  
 Freis, E. D., 242, 519  
 Fremont-Smith, K., 627  
 Friedman, M., 626
- Gargill, S. L., 44  
 Garlett, E. L., 431  
 Geffer, W. I., 431  
 Gendel, B. R., 267  
 Ginsberg, V., 521  
 Gitlow, S. E., 521  
 Gofman, J. W., 358  
 Gordon, D., 242  
 Gordon, G. S., 629  
 Gordon, V. H., 246  
 Gottstein, W. K., 403  
 Gould, R. G., 209



- Graettinger, J. S., 3  
 Graham, D. M., 358  
 Graybiel, A., 3  
 Green, H. D., 238  
 Griffith, G. C., 31  
 Grisson, K. S., 241  
 Grollman, A., 247  
 Gruhn, J. G., 770  
 Gutman, A. B., 765  
 Guyton, A. C., 242
- Hallman, B. L., 619, 724  
 Hamwi, G. J., 261  
 Handley, C. A., 615  
 Harkins, H. N., 628  
 Harris, J., 250, 518  
 Harrison, J. H., 55  
 Hecht, H. H., 625  
 Heck, F. J., 202  
 Herwitz, G. K., 442  
 Heyman, A., 239, 249, 622  
 Hickam, J. B., 243  
 Hill, E. H., 520  
 Holland, B. C., 243  
 Horn, J., 517  
 Horwitt, B., 242  
 Huebner, R. J., 522  
 Hufnagel, C. A., 243  
 Huie, R. A., Jr., 615  
 Hurst, W. W., 629  
 Hutaff, L. W., 243
- Iglauder, A., 442  
 Ingbar, S. H., 238  
 Iseri, L. T., 517  
 Iversen, P., 324
- Jackson, C. E., 237, 618  
 Jackson, G. W., 244  
 Jacox, R. F., 518  
 James, D. F., 244  
 James, G. W., 521, 621  
 Jawetz, E., 626  
 Jenkins, D. E., 620  
 Jensen, L. C., Jr., 621  
 Johnson, R. L., 242, 519  
 Johnson, T. D., 244  
 Jones, F., 247  
 Jones, H. B., 358  
 Jungck, E. C., 244
- Kallsen, R. A., 624  
 Kaufman, J., 442  
 Keefer, C. S., 665  
 Kellaway, P. E., 619  
 Kelly, K. H., 625  
 Kennamer, S. R., 410  
 Kennedy, T. J., Jr., 274  
 Kerby, G. P., 623
- Kern, F., Jr., 67  
 Keye, J. D., Jr., 621  
 Kinsell, L. W., 628  
 Kirby, W. M. M., 179, 627  
 Knight, V. H., 237  
 Kruger, H. E., 410  
 Krumbhaar, E. B., 540  
 Kunkle, E. C., 249  
 Kyle, L. H., 244, 252, 619, 622
- Laken, B., 157  
 Lambeth, W. A., 240  
 Lange, J., 628  
 Lauson, H. D., 135  
 Lawson, H. C., 240  
 Layne, J. A., 629  
 Lemley, J. M., 245  
 Leslie, S. H., 157  
 Lesses, M. F., 44  
 Levin, W. C., 620  
 Levinson, D. C., 31  
 Levitt, M. F., 715  
 Lewis, J. H., 245  
 Lichtman, H., 521  
 Lieberman, D., 626  
 Lindgren, F. T., 358  
 Lindsay, A., 247  
 Lloyd, C. W., 518  
 Loew, E. R., 461  
 Lorber, S. H., 520  
 Lyon, T. P., 358  
 Lyons, R. H., 517, 518
- Madison, L., 246, 250  
 Malbin, B., 312  
 Manchester, B., 520  
 Marble, A., 590  
 Margen, S., 628  
 Marlow, A. A., 624  
 Martin, S. P., 246, 623  
 Marvin, H. N., 241, 246  
 Mason, M. F., 247  
 Mason, R. E., 524  
 Master, A. M., 520  
 Matteucci, W. V., 517  
 Mayerson, H. S., 248  
 McCabe, J., 517  
 McCaughey, R. S., 517  
 McCawley, E. L., 625  
 McClement, J. H., 667  
 McMillan, G. C., 92  
 Meadows, J. C., 251  
 Meneely, G. R., 245  
 Merchant, W. R., 518  
 Merrill, A. J., 615  
 Metcalf, B. H., 241  
 Michael, M., Jr., 248, 252  
 Michaels, G., 628  
 Michtom, R. J., 686
- Miller, H., 240  
 Moore, A. E., 521  
 Moser, M., 520  
 Most, H., 639  
 Moyer, J. H., 247, 615, 617  
 Mudge, G. H., 302  
 Muirhead, E. E., 247  
 Muller, J. C., 253  
 Mundy, W. L., 630  
 Myers, G. B., 517  
 Myers, J. D., 248, 251
- Neubauer, R. A., 615  
 Newman, W., 706  
 Nichols, A. V., 358  
 Nickerson, M., 123  
 Nouse, D. C., 517  
 Nowinski, W., 616
- Oblath, R. W., 410  
 Ochs, L., 246, 250  
 O'Neill, J. F., 571  
 Orloff, J., 274  
 Ory, A. A., 617  
 Osborne, J. A., 410  
 Osgood, E. E., 624  
 Osserman, K. E., 170  
 Otto, J. C., 557
- Packard, J. M., 3  
 Pannill, F. C., 616  
 Pappenheimer, A. M., 523  
 Patterson, J. L., Jr., 249, 622  
 Paul, J. R., 577  
 Peers, J. H., 522  
 Pepper, O. H. P., 572  
 Perkoff, G. T., 628  
 Pessotti, R. L., 237  
 Pfeiffer, J. B., 249  
 Pines, K. L., 302  
 Pizzolato, P., 237  
 Plitman, G. I., 267  
 Pordy, L., 520  
 Porter, R. R., 615  
 Poutasse, E. F., 55  
 Prinzmetal, M., 410
- Rabkin, B., 520  
 Radding, R. S., 653  
 Ralli, E. P., 157  
 Ransmeier, J. C., 620  
 Rast, C. L., 253  
 Relman, A. S., 238  
 Renzetti, A. D., Jr., 667  
 Riley, R. L., 667  
 Rose, J. C., 242  
 Rosenberg, B. L., 75  
 Roth, D., 128  
 Rubitsky, H. J., 448

Rundles, R. W., 249, 253  
Rush, B., 546, 551  
Ruskin, A., 616  
Ruskin, B., 616  
Russ, E. M., 468, 480

Sagemen, C. B., 520  
Sauvage, L. R., 628  
Scaparone, M., 627  
Schaeffer, M., 619  
Schemm, F. R., 629  
Schilero, A. J., 531  
Schimmel, N. H., 517  
Schnaper, H. W., 242, 519  
Schneider, R. G., 620  
Schoen, A. M., 250  
Schottenfeld, A., 770  
Schottstaedt, M. F., 626  
Schroeder, H. A., 240, 252  
Schultz, M., 686  
Schwartz, L., 516  
Sealy, W. C., 250  
Seaman, A. J., 624  
Segaloff, A., 242  
Sellers, A. L., 410  
Sellers, S., 239  
Sensenbach, W., 246, 250  
Shaffer, C. F., 26  
Shay, H., 520  
Sheedy, J. A., 403  
Sherry, S., 283  
Shulman, N. R., 519  
Shure, M., 239  
Siegal, S., 196

Sikes, C. R., 616  
Silver, S., 40, 516  
Sisson, J. H., 238  
Skaggs, R. H., 616  
Sleisenger, M. H., 255  
Smith, C. W., 398  
Smith, L. A., 410  
Sokolow, M., 626  
Southam, C. M., 521  
Spain, D. M., 128  
Sternheimer, R., 312  
Stevens, A. R., Jr., 629  
Stevens, J. E., 202  
Stolk, N. J., 67  
Stone, C. A., 461  
Strisower, B., 358  
Stueck, G. H., Jr., 157  
Sugar, B., 627  
Sugar, S. J. N., 516  
Sutherland, C. G., 244  
Sutherland, V. C., 624

Tagnon, H. J., 519  
Tarver, H., 628  
Taylor, W. J., 251  
Taymor, R. C., 520  
Thomas, H. M., Jr., 581  
Thompson, R. T., 519  
Tivey, H., 624  
Tocantins, L. M., 568, 571  
Tuohy, E., 519  
Tyler, F. H., 628  
Tyor, M. P., 237  
Ureles, A. L., 44

Voigt, E. H., 251  
Volwiler, W., 627

Walker, W. G., Jr., 630  
Walsh, W. P., 244, 252, 619, 622  
Warren, J. V., 241, 253, 622  
Wasserman, K., 248  
Watson, J., 521  
Wessler, S., 331  
Westphal, R. D., 619  
Whorton, C. M., 248, 252  
Willett, R. W., 249, 253  
Williams, A. H., 252  
Wilson, J. S., 241, 253  
Wise, R. A., 619  
Wishinsky, H., 622  
Witham, A. C., 618  
Woldow, A., 516  
Wolfe, S., 521  
Womack, C. R., 247  
Woodin, W. G., 188  
Woodruff, C. W., 253

Yohalem, S. B., 40  
Youmans, J. B., 133  
Young, N. F., 615  
Young, W. G., Jr., 250  
Yow, E. M., 254, 620  
Yü, T. F., 765

Zazeela, H., 409  
Zimmerman, H. J., 518  
Zinn, W. J., 31  
Zoll, P. M., 331

# SUBJECT INDEX VOLUME XI

(ab.) = Abstract; (CPC) = Clinico-pathologic Conference; (E.) = Editorial

## Acidification of urine and potassium metabolism, 274

### ACTH

- adrenal response to (ab.), 242
- and cortisone, use of preoperatively (ab.), 619
- and remission of thrombocytopenic purpura (ab.), 624
- and rheumatic carditis, 128
- effect of, on fluid and electrolyte distribution, 715
- therapy in periarteritis nodosa, 630
- Adrenalin test in hypersplenism (ab.), 624
- Adrenergic blockade for cardiospasm, 123
- Adrenocorticotrophic hormone in trichinosis, 639
- Allylmercaptomethyl penicillin, 188
- Alveolar-capillary diffusion, impairment of, 667
- Amyloid deposition, localized, in respiratory tract, 770

### Anemia

- effects of, on hepatic blood flow and splanchnic metabolism (ab.), 248
- in renal insufficiency (ab.), 247
- megaloblastic, response of, to citrovorum factor (ab.), 253
- pernicious, citrovorum factor in (ab.), 521
- Aneurysm, abdominal, rupture of, 531
- Angina pectoris, 331
- Antacid action, influence of gastric function on (ab.), 250
- Antibiotic therapy
  - and alterations in bacterial flora (E.), 665
  - in Friedländer pneumonia, 179
- Antidiuretic hormone, secretion of, 135
- Antihyaluronidase, streptococcal, in rheumatic fever, arthritis and glomerulonephritis (ab.), 519
- Antirabies vaccine, fatal reaction to, 228
- Antithyroid therapy and exophthalmos, 581
- Aortic insufficiency with cardiac failure and recurrent abdominal pain, 507

### Arachnodactyly, 261

### Arteriosclerosis

- clinical implications of, 730
- diet and lipotropic agents in, 736
- of aorta, coronary and cerebral arteries (CPC), 507
- seminars on, 92, 209, 358, 468, 600, 730, 736

### Arteriosclerotic heart disease, 26

- Ascorbic acid, effect of, in depression of respiration of heart and kidney slices (ab.), 616

### Aspiration biopsy of kidney, 324

### Aspirin intoxication (ab.), 622

### Asthma treated with psychotherapy, 117

### Atherosclerosis, 92

- and blood lipids (E.), 271
- and lipid metabolism, 209
- lipoproteins in, 358
- predisposition to, and hypercholesteremia, 600

### Aureomycin

- and terramycin in amebic colitis (ab.), 620
- in congenital fibrocystic disease (ab.), 627

### Auricle, right, ball thrombus of, 653

### Auricular

- fibrillation in hyperthyroidism, 40
- flutter, 410

## Bacteria, effect of, on human leukocytes (ab.), 246

- Bacterial flora, normal, alterations in, during antibiotic therapy (E.), 665

### Ball thrombus of right auricle, 653

### Ballistocardiogram, evaluation of (ab.), 520

### Banthine

- and intact human colon, 67

## Banthine, effects of, on gastric secretion and peptic ulceration (ab.), 628

- Benemid, effect of, on serum penicillin concentrations (ab.), 627

### Benzodioxane tests, false negative, 524

### Bilious remitting fever (reprint), 546

### Biopsy, aspiration, of kidney, 324

### Blood

- flow in foot after hexamethonium (ab.), 519
- peripheral, evaluation of (ab.), 238
- lipids and atherosclerosis (E.), 271
- spectrophotometric oxygen determination in (ab.), 243

### Bone marrow

- absorption of substances injected into, 568
- influence of high oxygen tension on metabolism of (ab.), 241
- infusion of blood via (reprint), 571

### Book reviews

- Personality in Peptic Ulcer (Sullivan and McKell), 132
- Principles of Internal Medicine (Harrison), 270

## Calcinosis circumscripta, idiopathic, 267

### Canicola fever, 75

### Carbonic anhydrase inhibition on potassium excretion, 274

### Carcinoma

- of cervix, response of peripheral blood and bone marrow to x-ray therapy in (ab.), 243
- of prostate, 55
- of thyroid and I<sup>131</sup>, 44

### Cardiac

- and clinical catheterization, 31
- failure after aortic-pulmonary anastomosis, 403
- valvular prosthesis, plastic (ab.), 243

### Cardiospasm, adrenergic blockade for, 123

### Catheterization, clinical and cardiac, 31

### Cation exchange resin

- effect of, on electrolyte balance (ab.), 616
- in control of sodium balance (ab.), 517

### Cerebral

- circulation, oximetric technic in study of (ab.), 622
- vascular disease, cerebral functions in (ab.), 622

### Circulation, effect of isuprel on, 442

### Cirrhosis

- nodular, and carcinoma of liver (CPC), 387
- of liver, salt and water excretion in, 157
- serum and urine constituents in, 157

### Clinic on psychosomatic problems (Massachusetts General Hosp.)

- Case of asthma treated with psychotherapy, 117

### Psychotherapy of a psychosomatic illness, 381

### Clinico-pathologic conference (Washington Univ.)

- Aortic insufficiency with cardiac failure and recurrent abdominal pain, 507

### Fatal reaction to antirabies vaccine, 228

### Miliary pulmonary infiltration with hydrothorax and ascites, 757

### Recurrent hematemesis, abdominal pain and hepatic failure, 387

### Rheumatic heart disease with acute rheumatic fever, 109

### Cold pressor studies in hypertension and pheochromocytoma, 448

### Colon, intact, and banthine, 67

### Combined staff clinic (Columbia Univ.)

### Hypersplenism, 494



- Common cold (ab.), 522  
 Conference on therapy (Cornell Univ.)  
   Use of curare and curare-like agents, 749  
 Convalescent care (E.), 409  
 Cortisone  
   alteration of inflammatory response by (ab.), 248  
   and rheumatic carditis, 128  
   effects of, in bacterial infections (ab.), 252  
     on fluid and electrolyte distribution, 715  
   therapy in periarteritis nodosa, 630  
 Cranial arterial tone, fluctuations in (ab.), 249  
 Cytology, quantitative, exfoliative (ab.), 621
- D**  
 Defibrillation, electrical, of heart (ab.), 242  
 Detoni-Fanconi syndrome (ab.), 244  
 Dextran  
   effect of, on cardiac output and circulation (ab.), 618  
   for expansion of plasma volume (ab.), 238  
   gastrointestinal excretion of (ab.), 618  
   use of, in shock (ab.), 253  
 Diabetes, severe, intermediate insulins in control of (ab.), 516  
 Dicumarol-induced hemorrhage and vitamin K<sub>1</sub> oxide (ab.), 244  
 Diet and lipotropic agents  
   in arteriosclerosis, 736  
   influence of, on phospholipid turnover in plasma (ab.), 240  
 Digitoxin, intravenous, effect of, on fluid distribution (ab.), 237  
 Diphenhydramine in auricular fibrillation (ab.), 625  
 Diverticulosis of jejunum with hemorrhage, 398  
 Drugs, antispasmodic, and intact human colon, 67  
 Duodenal stump ulcer (ab.), 520  
 Duties of a physician (reprint), 551
- E**  
 Ectopic rhythms, pronestyl in, 431  
 Eisenmenger's complex, 31  
 Electrocardiogram  
   exercise, 26  
   extremity leads of, 3  
 Electroencephalographic abnormalities in liver disease (ab.), 619  
 Electrolyte  
   and water content of cardiac and skeletal muscle (ab.), 517  
   distribution, effect of cortisone and ACTH on, 715  
 Endocarditis, chronic, and pancarditis, rheumatic (CPC), 109  
 Ethyl chloride, effects of, on cardiac rhythm (ab.), 239  
 Exercise, effect of, on sodium excretion (ab.), 615  
 Exophthalmos and antithyroid therapy, 581
- F**  
 Fat tolerance in patients recovering from myocardial infarction (ab.), 516  
 Fatal reaction to antirabies vaccine, 228  
 First five years of American Journal of Medicine (E.), 1  
 Friedländer pneumonia, antibiotic therapy in, 179
- G**  
 Glomerular filtration rate and urinary excretion, variations in (ab.), 239  
 Glucose tolerance in glycosuria (ab.), 247  
 Gouty tophi, mobilization of, by uricosuric agents, 765
- H**  
 Heart  
   electric field of, and extremity leads of electrocardiogram, 3  
   failure, congestive, response to elevation of temperature and humidity in (ab.), 617  
   response of, to vagal stimulation (ab.), 250
- Hemochromatosis, 590  
 Hemorrhage and diverticulosis of jejunum, 398  
 Hemorrhagic disposition in certain families (reprint), 557  
 Hepatic disease, serum cholinesterase in (ab.), 627  
 Herpangina (ab.), 522  
 Hexamethonium, evaluation of (ab.), 242  
 Hormone therapy in carcinoma of prostate, 55  
 Horns of a dilemma (E.), 133  
 Human serum, reaction of, with quaternary ammonium salt (ab.), 518  
 Hyaluronidase, use of, in administration of aureomycin (ab.), 627  
 Hypercholesteremia with predisposition to atherosclerosis, 600  
 Hypersplenism, 494  
 Hypertension  
   canine, peripheral resistance in (ab.), 252  
   drug therapy of (ab.), 616  
   malignant, course of (ab.), 626  
   pharmacologic and cold pressor studies in, 448  
   pulmonary, 686  
 Hyperthyroidism  
   and myxedema, effect of, on cerebral blood flow and metabolism (ab.), 246  
   diagnosis of, with radioactive iodine (ab.), 516  
   masked, in cardiac patients, 40  
   methimazole (tapazole) in, 724  
 Hypoproteinemia, idiopathic, metabolic defect in (ab.), 628
- I**  
 Imidazoline as hypotensive agent (ab.), 617  
 Infarction of stomach, 645  
 Infusion of blood into circulation, 571  
 Insulin tolerance tests (ab.), 628  
 Iodine, radioactive, for thyroid carcinoma, 44  
 Iron  
   metabolism in hemochromatosis (ab.), 521  
   stores in man (ab.), 629  
 Irritable heart (reprint), 559  
 Isuprel, effect of, on circulation, 442
- J**  
 Jejunum, diverticulosis of, 398
- K**  
 Kidney, aspiration biopsy of, 324
- L**  
 Leads, extremity, of electrocardiogram, 3  
 Leukemia  
   granulocytic, clinical appearance of, 658  
   uptake of P<sup>32</sup> in DNA fraction in (ab.), 624  
 Lipid metabolism  
   and atherosclerosis, 209  
   inborn error of, 600  
 Lipoproteins in atherosclerosis, 358  
 Liver  
   cell necrosis, central, role of shock in, 170  
   physiology, studies in (ab.), 237  
 L-nor-epinephrine, effect of, on cerebral circulation (ab.), 250  
 Long follow-up, 572  
 Low potassium syndrome, 283
- M**  
 Marfan's syndrome, 261  
 Meningitis, aseptic, epidemic of, due to *Leptospira pomona* (ab.), 619  
 Methimazole (tapazole) in hyperthyroidism, 724  
 Methylandrostenedial (ab.), 629  
 Miliary pulmonary infiltration with hydrothorax and ascites, 757  
 Mononucleosis, infectious, 202  
 Myeloma, serum and urinary proteins in (ab.), 249

- Neoplasms**, metastatic, of liver and hepatic arteriography (ab.), 625
- Nodal rhythms**, experimental production of (ab.), 625
- Osteomalacia and renal tubular acidosis**, 302
- Oxygen therapy**, effects of, on cerebral functions (ab.), 249
- Para-amino-benzoic acid**, acetylation of, by liver (ab.), 247
- Paritol**, effects of, on prothrombin and Lee-White clotting times (ab.), 617
- Paritol-C**, a new anticoagulant (ab.), 520
- Pathology of atherosclerosis**, 92
- Penicillin "O"**, antibacterial properties of, 188
- Pennsylvania Hospital**  
during its first century, 540  
200th anniversary, 535
- Periarteritis nodosa**, 630
- Pharmacologic**  
basis of tests used in pheochromocytoma, 461  
studies in hypertension and pheochromocytoma, 448
- Pheochromocytoma**  
pharmacologic and cold pressor studies in, 448  
basis for tests in, 461  
studies on (ab.), 252  
with false negative benzodioxane tests, 524
- Pitressin tannate**, sodium and water metabolism studies during administration of (ab.), 518
- Pituitary**  
anterior, cellular changes in, during constriction of renal arteries or nephrectomy (ab.), 251  
posterior, influence of, on sodium and water excretion (ab.), 243
- Plasma**  
and interstitial fluid, measurement of (ab.), 629  
human, protein-lipid relationships in, 468, 480
- Polio-myelitis research**, recent trends in, 577
- Polycythemia vera**, pulmonary function studies in, 706
- Polymyxin**, oral, in chronic shigella infections (ab.), 626
- Polyradiculitis**, experimental (ab.), 523
- Potassium metabolism and acidification of urine**, 274
- Pressures in occluded hepatic venule and portal venous pressure** (ab.), 251
- Pressure-volume relationships in pulmonary circuit** (ab.), 241
- Primary amines and pressor substances in urine** (ab.), 240
- Pro fibrinolysin**, serum, activation of (ab.), 245
- Pronestyl** in ectopic rhythms, 431
- Prostate, carcinoma of**, 55
- Protein-bound iodine**, clinical use of (ab.), 619
- Protein-lipid relationships in human plasma**, 468, 480
- Pseudomonas aeruginosa**, infections due to (ab.), 254
- Psychiatry in medicine**, 588
- Psychotherapy**  
for asthma, 117  
of a psychosomatic illness, 381
- Pulmonary**  
blood volume and vital capacity (ab.), 240  
disease, types of, and impairment of alveolar-capillary diffusion, 667  
function studies in polycythemia vera, 706  
hypertension, 686
- Pyelonephritis**, recognition of, 312
- Recurrent hematemeses**, abdominal pain and hepatic failure, 387
- Regitine and benodaine**, adrenolytic blocking effects of, (ab.), 241
- Renal**  
clearances of penicillin, PSP and PAH (ab.), 517  
disease, chronic, electrolyte therapy in (ab.), 615  
excretion of digitoxin (ab.), 626  
of sodium solutions in Addison's disease (ab.), 238  
tubular acidosis and osteomalacia, 302  
mechanisms and low potassium syndrome, 283  
vascular shunts, problem of (ab.), 615  
Respiration of human cerebral cortex slices (ab.), 624  
Respiratory tract, localized amyloid deposition in, 770
- Rheumatic**  
carditis, histopathology of, 128  
disease, turbidimetric estimation of gamma globulin in (ab.), 518  
fever and rheumatic heart disease, 109  
heart disease with acute rheumatic fever, 109  
Rupture of abdominal aneurysm into gastrointestinal tract, 531
- Seminars on arteriosclerosis**, 92, 209, 358, 468, 600, 730, 736
- Serum antiproteolytic activity in disease** (ab.), 518
- Shock in central liver cell necrosis**, 170
- Sickle cell anemia**, antiserum against erythrocytes from (ab.), 620
- Splanchnic removal of bacteria from blood stream in leukopenia** (ab.), 623
- Stomach**, infarction of, 645
- Stress and ACTH**, effects of, on plasma amino acid nitrogen levels (ab.), 239
- Terramycin**, studies on (ab.), 620
- Tetralogy of Fallot**, 403
- Thecoma with feminization** (ab.), 246
- Thorotrast interference with removal of bacteria from splanchnic circulating blood** (ab.), 623
- Thyroid**  
carcinoma of, and  $I^{131}$ , 44  
gland  $I^{131}$  uptake, 255  
normal, and 6-n-propylthiouracil (ab.), 244
- Thyrotoxicosis factitia**, 255
- Torulosis**, visceral, without nervous system involvement, 658
- Trichinosis**, 639
- Triethylene melamine**  
in lymphomas and leukemias (ab.), 621  
therapy in leukemia (ab.), 253
- Tuberculosis**  
miliary, and pyelonephritis (CPC), 757  
study of, with rabbit eye (ab.), 523
- Uricosuric agents**, mobilization of gouty tophi with, 765
- Urinary sediments**, new stain for, 312
- Use of curare and curare-like agents**, 749
- Vaccine**, antirabies, fatal reaction to (CPC), 228
- Vectorcardiograms**  
analysis of (ab.), 618  
spatial, in right bundle branch block (ab.), 237
- Vitamin B<sub>12</sub>**, crystalline, in pernicious anemia (ab.), 521, 621
- Volume changes and mobilization of interstitial proteins by saline infusions** (ab.), 248
- Water**  
content of heart in anoxia (ab.), 245  
tolerance tests in cirrhosis of liver, 157
- West Nile and Ilheus virus infections in neoplastic disease** (ab.), 521

# The American Journal of Medicine

Editor ALEXANDER B. GUTMAN, M. D.

*Professor of Medicine*

COLUMBIA UNIVERSITY COLLEGE OF PHYSICIANS AND SURGEONS, NEW YORK  
DIRECTOR OF MEDICAL RESEARCH AND PHYSICIAN TO THE MOUNT SINAI HOSPITAL, NEW YORK

---

## ADVISORY BOARD

DAVID P. BARR, M.D., *Professor of Medicine, Cornell University Medical College, New York*; ARTHUR L. BLOOMFIELD, M.D., *Professor of Medicine, School of Medicine, Stanford University, San Francisco*; FRANCIS G. BLAKE, M.D., *Sterling Professor of Medicine, Yale University School of Medicine, New Haven*; EUGENE A. STEAD, JR., M.D., *Professor of Medicine, School of Medicine, Duke University, Durham*; JOSEPH T. WEARN, M.D., *Professor of Medicine, School of Medicine, Western Reserve University, Cleveland*.

## ASSOCIATE EDITORS

HERRMAN L. BLUMGART, M.D., *Boston*; HARRY GOLD, M.D., *New York*; A. McGEHEE HARVEY, M.D., *Baltimore*; GEORGE H. HOUCK, M.D., *Palo Alto*; CHESTER S. KEEFER, M.D., *Boston*; T. GRIER MILLER, M.D., *Philadelphia*; WALTER L. PALMER, M.D., *Chicago*; OSWALD H. ROBERTSON, M.D., *Stanford*; EPHRAIM SHORR, M.D., *New York*; GEORGE W. THORN, M.D., *Boston*; WILLIAM S. TILLET, M.D., *New York*; ROY H. TURNER, M.D., *New Orleans*; RUSSELL M. WILDER, M.D., *Bethesda, Md.*; M. M. WINTROBE, M.D., *Salt Lake City*; W. BARRY WOOD, M.D., *St. Louis*; JOHN B. YOUMANS, M.D., *Nashville*.

*Volume XI*  
JULY TO DECEMBER  
1951

THE AMERICAN JOURNAL OF MEDICINE, INC.

NEW YORK

MCMLI



# CONTENTS OF VOLUME XI

## ORIGINAL ARTICLES

The First Five Years . . . . .	<i>The Editor</i> . . . . .	1
A New Method of Equating and Presenting Bipolar and Unipolar Extremity Leads of the Electrocardiogram. Advantages Gained in Visualization of Their Common Relationship to the Electric Field of the Heart. . . . .	( <i>Lieut. John S. Graettinger</i> . . . . . <i>Lieut. John M. Packard</i> . . . . . <i>Capt. Ashton Graybiel</i> . . . . .)	3
Exercise Electrocardiogram. An Aid in the Diagnosis of Arteriosclerotic Heart Disease in Persons Exhibiting Abnormally Large Q <sub>3</sub> Waves . . . . .	( <i>Carl F. Shaffer</i> . . . . . <i>Don W. Chapman</i> . . . . .)	26
Clinical and Cardiac Catheterization Studies in Four Cases of Eisenmenger's Complex . . . . .	( <i>Richard S. Cosby</i> . . . . . <i>David C. Levinson</i> . . . . . <i>George C. Griffith</i> . . . . . <i>Willard J. Zinn</i> . . . . . <i>Sim P. Dimitroff</i> . . . . .)	31
Diagnosis of Masked Hyperthyroidism in Cardiac Patients with Auricular Fibrillation . . . . .	( <i>Mortimer M. Bortin</i> . . . . . <i>Solomon Silver</i> . . . . . <i>Stephen B. Yohalem</i> . . . . . <i>A. Stone Freedberg</i> . . . . .)	40
Treatment of Thyroid Carcinoma with Radioactive Iodine (I <sup>131</sup> ) . . . . .	( <i>Alvin L. Ureles</i> . . . . . <i>Mark F. Lesses</i> . . . . . <i>Samuel L. Gargill</i> . . . . .)	44
Management of Carcinoma of the Prostate. Study of Hormonal and Surgical Therapy in 100 Patients . . . . .	( <i>J. Hartwell Harrison</i> . . . . . <i>Eugene F. Poutasse</i> . . . . .)	55
Effects of Certain Antispasmodic Drugs on the Intact Human Colon, with Special Reference to Banthine ( $\beta$ -Diethylaminoethyl Xanthene-9-Carboxylate Methobromide) . . . . .	( <i>Fred Kern, Jr.</i> . . . . . <i>Thomas P. Almy</i> . . . . . <i>Natalie J. Stolk</i> . . . . .)	67
Canicola Fever. Review, with Report of Two New Cases . . . . .	<i>Bernard L. Rosenberg</i> . . . . .	75
Pathology of Atherosclerosis . . . . .	( <i>G. Lyman Duff</i> . . . . . <i>Gardner C. McMillan</i> . . . . .)	92
Rheumatic Heart Disease with Acute Rheumatic Fever. . . . .		109
A Case of Asthma Treated with Psychotherapy . . . . .		117
Treatment of Cardiospasm with Adrenergic Blockade . . . . .	( <i>Mark Nickerson</i> . . . . . <i>Lloyd S. Call</i> . . . . .)	123
Effect of Cortisone and ACTH on the Histopathology of Rheumatic Carditis. Report of a Necropsied Case. . . . .	( <i>David M. Spain</i> . . . . . <i>Daniel Roth</i> . . . . .)	128
The Horns of a Dilemma . . . . .	<i>John B. Youmans</i> . . . . .	133
Problem of Estimating the Rate of Secretion of Antidiuretic Hormone in Man . . . . .	<i>Henry D. Lauson</i> . . . . .	135
Studies of the Serum and Urine Constituents in Patients with Cirrhosis of the Liver during Water Tolerance Tests . . . . .	( <i>Elaine P. Ralli</i> . . . . . <i>Stephen H. Leslie</i> . . . . . <i>George H. Stueck, Jr.</i> . . . . . <i>Bertram Laken</i> . . . . .)	157
Role of Shock in the Production of Central Liver Cell Necrosis . . . . .	( <i>Max Ellenberg</i> . . . . . <i>Kermit E. Osserman</i> . . . . .)	170
Antibiotic Therapy of Friedländer Pneumonia . . . . .	( <i>William M. M. Kirby</i> . . . . . <i>Daniel H. Coleman</i> . . . . .)	179
Clinical and Pharmacologic Studies with Allylmercaptomethyl Penicillin (Penicillin "O") . . . . .	( <i>Charles V. Adair</i> . . . . . <i>William G. Woodin</i> . . . . . <i>Paul A. Bunn</i> . . . . .)	188

# Contents

iii

Allergy to Penicillin O and the Management of Penicillin Sensitivity . . . . .	<i>Sheppard Siegal . . . . .</i>	196
Infectious Mononucleosis. A Study of 210 Sporadic Cases . . . . .	<i>{ John E. Stevens . . . . . Edwin D. Bayrd . . . . . Frank J. Heck . . . . . }</i>	202
Lipid Metabolism and Atherosclerosis . . . . .	<i>R. Gordon Gould . . . . .</i>	209
Fatal Reaction to Antirabies Vaccine . . . . .		228
Southern Society for Clinical Research—Abstracts of Papers Presented at the Fifth Annual Meeting, New Orleans, Louisiana, January 27, 1951 . . . . .		237
Thyrotoxicosis Factitia. Measurement of the Thyroid Gland I <sup>131</sup> Uptake and Serum Protein-bound I <sup>131</sup> . . . . .	<i>{ David L. Chamovitz . . . . . Marvin H. Slesinger . . . . . A. Stone Freedberg . . . . . }</i>	255
Marfan's Syndrome (Arachnodactyly). . . . .	<i>George J. Hamwi . . . . .</i>	261
Idiopathic Calcinosis Circumscripta . . . . .	<i>{ Gerald I. Plitman . . . . . Benjamin R. Gendel . . . . . }</i>	267
Blood Lipids and Atherosclerosis . . . . .	<i>Herrman L. Blumgart . . . . .</i>	271
Relationship between Acidification of the Urine and Potassium Metabolism. Effect of Carbonic Anhydrase Inhibition on Potassium Excretion . . . . .	<i>{ Robert W. Berliner . . . . . Thomas J. Kennedy, Jr. . . . . Jack Orloff . . . . . }</i>	274
Low Potassium Syndrome Due to Defective Renal Tubular Mechanisms for Handling Potassium . . . . .	<i>{ David P. Earle . . . . . Sol Sherry . . . . . Ludwig W. Eichna . . . . . Neal J. Conan . . . . . }</i>	283
Renal Tubular Acidosis with Osteomalacia. Report of Three Cases . . . . .	<i>{ Kermit L. Pines . . . . . Gilbert H. Mudge . . . . . }</i>	302
Clinical Recognition of Pyelonephritis, with a New Stain for Urinary Sediments . . . . .	<i>{ Richard Sternheimer . . . . . Barney Malbin . . . . . }</i>	312
Aspiration Biopsy of the Kidney . . . . .	<i>{ Poul Iversen . . . . . Claus Brun . . . . . }</i>	324
Angina Pectoris. A Clinical and Pathologic Correlation. . . . .	<i>{ Paul M. Zoll . . . . . Stanford Wessler . . . . . Herrman L. Blumgart . . . . . Hardin B. Jones . . . . . John W. Gofman . . . . . Frank T. Lindgren . . . . . }</i>	331
Lipoproteins in Atherosclerosis . . . . .	<i>{ Thomas P. Lyon . . . . . Dean M. Graham . . . . . Beverly Strisower . . . . . Alex V. Nichols . . . . . }</i>	358
Psychotherapy of a Psychosomatic Illness: Essential Hypertension . . . . .		381
Recurrent Hematemesis, Abdominal Pain and Hepatic Failure . . . . .		387
Diverticulosis of Jejunum with Hemorrhage . . . . .	<i>{ Edmund H. Berger . . . . . Clarence W. Brunkow . . . . . Clarence W. Smith . . . . . }</i>	398
Cardiac Failure after Aortic-pulmonary Anastomosis in Tetralogy of Fallot . . . . .	<i>{ John A. Sheedy . . . . . Werner K. Gottstein . . . . . George K. Fenn . . . . . }</i>	403

Convalescent Care . . . . .	Herman Zazeela . . . . .	409
	(Myron Prinzmetal . . . . .)	
	(Eliot Corday . . . . .)	
	(Robert W. Oblath . . . . .)	
	(H. E. Kruger . . . . .)	
	(I. C. Brill . . . . .)	
Auricular Flutter . . . . .	(Joshua Fields . . . . .)	410
	(S. Rexford Kennamer . . . . .)	
	(John A. Osborne . . . . .)	
	(L. Allen Smith . . . . .)	
	(Alvin L. Sellers . . . . .)	
	(Walter Flieg . . . . .)	
	(Eileen Finston . . . . .)	
Use of Pronestyl in the Treatment of Ectopic Rhythms.	(Keehn Berry . . . . .)	
Treatment of Ninety-eight Episodes in Seventy-eight	(E. Lee Garlett . . . . .)	431
Patients . . . . .	(Samuel Bellet . . . . .)	
	(William I. Gefter . . . . .)	
Effect of Isuprel (Isopropylpinephrine) on Circulation of	(Joseph Kaufman . . . . .)	
Normal Man . . . . .	(Arnold Iglauer . . . . .)	442
	(Gisela K. Herwitz . . . . .)	
Re-evaluation of the Reliability of Pharmacologic and	(James A. Evans . . . . .)	
Cold Pressor Studies in Hypertension and Pheochromocytoma . . . . .	(Hyman J. Rubitsky . . . . .)	448
	(Carl C. Bartels . . . . .)	
	(Elmer C. Bartels . . . . .)	
Pharmacologic Basis of Various Tests Used in the Diag-	(George Entwisle . . . . .)	
nosis of Pheochromocytoma . . . . .	(Clement A. Stone . . . . .)	461
	(Earl R. Loew . . . . .)	
Protein-lipid Relationships in Human Plasma		
I. In Normal Individuals . . . . .	(Ella M. Russ . . . . .)	
	(Howard A. Eder . . . . .)	468
	(David P. Barr . . . . .)	
II. In Atherosclerosis and Related Conditions . . . . .	(David P. Barr . . . . .)	
	(Ella M. Russ . . . . .)	480
	(Howard A. Eder . . . . .)	
Hypersplenism . . . . .		494
Aortic Insufficiency with Cardiac Failure and Recurrent		
Abdominal Pain . . . . .		507
American Federation for Clinical Research—Abstracts of		
Papers Presented at the Eastern Sectional Meeting in		
Washington, D.C., December 9, 1950 . . . . .		516
Pheochromocytoma with False Negative Benzodioxane		
Tests . . . . .	Robert E. Mason . . . . .	524
Rupture of Abdominal Aneurysm into the Gastrointes-	(Eli Antzis . . . . .)	
tinal Tract . . . . .	(James Dunn . . . . .)	531
	(Anthony J. Schilero . . . . .)	
Pennsylvania Hospital—200th Anniversary . . . . .	Garfield G. Duncan . . . . .	535
Days at the Pennsylvania Hospital during Its First Cen-		
tury . . . . .	Edward B. Krumbhaar . . . . .	540



# Contents

v

An Account of the Bilious Remitting Fever, as It Appeared in Philadelphia, in the Summer and Autumn of the Year 1780 . . . . .	<i>Benjamin Rush</i> . . . . .	546
Observations on the Duties of a Physician, and the Methods of Improving Medicine. Accommodated to the Present State of Society and Manners in the United States . . . . .	<i>Benjamin Rush</i> . . . . .	551
An Account of an Hemorrhagic Disposition Existing in Certain Families . . . . .	<i>John C. Otto</i> . . . . .	557
On Irritable Heart. A Clinical Study of a Form of Functional Cardiac Disorder and Its Consequences. . . . .	<i>J. M. Da Costa</i> . . . . .	559
Rapid Absorption of Substances Injected into the Bone Marrow . . . . .	<i>L. M. Tocantins</i> . . . . .	568
Infusion of Blood and Other Fluids into the Circulation via the Bone Marrow. . . . .	{ <i>L. M. Tocantins</i> . . . . . <i>J. F. O'Neill</i> . . . . . }	571
The Long Follow-up: A Compensation of the Aging Physician . . . . .	<i>O. H. Perry Pepper</i> . . . . .	572
Recent Trends in Poliomyelitis Research . . . . .	<i>John R. Paul</i> . . . . .	577
Exophthalmos in the Light of Current Anti-thyroid Therapy . . . . .	<i>Henry M. Thomas, Jr.</i> . . . .	581
The Place of Psychiatry in Medicine . . . . .	<i>E. D. Bond</i> . . . . .	588
Hemochromatosis . . . . .	{ <i>Alexander Marble</i> . . . . . <i>C. Cabell Bailey</i> . . . . . }	590
Hypercholesteremia with Predisposition to Atherosclerosis. An Inborn Error of Lipid Metabolism . . . . .	<i>David Adlersberg</i> . . . . .	600
American Federation for Clinical Research—Abstracts of Papers Presented at the Southern Sectional Meeting in New Orleans, Louisiana, January 26, 1951. . . . .		615
American Federation for Clinical Research—Abstracts of Papers Presented at the Western Sectional Meeting in Seattle, Washington, January 25, 1951 . . . . .		624
Periarthritis Nodosa. Report of a Case Treated with ACTH and Cortisone. . . . .	{ <i>William L. Mundy</i> . . . . . <i>William G. Walker, Jr.</i> . . . . <i>Hylan A. Bickerman</i> . . . . . <i>Gustav J. Beck</i> . . . . . }	630
Trichinosis. Case Report with Observations of the Effect of Adrenocorticotrophic Hormone . . . . .	{ <i>William Mabon Davis</i> . . . . . <i>Harry Most</i> . . . . . }	639
Infarction of the Stomach. Report of Three Cases of Total Gastric Infarction and One Case of Partial Infarction. . . . .	<i>Eli B. Cohen</i> . . . . .	645
Ball Thrombus of the Right Auricle . . . . .	<i>Robert S. Radding</i> . . . . .	653
Disseminated Visceral Torulosis without Nervous System Involvement. With Clinical Appearance of Granulocytic Leukemia . . . . .	{ <i>Samuel Zelman</i> . . . . . <i>Robert H. O'Neil</i> . . . . . <i>Alfred Plaut</i> . . . . . }	658
Alterations in Normal Bacterial Flora of Man and Secondary Infections during Antibiotic Therapy . . . . .	<i>Chester S. Keefer</i> . . . . .	665
Clinical and Physiologic Features of Some Types of Pulmonary Diseases with Impairment of Alveolar-capillary Diffusion. The Syndrome of "Alveolar-capillary Block"	{ <i>Robert Austrian</i> . . . . . <i>John H. McClement</i> . . . . . <i>Attilio D. Renzetti, Jr.</i> . . . . <i>Kenneth W. Donald</i> . . . . . <i>Richard L. Riley</i> . . . . . <i>André Cournand</i> . . . . . }	667

Primary Pulmonary Hypertension. I. Clinical and Hemodynamic Study . . . . .	{ David T. Dresdale . . . . . Martin Schultz . . . . . Robert J. Michtom . . . . . }	686
Pulmonary Function Studies in Polycythemia Vera. Results in Five Probable Cases . . . . .	{ Walter Newman . . . . . James A. Feltman . . . . . Blanche Devlin . . . . . }	706
Effect of Cortisone and ACTH on Fluid and Electrolyte Distribution in Man . . . . .	{ Marvin F. Levitt . . . . . Mortimer E. Bader . . . . . }	715
Experience with Methimazole (Tapazole) in the Treatment of Hyperthyroidism. A Report of Thirty-five Cases . . . . .	{ Bernard L. Hallman . . . . . Philip K. Bondy . . . . . }	724
Arteriosclerosis—Some Clinical Implications . . . . .	William Dock . . . . .	730
Diet and Lipotropic Agents in Arteriosclerosis . . . . .	Jack D. Davidson . . . . .	736
Use of Curare and Curare-like Agents. . . . .		749
Miliary Pulmonary Infiltration with Hydrothorax and Ascites . . . . .		757
Mobilization of Gouty Tophi by Protracted Use of Uricosuric Agents . . . . .	{ T. F. Yü . . . . . Alexander B. Gutman . . . . . Aaron Schottenfeld . . . . . }	765
Localized Amyloid Deposition in the Lower Respiratory Tract . . . . .	{ Leon M. Arnold . . . . . John G. Gruhn . . . . . A. David Etess . . . . . }	770

COPYRIGHT, 1951

BY THE AMERICAN JOURNAL OF MEDICINE, INC.

*All Rights Reserved*



## Prelude to asthma?

*not necessarily...*

Tedral, taken at first sign of attack, often forestalls severe symptoms.

*in 15 minutes...* Tedral brings symptomatic relief with a definite increase in vital capacity. Breathing becomes easier as Tedral relaxes smooth muscle, reduces tissue edema, provides mild sedation.

*for 4 full hours...* Tedral maintains more normal respiration for a sustained period—not just a momentary pause in the attack.

*Prompt and prolonged relief* with Tedral can be initiated any time, day or night, whenever needed without fear of incapacitating side effects.

*Tedral provides:*

theophylline .....	2 gr.
ephedrine .....	$\frac{3}{8}$ gr.
phenobarbital .....	$\frac{1}{8}$ gr.

*in boxes of 24, 120 and 1000 tablets*

# Tedral

**CHILCOTT**  
*Laboratories*


DIVISION OF The Maltine Company MORRIS PLAINS, NEW JERSEY



# AUREOMYCIN

Hydrochloride Crystalline

*Effective against many bacterial and  
rickettsial infections, as well as certain  
protozoal and large viral diseases.*



The Geriatriist looks always for a treatment which shall act effectively to curb infection, without unduly upsetting normal metabolic processes and immunologic responses. Aureomycin provides a maximum anti-infectious effect with a minimum of disturbance. Infection in the elderly is more apt to be subacute, or chronic, than acute; and of mixed rather than pure type. Under such conditions, the oral effectiveness and broad activity of aureomycin make it of exceptional value.


*Capsules: 50 mg.—Bottles of 25 and 100, 250 mg.— Bottles of 16 and 100.*

*Ophthalmic: Vials of 25 mg. with dropper; solution prepared by adding 5 cc. of distilled water.*

LEDERLE LABORATORIES DIVISION

*AMERICAN Cyanamid COMPANY*

30 Rockefeller Plaza, New York 20, N. Y.



IN THE **COSTLY** CARE OF CHRONIC OSTEOMYELITIS

**Tryptar**\*

#### OTHER INDICATIONS

Amputation Stumps  
Burns  
Carcinomatous Ulcers  
Chest Surgery  
Decubitus Ulcers  
Diabetic Gangrene  
Empyema  
Infected Compounded  
Fractures  
Soft Tissue Abscesses,  
Sinuses, Fistulae  
Subcutaneous Hematoma  
Varicose Ulcers

In chronic osteomyelitis, therapy that hastens recovery is an economy reflected in many ways. Tryptar is dramatic in reducing recovery time from years to months or even weeks.

Tryptar, selectively, causes dissolution of necrotic tissue and removes debris without injury to normal tissue. With topical administration, Tryptar is non-toxic and completely non-antigenic.

Tryptar in the treatment of osteomyelitis may be applied in the form of dry powder, in wet dressings, in gelatin capsules or by irrigation, depending upon the location of the lesion.

Tryptar is supplied as a two vial preparation: one 30 cc. vial contains 250,000 Armour Units (250 mg. of Tryptic Activity) of highly purified crystalline trypsin; the companion 30 cc. vial contains 25 cc. of Tryptar Diluent (Sorensen's Phosphate Buffer Solution) pH 7.1. Plus plastic adapter for use with powder blower.

\*THE ARMOUR LABORATORIES BRAND OF PURIFIED CRYSTALLINE TRYPSIN



**THE ARMOUR LABORATORIES • CHICAGO 11, ILLINOIS**

*—world-wide dependability—*

PHYSIOLOGIC THERAPEUTICS THROUGH BIORESEARCH

Highly effective in an unusually wide range  
of common skin disorders

# Pragmatar

the outstanding tar-sulfur-salicylic acid ointment



## *In seborrheic dermatitis*

Pruritic axillary seborrheic dermatitis of 6 years standing, April 5, 1951. Treatment over the years with various medicaments had failed.

At termination of therapy with Pragmatar, May 24, 1951. (From the first day of treatment the itching had stopped. By the end of one week there was a marked clinical improvement in the lesion.)

Formula: Cetyl alcohol-coal tar distillate, 4%; near-colloidal sulfur, 3%; salicylic acid, 3%—incorporated in a special washable base.

'Pragmatar' T.M. Reg. U.S. Pat. Off.

*Smith, Kline & French  
Laboratories, Philadelphia*





# *Only Terramycin* *in Oral Drops* offers *all* these advantages

**1*****convenience of a liquid concentrate***

Crystalline Terramycin Hydrochloride Oral Drops provide 200 mg. per cc., 50 mg. in each 9 drops—the only broad-spectrum antibiotic available as a liquid concentrate affording optimal convenience and flexibility in dosage schedules.

**2*****miscibility with foods and fluids***

Terramycin Oral Drops are miscible with most foods, milk and fruit juices; can be taken "as is" or mixed. Potent oral drops offer rapid broad-spectrum antibiotic activity in a form permitting the utmost simplicity in the therapeutic regimen.

**3*****pure crystalline compound—well tolerated***

Terramycin Oral Drops are prepared from pure crystalline material, free of impurities which may contribute to adverse reactions.


***supplied:***

2.0 Gm. with 10 cc. of diluent,  
and specially calibrated dropper.

ANTIBIOTIC DIVISION



CHAS. PFIZER &amp; CO., INC., Brooklyn 6, N. Y.



*This is what happens to spasm  
shortly following the administration  
of that excellent combination  
of natural belladonna alkaloids  
with minimal phenobarbital\*...*



**Donnatal®**

is a product of A. H. ROBINS CO., INC.


RICHMOND 20, VA.

Ethical Pharmaceuticals of Merit since 1878

\*In each tablet, capsule,  
or 5 cc. (100.) of Elixir:

Hyoscyamine Sulfate ..... 0.1037 mg.  
Atropine Sulfate ..... 0.0194 mg.  
Hyoscine Hydrobromide.. 0.0065 mg.  
Phenobarbital, 1/4 gr..... 16.2 mg.

# NEO-IOPAX



## NEO-IOPAX® (Sodium Iodomethamate U.S.P.)

A safe urographic  
contrast medium  
remarkably free  
from hazard for  
the patient.

## Pictures without penalties

NEO-IOPAX urograms and PRIODAX cholecystograms  
give definitive information for diagnosing certain pathologic  
conditions of the urinary and biliary tracts, respectively—without penalty.



# PRIODAX



## PRIODAX® (Iodoliphoic Acid U.S.P.)


A well tolerated  
cholecystographic  
agent, conveniently  
administered,  
pleasant to take.

*Schering* CORPORATION  
BLOOMFIELD, N. J.



# NEW!

*an oral therapeutic hematinic*



**IN RECOMMENDED DOSAGE** — 1 capsule t.i.d.  
— Rubragran supplies *truly therapeutic amounts* of four fundamental factors in normal red blood cell production.

<b>RUBRAGRAN FORMULA</b>	<b>1 capsule supplies</b>	<b>1 capsule t.i.d. supplies</b>
Vitamin B <sub>12</sub>	25 mcgm.	75 mcgm.
Folic Acid	1.67 mg.	5 mg.
Ascorbic Acid	100 mg.	300 mg.
Iron (elemental)	67 mg.	200 mg.
(as ferrous sulfate exsic.)	230 mg.	690 mg.
Desiccated Liver	100 mg.	300 mg.

*Bottles of 100 and 500*

Because of the serious nature of the disease, true Addisonian pernicious anemia should not be treated with oral preparations. Parenteral Rubramin or Liver Injection should be used.

'RUBRAGRAN' IS A TRADEMARK OF E. R. SQUIBB & SONS

## SQUIBB



**Ciba  
Presents**

**A New Advance  
in Sulfonamide Safety ...**

**ELKOSIN®**

BRAND OF SULFADIMETINE

Scored 0.5 Gm. tablets.  
Bottles of 100 and 1000.

- Remarkably low incidence of side effects—less than 5%
- Lowest acetylation yet reported—less than 10% in blood
- New improved solubility
- Renal complications rare—alkalis not needed
- High, sustained blood levels

WIDE ANTIBACTERIAL SPECTRUM

**Ciba** PHARMACEUTICAL PRODUCTS, INC., SUMMIT, N. J.

2-1890M



... In 28 Cases of  
**HERPES ZOSTER**  
 at Bethesda Naval  
 Hospital\*

William C. Marsh, Commander (MC) U.S.N., in a currently published paper<sup>1</sup>, "Treatment of Herpes Zoster with Protamide," now available to physicians as a reprint, presents these findings:

"Thirty-one cases of herpes zoster were treated with Protamide. Good to excellent results were obtained in twenty-eight. In those failing to respond other factors besides age of patient may have been involved.

"No controls were in our study as thousands of intramuscular injections of other drugs given to patients with herpes zoster in the past, with no appreciable benefit, would adequately serve as a control.

"Pain, not merely the discomfort or itching, was the indication for treatment. Protamide (1.3 cc, the contents of one ampul) was given daily intramuscularly.

No other local or systemic medications were given.

"The relief of pain was superior to that obtained when using either pituitrin, thiamine chloride, autohemotherapy, sodium iodide, or high voltage Roentgen therapy.


"The advantages of Protamide are the simplicity and absence of pain in administration, lack of reactions, and apparent safety.

"Costello<sup>2</sup> found that Protamide was effective in the relief of the posterior root pain of tabes dorsalis."


\* U. S. Naval Hospital, National Naval Medical Center, Bethesda, Maryland.

1. U.S. Armed Forces Med. Journal, September, 1950.

2. Costello, R. T. New treatment for "lightning pains" of tabes dorsalis, Urol. and Cutan. Rev. 51: 260-263, May, 1947.



DOCTOR: Your prescription blank marked "Protamide" will bring you a compilation of papers together with folder—"FOR PROMPT RELIEF OF POSTERIOR NERVE ROOT PAIN." Send for them today.





improves the


**whole**

patient in

**hypertension**

Each PERTENAL tablet contains:  
 Veratrum Viride..... 100 mg. (1½ gr.)  
 Homatropine Methyl Bromide. 2.5 mg. (1/25 gr.)  
 Mannitol Hexanitrate ..... 30 mg. (½ gr.)  
 Phenobarbital ..... 15 mg. (¼ gr.)

**PERTENAL samples**  
on request



# PERTENAL®

*controls pressure and symptoms that send pressure soaring*

**more normal pressure**

*around the clock with two potent vasodilators.*

**eases stress on heart**

*by decreasing peripheral resistance.*

**relieves gastro-intestinal spasm**

*pain and other discomforts which aggravate pressure.*

**relaxes patient**

*mentally, physically; allays tension, worry, restlessness, insomnia.*

**from the first day of PERTENAL**

*therapy the whole patient is more comfortable, happier; able to live a more normal, perhaps longer life.*

CROOKES LABORATORIES, INC.



305 EAST 45 ST., NEW YORK, N. Y.

*In Corpus Luteum  
Hormone Deficiencies...*

## PROGESTONE HYPOGLOSSALS



10 Mgs.  
per tablet  
Bottles of  
50, 100, 1,000

Descriptive Literature and Specimens Available

*Progesterone in  
tablets specially  
prepared for  
sublingual or buccal  
administration  
and eminently  
suitable for oral  
(swallowed) use*

"There can be no doubt that progesterone sublingually is absorbed and utilized..."—Greenblatt, R. B.: J. Clin. Endocrinol. 4:7, 1944.

"The observation of greatest importance in this investigation is that the results indicate that progesterone itself is active when administered by the oral (swallowed) route. Furthermore, progesterone itself appears to be more active orally than anhydroxyprogesterone." — Bickers, W.: J. Clin. Endocrinol. 9:8, 1949.

**G. W. Catrick Co.** NEWARK 1, NEW JERSEY



## All Children Can Benefit from this Protective Hot Drink at Breakfast

In its widely distributed leaflet No. 268, "Eat a Good Breakfast," the U. S. Dept. of Agriculture states: "Summer or winter, there's something hot, as a rule, in a good breakfast.... Something hot is cheering and tones up the whole digestive route."



The problem of encouraging children to eat an adequately protective breakfast finds easier solution when Ovaltine in hot milk is recommended as a breakfast beverage. Many children clamor for a hot drink at the morning meal, and hot Ovaltine is the right kind of drink to recommend.

A cup of hot Ovaltine makes an excellent contribution of virtually all essential nutrients, adding substantially to the nutritional start for the day. It also serves in a gustatory capacity by enhancing the appeal of breakfast and making other foods more inviting.

The nutrient contribution made by a cup of Ovaltine is apparent from the table below. Note the wealth of essentials added to the nutritional intake by making the simple recommendation of adding a cup of hot Ovaltine to the child's breakfast.

THE WANDER COMPANY, 360 N. MICHIGAN AVE., CHICAGO 1, ILLINOIS

### Ovaltine

Here are the nutrients that a cupful of hot Ovaltine, made of ½ oz. of Ovaltine and 8 fl. oz. of whole milk,\* provides:

PROTEIN. . . . .	10.5 Gm.	IRON . . . . .	4 mg.	NIACIN. . . . .	2.3 mg.
FAT . . . . .	10.5 Gm.	COPPER . . . . .	0.2 mg.	VITAMIN C. . . . .	10 mg.
CARBOHYDRATE . . . . .	22 Gm.	VITAMIN A . . . . .	1000 I.U.	VITAMIN D. . . . .	140 I.U.
CALCIUM . . . . .	370 mg.	VITAMIN B <sub>1</sub> . . . . .	0.39 mg.	CALORIES. . . . .	225
PHOSPHORUS . . . . .	315 mg.	RIBOFLAVIN . . . . .	0.7 mg.		

\*Based on average reported values for milk.

## Up-to-Date Antitussive



*Sedative-antitussive effects of Methajade control sleep-robbing paroxysmal cough—allow patients to get needed rest.*

# SUBDUES VIOLENT COUGHING

METHAJADE® spares your patients the needless violence of exhausting, "unproductive" cough—without depriving them of the benefits of normal expectoration.

Because it contains *methadone*, the synthetic analgesic-antitussive, METHAJADE is capable of suppressing the irritability of the cough reflex without blocking it entirely.

Because it contains potassium citrate and diluted phosphoric acid, METHAJADE promotes the liquefaction and loosening of mucus, facilitating the expulsion of bronchial exudate and debris.

Because it relaxes smooth muscle, including that in the tracheobronchial tree, METHAJADE is effective in relieving the bronchial spasm often associated with coughs.

Together, these actions tend to *decrease* the frequency and *increase* the efficiency of coughing, and to relieve bronchial spasm. METHAJADE is therefore well qualified as a practical aid in achieving the modern objectives of cough control:

- (1) to preserve the natural physiologic benefits of the cough—
- (2) to reserve the act of coughing for "productive" expectoration.

**Composition**—METHAJADE is a sugar-free antitussive with a delicious, fresh, lime flavor. Each 30 cc. (1 fl. oz.) contains:

Methadone hydrochloride* (d,l-6-dimethylamino-4,4-diphenyl-3-heptanone hydrochloride)	10 mg.
WARNING: may be habit forming.	
Propadrine®, phenylpropanolamine hydrochloride	0.12 Gm.
Potassium citrate	1.2 Gm.
Diluted phosphoric acid	4.5 cc.
Alcohol 5%	

**Average Dose for Adults:** 1 to 2 teaspoonfuls every three or four hours.

#### CHILDREN:

(note: METHAJADE should not be administered to children under 2 years of age. In children 2 years of age or older, METHAJADE should be used only in cases of severe, intractable cough.)

Two years:  $\frac{1}{4}$  teaspoonful not more often than every four hours.

Five years:  $\frac{1}{2}$  teaspoonful not more often than every four hours.

Ten years: 1 teaspoonful not more often than every four hours.

**Packaging**—METHAJADE is supplied in pint Spasaver® bottles and gallon bottles.  
Sharp & Dohme, Philadelphia 1, Pa.

\*The analgesic potency of methadone hydrochloride is—weight for weight—approximately equivalent to that of morphine. It controls cough as well as, or better than, codeine. Methadone hydrochloride is subject to the provisions of the Harrison Narcotic Act.

# METHAJADE®

Antitussive



combined attack

A therapeutic attack on bacteria is more effective when several agents having harmonious antibacterial actions are employed simultaneously.

For a simple, effective strategy against susceptible organisms:

## R<sub>x</sub> Biosulfa\*

Bottles of 50 and 500 tablets.

Each tablet contains:

Penicillin G Potassium, Crystalline ..	100,000 units
Sulfadiazine .....	0.167 Gm.
Sulfamerazine .....	0.167 Gm.
Sulfamethazine .....	0.167 Gm.

\*Trademark, Reg. U. S. Pat. Off.

**Upjohn**

*Medicine... Produced with care... Designed for health*

THE UPJOHN COMPANY, KALAMAZOO, MICHIGAN

new

Schering ANNOUNCES

# DORMISON\*

*new, non-barbiturate hypnotic*

*for safe, sound sleep*

*without drug hangover*

*free from habit-forming properties  
of the barbiturates*

*safe*

free from habit-forming or addiction properties  
of barbiturates; rapidly metabolized; no cumulative action;  
no toxic effects on prolonged use

*acts gently and quickly in insomnia*

mild hypnotic action quickly induces restful sleep

*no prolonged suppressive effect*


action subsides after a few hours; patient continues to sleep naturally

*no drug hangover*

patient awakens refreshed with no "drugged" feeling

DORMISON is a substance new to pharmacology, completely different from barbiturates and other hypnotics. It contains only carbon, hydrogen and oxygen. It has no nitrogen, bromine, urea residues, sulfone groups or chemical configurations present in depressant drugs now in use.

The usual dose of DORMISON (methylparafynol†) is one or two capsules, taken just before the patient is ready for sleep. DORMISON's wide margin of safety allows liberal adjustment of dosage until the desired effect is obtained. DORMISON is supplied as 250 mg. soft gelatin capsules in bottles of 100.



Schering CORPORATION  
BLOOMFIELD, N. J.



\*T.M. †U.S. Pat. Pending

DORMISON

COMING  
1952



## CONGENITAL HEART DISEASE

Classification of Congenital Heart Disease.....RICHARD T. BING, Baltimore, Md.

Physiologic Methods of Study of Congenital Heart

Disease .....AARON HIMMELSTEIN AND ANDRE  
COURNAND, New York

Angiocardiography in Congenital Heart Disease.....ISRAEL STEINBERG AND CHARLES  
DOTTER, New York

Tetralogy of Fallot.....WILLIS J. POTTS, Chicago, Ill.

Patent Ductus Arteriosus.....ROBERT E. GROSS, Boston, Mass.

Pulmonary Stenosis .....R. C. BROCK, London, England

## A SYMPOSIUM

MULTIPLE SCLEROSIS and DEMYELINATING DISEASES—May 1952

*Guest Editor*, GEORGE A. SCHUMACHER, M.D., Associate Professor of Clinical Medicine,  
Cornell University Medical College, New York

*Subscription*: U.S.A. \$12: Canada and South America \$14: Abroad \$15

THE AMERICAN JOURNAL OF MEDICINE

49 WEST 45TH STREET, NEW YORK 36, N. Y.



# ATHEROSCLEROSIS

*Too Much Cholesterol or  
Too Little Phospholipids?*

**Plasma Ratio:**

PHOSPHOLIPID	mg.	%
CHOLESTEROL	mg.	%

The ratio is usually less than 1.0 in atherosclerosis, but in good health is 1.0 or more. 1 and 2

## INCREASE PHOSPHOLIPIDS

The new isotope technique<sup>3</sup> has demonstrated that:

- (a) choline deficiency occurs in man;
- (b) phospholipid production is increased in deficiency states by choline;
- (c) individuals differ in choline needs;

(Inositol also enters into synthesis of certain phospholipids.<sup>4</sup>)

## DECREASE CHOLESTEROL

Although cholesterol plasma levels may not be affected by dietary restrictions, the physical state of the cholesterol may be affected. Thus, "giant cholesterol molecules" associated with atherosclerosis may be eliminated by dietary control.<sup>5</sup>

## RATIONAL MEASURES

In the light of these recent studies, adequate lipotropic therapy and cholesterol-restricted diets are rational measures in:

**ATHEROSCLEROSIS  
CORONARY ARTERY DISEASES  
DIABETES**

**And Other Unbalanced Conditions  
of Disturbed Cholesterol Metabolism**

## GERICAPS

Supply Therapeutic Dosage of Lipotropics (700 mg.) with Vitamin A and B Complex factors to help compensate for deficiencies of low-cholesterol diets with Rutin and Vitamin C to prevent the retinopathy often associated with above diseases.

For Literature and Clinical Trial Package please write "Gericaps Literature" on post card, sign and send to us.



## BIBLIOGRAPHY

<sup>1</sup>Morrison L.M.: Tests for Atherosclerosis. Proc. AM. Soc. for Study of Atherosclerosis P 478 (1950)

<sup>2</sup>Gertler, M.M.; Stanley M.; Blund, E.F.; Age, Serum, Cholesterol and Coronary Artery Disease. Circulation 2:517 (1950)


<sup>3</sup>Science 109:613 (June 17) 1949

<sup>4</sup>Biochem. J. 40:494 (1946)

<sup>5</sup>Gofman, J.W.; et al.; Science. 111:166 (1950)

## SHERMAN LABORATORIES

Detroit 15, Michigan  
Windsor, Ont. Los Angeles 5, Cal.



FOR  
**Christmas** . THE IDEAL GIFT FOR  
YOUR COLLEAGUE OR DOCTOR FRIEND

An up-to-date medical journal any doctor will take pride in having in his library—beautifully printed and illustrated—a publication that will provide both pleasure and inspiration.

The American Journal of Medicine  
49 West 45th St. New York 36


Yearly Subscription \$10.00 U.S.A.

Canada and Pan-American Union \$12.00

Foreign \$15.00



**PONDETS**  
contain  
**BOTH PENICILLIN**  
and  
**BACITRACIN**



In a delicious hard  
candy base that  
completely masks  
antibiotic taste.

**ADVANTAGES:**

More potent  
antibiotic action

Penicillin and bacitracin exhibit true  
synergism.<sup>1,2</sup>

Wider antibacterial  
Spectrum

Organisms with little or "borderline"  
sensitivity to either antibiotic alone, are  
often readily susceptible to this com-  
bination.

Effective Oral  
Levels

Lasting at least one-half hour in most  
patients.

1. Eagle, H., and Fleischman, R.: Proc. Soc. Exper. Biol.  
& Med. 68:415, 1948
2. Bachman, M. C.: J. Clin. Invest. 28:864, 1949

In each troche: 20,000 units Crystalline Potassium  
Penicillin-G, and 50 units Bacitracin.

**PONDETS®**

PENICILLIN-BACITRACIN TROCHES WYETH

SUPPLIED: Vacuum-packed tins of 48 troches.

*Wyeth* INCORPORATED, PHILADELPHIA 2, PA.





# MERCUHYDRIN<sup>®</sup>

SODIUM

*effective, well tolerated ...*

*intramuscularly, intravenously and subcutaneously*

"Reaser and Burch, using radioactive isotopes, have confirmed earlier evidence that mercurial diuretics greatly increase the urinary loss of sodium. The sodium diuresis precedes water diuresis by two to four hours, and sodium excretion per day may be increased sevenfold while water excretion is merely doubled."

Editorial: J.A.M.A. 135:576,  
Nov. 1, 1947.

"... The concentration of sodium in the urine was increased nearly two and one-half times by the injection of the mercurial diuretic, while the average total excretion of sodium in 24 hours was increased more than four times by MERCUHYDRIN injections."

Griggs, D. E., and Johns, V. J.:  
California Med. 69:133, Aug. 1948.

**MERCUHYDRIN** Sodium (brand of meralluride sodium) is available in 1 cc. and 2 cc. ampuls.

*L*akeside  
laboratories, INC., MILWAUKEE 1, WISCONSIN

